

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 June 2003 (26.06.2003)

PCT

(10) International Publication Number
WO 03/051868 A1

(51) International Patent Classification⁷: C07D 401/14,
401/04, 491/10, 405/14, 413/14, A61K 31/4545, A61P
13/10

and Development, Ramsgate Road, Sandwich, Kent CT13
9NJ (GB).

(21) International Application Number: PCT/IB02/05234

(74) Agents: HAYLES, James, R. et al.; Pfizer Limited, Rams-
gate Road, Sandwich, Kent CT13 9NJ (GB).

(22) International Filing Date: 6 December 2002 (06.12.2002)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130261.1 18 December 2001 (18.12.2001) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Burasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for GB only*): PFIZER LIMITED [GB/GB];
Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).


(71) Applicant (*for all designated States except GB, US*):
PFIZER INC. [US/US]; 235 East 42nd Street, New York,
NY 10017 (US).

Published:
— with international search report

(72) Inventors; and
(75) Inventors/Applicants (*for US only*): MIDDLETON,
Donald, Stuart [GB/GB]; Pfizer Global Research and
Development, Ramsgate Road, Sandwich, Kent CT13 9NJ
(GB). STOBIE, Alan [GB/GB]; Pfizer Global Research

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: LACTAMS AS TACHYKININ ANTAGONISTS



(I)

(57) Abstract: Compounds of the formula (I) or a pharmaceutically acceptable salt, prodrug, solvate or polymorph thereof, wherein: R is het^a; R¹ is phenyl optionally substituted by one or more substituents; m is 1-4; Z is selected from: a) N(R³)(R⁴X) wherein X is NR³R⁵, OR³, Oaryl¹, Ohet^b, Ohet^c, aryl¹, het^b or het^c; b) N(R³)Y wherein Y is aryl¹, het^b or het^c; and c) a 4-7 membered N containing saturated or partially saturated heterocycle said heterocycle attached to the alkylene link via said nitrogen atom, said heterocycle optionally containing an

additional 1-3 groups, each independently selected from C=O, NH, S(O)_p and O; optionally, said heterocycle is: (i) spirofused with het^b, such that both rings share 1 atom; or (ii) optionally independently substituted by 1-3 groups; wherein R³ and R⁵ are both independently selected from H and C₁₋₆alkyl; wherein R⁴ is selected from C₁₋₆alkylene; wherein R⁵ is selected from C(O)OR³, S(O)_pR³, S(O)_paryl¹, C(O)R³, and C(O)NR³R⁶; het^b is a 4-7 membered heterocycle containing 1-3 heteroatoms, each independently selected from N, O and S, said N being optionally substituted with O, said ring optionally containing 1-2 C=O groups, said ring being saturated or partially saturated, said ring being optionally benzofused, said ring being optionally substituted by 1-3 substituents; het^c and het^d are a 5-7 membered aromatic heterocycle containing 1-3 heteroatoms each independently selected from N, O and S, said ring being optionally benzofused, said ring system as a whole being optionally substituted by 1-3 substituents; aryl¹ is phenyl or naphthyl, each being optionally substituted by 1-3 substituents; p is 0, 1 or 2; and n is 0-4; are useful in treating or preventing a condition for which an NK₂ antagonist is efficacious.

WO 03/051868 A1

Lactams as Tachykinin Antagonists


This invention relates to therapeutic agents of the lactam family and to processes for the preparation of, intermediates used in the preparation of, compositions
5 containing and uses of, such derivatives.

International Patent Application Publication Number WO 96/05193 discloses various (azetidin-1-ylalkyl)lactams as tachykinin antagonists.

10 International Patent Application Publication Number WO97/25322 discloses various azetidinyalkyl derivatives of N-substituted nitrogen heterocycles as tachykinin antagonists.

The therapeutic agents of the present invention are antagonists of tachykinins,
15 including neurokinin A (NKA), neurokinin B (NKB) and Substance P, acting at the human neurokinin-1 (NK₁), neurokinin-2 (NK₂) or neurokinin-3 (NK₃) receptor, or a combination of two or more thereof. They are therefore useful for preventing or treating inflammatory disease, a central nervous system (CNS) disorder, a
gastro-intestinal (GI) disorder, a disease caused by *Helicobacter pylori* or other
20 urease positive Gram negative bacteria, urological conditions, a pulmonary disorder, an allergy, a hypersensitivity disorder, a vasospastic disease, a proliferative disorder, a fibrosing or collagen disease, reflux sympathetic dystrophy, an addiction disorder, a stress-related somatic disorder, a peripheral neuropathy, a neuropathological disorder, a disorder related to immune
25 enhancement or suppression, a rheumatic disease, an opthalmic disease, acute and chronic pain or a viral disease.

The present invention provides a compound of formula (I):



or a pharmaceutically acceptable salt, prodrug, solvate or polymorph thereof, wherein:

R is het^a;

5

R¹ is phenyl optionally substituted by one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, and C₁₋₆ alkyl optionally substituted by one or more halogen;

10 m is 1-4;

Z is selected from:

a) N(R³)(R⁴X)

wherein X is NR³R⁵, OR³, Oaryl¹, Ohet^b, Ohet^c, aryl¹, het^b or het^c;

15 b) N(R³)Y

wherein Y is aryl¹, het^b or het^c; and

c) a 4-7 membered N containing saturated or partially saturated heterocycle said heterocycle attached to the alkylene link via said nitrogen atom, said heterocycle optionally containing an additional 1-3 groups, each independently
20 selected from C=O, NH, S(O)_p and O; optionally, said heterocycle is:

i) spirofused with het^b, such that both rings share 1 atom; or

ii) substituted by 1-3 groups each independently selected from het^b, het^c, aryl¹, R³, R⁴OR³, R⁴C(O)R³, OR³, OR⁷OR³, OR⁴OC(O)R³, OR⁴OC(O)NR³R⁶, S(O)_pR⁴, C(O)R³, C(O)NR³R⁶, C(O)OR³, R⁷C(O)OR³,
25 C(O)R⁷OR³, C(O)OR⁷OR³, CF₃, NR³R⁶, R⁴NR³R⁵, OC(O)NR³R⁴ and NR³R⁵;

wherein R³ and R⁶ are both independently selected from H and C₁₋₆ alkyl;

wherein R⁴ and R⁷ are both independently selected from C₁₋₆ alkylene;

30 wherein R⁵ is selected from C(O)OR³, S(O)_pR³, S(O)_paryl¹, C(O)R³, and C(O)NR³R⁶;

het^b is a 4-7 membered heterocycle containing 1-3 heteroatoms, each independently selected from N, O and S, said N being optionally substituted with O, said ring optionally containing 1-2 C=O groups, said ring being saturated or partially saturated, said ring being optionally benzofused, said ring being
5 optionally substituted by 1-3 substituents selected from halo, R³, OR³, C(O)NR³R⁶, R⁷NR³R⁶, NR³R⁵, NHS(O)_pR⁴, S(O)_pNR³R⁶, S(O)_pR⁴, CN, NR³R⁶ and aryl¹;

het^a and het^c independently represent a 5-7 membered aromatic heterocycle
10 containing 1-3 heteroatoms each independently selected from N, O and S, said ring being optionally benzofused, said ring system as a whole being optionally substituted by 1-3 substituents, each independently selected from: halo, R³, OR³, C(O)NR³R⁶, R⁴NR³R⁶, NR³R⁵, NHS(O)_pR⁴, S(O)_pNR³R⁶, S(O)_pR⁴, CN, NR³R⁶, and R⁴NR³S(O)_pR³;

15 aryl¹ is phenyl or naphthyl, each optionally substituted by 1-3 substituents, each independently selected from: halo, R³, OR³, C(O)NR³R⁶, R⁷NR³R⁶, NR³R⁵, NHS(O)_pR⁴, S(O)_pNR³R⁶, S(O)_pR⁴, CN;

20 p is 0, 1 or 2; and

n is 0-4.

Halo includes fluoro, chloro, bromo and iodo groups.

25

Alkyl and alkylene include both straight chain and branched chain.

A pharmaceutically acceptable salt of a compound of the formula (I) may be readily prepared by mixing together solutions of a compound of the formula (I)
30 and the desired acid or base, as appropriate. The salt may precipitate from solution and be collected by filtration or may be recovered by evaporation of the solvent.

The pharmaceutically acceptable salts of the compounds of the formula (I) include the acid addition and the base salts thereof.

Suitable acid addition salts are formed from acids which form non-toxic salts and examples are the hydrochloride, hydrobromide, hydroiodide, sulphate, bisulphate,
5 nitrate, phosphate, hydrogen phosphate, acetate, maleate, fumarate, lactate, tartrate, citrate, gluconate, succinate, saccharate, benzoate, methanesulphonate, ethanesulphonate, benzenesulphonate, p-toluenesulphonate and pamoate salts.

10 Suitable base salts are formed from bases which form non-toxic salts and examples are the sodium, potassium, aluminium, calcium, magnesium, zinc and diethanolamine salts.

For a review on suitable salts see Berge et al, J. Pharm. Sci., 66, 1-19, 1977.

15

The pharmaceutically acceptable solvates of the compounds of the formula (I) include the hydrates thereof.


Also included within the present scope of the compounds of the formula (I) are
20 polymorphs thereof.

It will also be appreciated that the compounds of the invention will include prodrugs of (I) and pharmaceutically acceptable derivatives of (I) in which the functional groups explicitly recited above have been derivatised to provide
25 prodrugs which can be converted to the parent compound *in vivo*. Such prodrugs are discussed in Drugs of Today, 1983, 19, 499-538 and Annual Reports in Medicinal Chemistry, 1975, Vol. 10, Ch 31, 306-326.

A compound of the formula (I) contains one or more asymmetric carbon atoms
30 and therefore exists in two or more stereoisomeric forms. Where a compound of the formula (I) contains an alkenyl or alkenylene group, cis (E) and trans (Z) isomerism may also occur. The present invention includes the individual

stereoisomers of the compounds of the formula (I) and, where appropriate, the individual tautomeric forms thereof, together with mixtures thereof.

Those compounds of formula (I), which have the stereochemistry shown below
5 are particularly preferred.



Separation of diastereoisomers or cis and trans isomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or
10 H.P.L.C. of a stereoisomeric mixture of a compound of the formula (I) or a suitable salt or derivative thereof. An individual enantiomer of a compound of the formula (I) may also be prepared from a corresponding optically pure intermediate or by resolution, such as by H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the
15 diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable optically active acid or base, as appropriate.

Preferred embodiments of the present invention include compounds of formula (I) wherein:

20

Preferably R is pyridyl, optionally substituted by NR³R⁶, R³ or OR³.

More preferably R is pyridyl, optionally substituted by NMe₂, C₁₋₂ alkyl or OC₁₋₂ alkyl.

Yet more preferably R is pyridyl optionally substituted by methyl or ethyl.

25 Most preferably R is pyridyl optionally substituted by methyl.

Preferably the pyridyl moiety is substituted at the 2 position.

Preferably the lactam is attached to the pyridyl moiety at the 6 position of the
30 pyridyl group.

Preferably R^1 is phenyl optionally substituted by 1 or 2 halo substituents.

More preferably R^1 is phenyl, optionally substituted by 1 or 2 substituents selected from fluoro and chloro.

5

Yet more preferably R^1 is phenyl, 3,4-difluorophenyl, 3-chlorophenyl, 4-chlorophenyl or 3,4-dichlorophenyl.

Most preferably R^1 is 3,4-difluorophenyl, 4-chlorophenyl or 3,4-dichlorophenyl.

Most preferably R^1 is 3,4-dichlorophenyl.

10

Preferably m is 2-3

Most preferably m is 2.

Preferably n is 1-4.

15

More preferably n is 1-3.

Yet more preferably n is 1-2.

Most preferably n is 2.

Preferably R^3 is H or C_{1-4} alkyl; or

20

More preferably R^3 is H or C_{1-2} alkyl

Preferably R^4 is C_{1-4} alkylene

More preferably R^4 is C_{1-2} alkylene

25

Preferably R^5 is $C(O)OR^3$, $C(O)R^3$, $C(O)NR^3R^6$

Preferably R^6 is H, C_{1-4} alkyl; or

More preferably R^6 is H, C_{1-2} alkyl

Most preferably R^6 is H

30

Preferably R^7 is C_{2-8} alkylene

More preferably R^7 is C_{2-4} alkylene

Preferably Z is a piperidine or azetidine group optionally substituted by one or more of het^b , het^c , aryl^1 , OR^3 , R^3 and NR^3R^5 , wherein;

5 Het^b is a 5-6 membered saturated or partially saturated nitrogen containing heterocycle, said heterocycle optionally incorporating 1-2 groups each independently selected from O, C=O and N, said heterocycle being optionally benzofused, said heterocycle being optionally substituted by 1-2 substituents, each independently selected from OR^3 , R^3 , NR^3R^6 , NR^3R^5 , aryl^1 , SO_2R^4 and $\text{SO}_2\text{NR}^3\text{R}^6$;

10

Het^c is pyridyl, optionally substituted by 1 or 2 substituents each independently selected from halo and OR^3 ;

15

aryl^1 is phenyl, optionally substituted by 1 or 2 substituents each independently selected from halo and OR^3 ; and

R^3 , R^4 , R^5 and R^6 are as herein defined.

20

Preferably, the piperidine or aziridine group is substituted at the 4 or 3 position respectively.

25

Most preferably Z is a piperidine or azetidine group, optionally substituted by het^b , aryl^1 and NR^3R^5 ; wherein het^b is a morpholine or piperidine, optionally substituted at the 4 position by OH and or methyl; wherein;

aryl^1 is phenyl optionally substituted by OH; and

R^3 is H or methyl and R^5 is $\text{C}(\text{O})\text{CH}_3$.

30


Particularly preferred compounds include:

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 131)

- (5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-hydroxypiperidinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 135a)
- (5S)-5-(3,4-Dichlorophenyl)-5-{2-(4-methoxy-1-piperidinyl)ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 61)
- 5 (5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{{2-[4-hydroxy-4-phenyl]-1-piperidinyl}ethyl}-2-piperidinone (Example 134)
- (5S)-5-(3,4-Dichlorophenyl)-5-{2-[4-hydroxy-4-(2-pyridyl)-1-piperidinyl]ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 92)
- N*-(1-{2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-phenyl-4-piperidinyl)-acetamide (Example 90)
- 10 {5S)-5-(3,4-Dichlorophenyl)-1-(6-methoxy-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 119)
- 5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-oxo-1-piperidinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 168)
- 15 (5S)-5-(3,4-Dichlorophenyl)-5-{2-[3-(4-hydroxy-1-piperidinyl)-1-azetidiny]ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 73)
- N*-(1-{2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-piperidinyl)-*N*-methylacetamide (Example 158)
- 20 The invention further provides methods for the preparation of the compounds of the invention, which are described below and in the examples and Preparations section. The skilled man will appreciate that the compounds of the invention could be made by methods other than those herein described and or adaption of a plethora of methods known in the art. It is to be understood that the synthetic
- 25 transformation methods specifically mentioned herein may be carried out in various different sequences in order that the desired substances can be efficiently assembled. The skilled chemist will exercise his judgement and skill as to the most efficient series of reactions for synthesis of a given target substance.
- 30 It will be apparent to those skilled in the art that sensitive functional groups may need to be protected and deprotected during synthesis of a substance of the invention. This may be achieved by conventional techniques, for example as

described in "Protective Groups in Organic Synthesis" by T. W. Greene and P. G. M. Wuts, John Wiley and Sons Inc, 1991.

5 The compounds of formula (I) may be prepared in accordance with the following scheme:



Compounds of formula (I) may be prepared from the compounds of formula (II) under the conditions of process step (a) a reductive amination. This involves the reaction of amine Z-H with aldehyde (II), with a suitable metal hydride reducing agent (to reduce intermediate imine), optionally in the presence of a suitable base and/or acid, optionally in the presence of a Lewis acid catalyst, in a suitable solvent at room temperature.

15 Suitable conditions include:

1eq aldehyde (or HCl salt of), 1-2 eq of amine, 1-3 eq suitable reducing agent (e.g. $NaCNBH_3$, $NaBH(OAc)_3$), optionally in the presence of a base (eg Hünigs, Et_3N), and/or an acid (eg $AcOH$), in a suitable solvent (eg dichloromethane (DCM), or tetrahydrofuran (THF)) at from between 15 minutes and 72 hours.

20 Or

(under Lewis acid catalysis)-1 eq aldehyde, excess of amine, an excess of reducing agent, in the presence of a base (eg Et_3N), in a suitable solvent (e.g. $EtOH$), using 10 eq of Lewis acid (eg $Ti(OiPr)_4$).


Particularly suitable are:

1 eq aldehyde, 1 to 1.5 eq amine, 1 to 3 eq $\text{NaBH}(\text{OAc})_3$, optionally in the presence of 1 to 12 eq Et_3N , and optionally in the presence of 2 to 30 eq of AcOH , in DCM or THF for between 15 minutes and 72 hours at rt.


5 Or

(under Lewis acid catalysis)-1 eq aldehyde, 1.1 eq amine, 1.5 eq $\text{NaBH}(\text{OAc})_3$, in the presence of 2.5 eq Et_3N , in EtOH using 10 eq of $\text{Ti}(\text{OiPr})_4$.

10 The compounds of formula (II) may be prepared in accordance with the following scheme.



15 Compounds of formula (IV) may be prepared from the compounds of formula (III) under the conditions of process step (b), an alkylation reaction. Suitable conditions include using an excess of alkylating agent; in a preferred embodiment the conditions include an excess of a compound of formula (V)



where L is halo, with an excess of suitable base (typically an alkali metal salt), in a suitable solvent (eg ethylene glycol dimethyl ether (DME), 1-methyl-2-pyrrolidinone (NMP)), optionally in the presence of a catalyst (e.g. CuI), at the
5 reflux temp of the reaction for 1 to 24 hours.


Preferably a class of alkylation reaction known as the Goldberg reaction is used. This comprises 1.5 to 3 eq alkylating agent, V, where L is F, Cl or Br, 1.1 to 1.5
10 eq of K₂CO₃, or KO^tBu in NMP or DME at reflux for 1 to 24 hours.

Compounds of formula (II) may be prepared from the compounds of formula (IV) under the conditions of process step (c), a dioxalan hydrolysis reaction. Suitable conditions include hydrolysis under acidic conditions, such as 2.5N HCl in THF at
15 rt for 24 hours.

Process steps (b) and (c) may be carried out using "one pot" methodology without the isolation of compounds of formula (IV)

20 The compounds of formula (II) may also be prepared in accordance with the following scheme:

12




Compounds of formula (VI) may be prepared from the compounds of formula (III) under the conditions of process step (d), a dioxalan cleavage reaction. This is conducted under non-aqueous strongly acidic conditions, for example, Amberlyst® 15 resin in MeOH at rt for 18 hours.

Compounds of formula (II) may be prepared from the compounds of formula (VI) under the conditions of process step (e), an acetal hydrolysis reaction. Suitable conditions include hydrolysis under acidic conditions to give the aldehyde, for example 1.1-2.5N HCl in THF at rt for 18-24 hours.

The compounds of formula (I) may also be prepared in accordance with the following scheme:

13



Compounds of formula (VIII) may be prepared from the compounds of formula (VII) under the conditions of process step (a), a reductive amination reaction as discussed earlier.

5

Compounds of formula (I) may be prepared from the compounds of formula (VIII) under the conditions of process step (b), an alkylation reaction as discussed earlier.

- 10 In addition to the process routes already described, compounds of formula (I) where Z is $N(R^2)(R^3X)$, where R^2 is C_{1-6} alkyl, R^3 is C_{1-6} alkyl and X is het^a or het^b , may be prepared from the corresponding compounds of formula (I) where R^2 is hydrogen, by process step (a) a reductive amination reaction as discussed earlier.

15

All of the above reactions and the preparations of novel starting materials used in the preceding methods are conventional and appropriate reagents and reaction

conditions for their performance or preparation as well as procedures for isolating the desired products will be well-known to those skilled in the art with reference to literature precedents and the Examples and Preparations hereto.

- 5 The affinity of the compounds of formula (I), and their salts, for the human NK₂ receptor can be assessed *in vitro* by testing their ability to compete with [³H] or [¹²⁵I] NKA (neurokinin A) for binding to membranes prepared from Chinese hamster ovary cells expressing the cloned human NK₂ receptor using a modification of the method described in McLean, S. *et al*, J. Pharm. Exp. Ther.,
 10 267, 472-9 (1993). The membranes are incubated (90min, 25°C) with [³H] or [¹²⁵I] NKA and a range of concentrations of the test compound. Non specific binding was determined in the presence of 1μM SR-48968 (N-[(2S)-4-[4-(acetylamino)-4-phenyl-1-piperidiny]-2-(3,4-dichlorophenyl)butyl]-N-methylbenzamide).
- 15 All the compounds of the present invention (as exemplified herein) had a binding affinity for NK₂ receptors with Ki <1000nM.

The binding Ki expressed in nM for selected compounds of the present invention are expressed below:

20

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 131) Ki = 3.8

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-hydroxypiperidinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 135a)

25 Ki = 3.7

(5S)-5-(3,4-Dichlorophenyl)-5-{2-(4-methoxy-1-piperidinyl)ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 61) Ki = 12

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[4-hydroxy-4-phenyl]-1-piperidinyl}ethyl}-2-piperidinone (Example 134) Ki = 1.9

30 (5S)-5-(3,4-Dichlorophenyl)-5-{2-[4-hydroxy-4-(2-pyridyl)-1-piperidinyl]ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 92) Ki = 7

- N*-(1-{2-[(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-phenyl-4-piperidinyl)-acetamide (Example 90) $K_i = 2.4$
- (5*S*)-5-(3,4-Dichlorophenyl)-1-(6-methoxy-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 119) $K_i = 5.4$
- 5 5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-oxo-1-piperidinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 168) $K_i = 2.6$
- (5*S*)-5-(3,4-Dichlorophenyl)-5-{2-[3-(4-hydroxy-1-piperidinyl)-1-azetidiny]ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 73) $K_i = 2.2$
- N*-(1-{2-[(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-piperidinyl)-*N*-methylacetamide (Example 158) $K_i = 8$
- 10

The NK_2 receptor antagonist activity of the compounds of formula (I), and their salts, can be assessed *in vitro* by testing their ability to antagonise the contractile effects of the selective NK_2 receptor agonist [β Ala⁸] NKA₍₄₋₁₀₎ in human bladder

15 tissue or in rabbit pulmonary artery using the method of Patacchini and Maggi, Eur. J. Pharm., 236, 31-37 (1993).

Those compounds of the present invention, with binding affinity $K_i < 10\text{nM}$ (using the modified method described in McLean, S. *et al*, J. Pharm. Exp. Ther., 267,

20 472-9 (1993)) were profiled using the method of Patacchini. These especially preferred compounds had $K_b < 10\text{nM}$ or $pA_2 > 8$.

The compounds of formula (I), and their salts, can be tested for NK_2 receptor antagonist activity, *in vivo*, by testing their ability to inhibit bronchoconstriction induced by [β Ala⁸] NKA₍₄₋₁₀₎ in the anaesthetised guinea pig using the method

25 described by Mural *et al*, J. Pharm. Exp. Ther., 262, 403-8 (1992) or Metcalfe *et al*, Br. J. Pharmacol., 112, 563P (1994).

The compounds of formula (I), and their salts, can be tested for NK_3 receptor antagonist activity, *in vitro*, by testing their ability to compete with [³H] senktide (a selective NK_3 receptor agonist) on membranes prepared from guinea pig cortex, using the method described in Chretein, *et al*, Eur. J. Pharmacol, 256, 73-78

30 (1994).

The compounds of the present invention have been found to be potent NK₂ antagonists.

- 5 The present invention provides for the use of a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof as a medicament.

It further provides the use of compounds of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof in the preparation of a medicament for
10 the treatment of a condition for which an NK₂ antagonist is efficacious.

As NK₂ antagonists, the therapeutic agents of the present invention are therefore useful for preventing or treating an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system
15 (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis, ulcerative colitis or Crohn's disease, a disease caused by *Helicobacter pylori* or other urease positive Gram negative bacteria, urological conditions, ie a bladder
20 disorder or a urogenital tract disorder (such as incontinence, hyperreflexia, impotence or cystitis) or associated conditions such as benign prostatic hyperplasia, over active bladder and lower uterine tract symptoms, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a
25 hypersensitivity disorder such as to poison ivy, a vasospastic disease such as angina or Reynaud's disease, a proliferative disorder such as cancer or a disorder involving fibroblast proliferation, a fibrosing or collagen disease such as scleroderma or eosinophilic fasciitis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-
30 related somatic disorder, a neuropathological disorder such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a rheumatic

disease such as fibrositis, emesis, cough, migraine, an ophthalmic disease such as proliferative retinopathy, ocular inflammation, conjunctivitis, or a viral disease such as influenza or a cold.

- 5 The compounds of the present invention are also useful in the prevention and treatment of pain; both acute and chronic.

Acute pain is short-lived (e.g. post-operative pain). Chronic pain is usually defined as pain persisting from 3 to 6 months and includes somatogenic pains
10 and psychogenic pains. Psychogenic pain is that which occurs without an organic origin such as low back pain, atypical facial pain and chronic headache.

Other types of pain are caused by injury or infection of peripheral sensory nerves. It includes, but is not limited to pain from peripheral nerve trauma, herpes virus
15 infection, diabetes mellitus, causalgia, plexus avulsion, neuroma, limb amputation and vasculitis. Neuropathic pain is also caused by nerve damage from chronic alcoholism, human immunodeficiency virus infection, hypothyroidism, uremia or vitamin deficiencies.

20 Further types of pain are: neuropathic pain for example, AIDS neuropathy, post herpetic neuralgia, diabetic neuropathy and trigeminal neuralgia, fibromyalgia, pain associated with somatoform disorders, arthritic pain, cancer pain, neck pain, shoulder pain, back pain, cluster headaches, tension-type headache, migraine, herpes neuralgia, phantom limb pain, central pain, dental pain, NSAID-resistant
25 pain, visceral pain, surgical pain, post-operative pain, bone injury pain, pain during labor and delivery, pain resulting from burns, including sunburn, post-partum pain, angina pain, and genitourinary tract-related pain including cystitis. The term pain shall also preferably refer to nociceptive pain or nociception. Examples of acute pain include, in particular, post-operative pain such as pain
30 following a dental extraction, migraine, headache and trigeminal neuralgia.

Examples of chronic pain include, in particular, musculoskeletal pain or pain associated with musculo-skeletal disorders such as rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, sero-negative (non-rheumatoid)

- arthropathies, non-articular rheumatism and peri-articular disorders, and pain associated with cancer; pain with an inflammatory component such as rheumatic pain, secondary inflammatory osteoarthritis, dental pain and dysmenorrhoea; back pain such as low back pain e.g. spinal stenosis, prolapsed disc or sciatica;
- 5 trauma; herpes zoster; neuropathic pain such as post-herpetic neuralgia, trigeminal neuralgia, segmental or intercostal neuralgia, fibromyalgia, causalgia, peripheral neuropathy, diabetic neuropathy, chemotherapy-induced neuropathy, AIDS related neuropathy, occipital neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, reflex sympathetic dystrophy, phantom limb pain;
- 10 gastrointestinal pain such as functional bowel disorders, which include non-ulcer dyspepsia, non-cardiac chest pain and irritable bowel syndrome. (Irritable bowel syndrome is a gastrointestinal disorder characterised by the presence of abdominal pain and altered bowel habits without any evidence of organic disease); traumatic pain such as postoperative pain; traumatic avulsion pain such
- 15 as brachial plexus; chronic pain such as arthritic pain such as occurring in osteo-, rheumatoid or psoriatic arthritis; various forms of headache such as migraine, acute or chronic tension headache, temporomandibular pain, maxillary sinus pain, cluster headache; odontalgia; pain of visceral origin; nerve entrapment pain; sport's injury pain; menstrual pain; meningitis; arachnoiditis; angina; gout; burns;
- 20 scar pain; itch; and thalamic pain such as post stroke thalamic pain.

- Accordingly the present invention provides the use of compounds of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof in the preparation of a medicament for the treatment of a condition selected from: inflammatory
- 25 disease, a central nervous system (CNS) disorder, a gastro-intestinal (GI) disorder, a disease caused by *Helicobacter pylori* or other urease positive Gram negative bacteria, urological conditions, a pulmonary disorder, an allergy, a hypersensitivity disorder, a vasospastic disease, a proliferative disorder, a fibrosing or collagen disease, reflux sympathetic dystrophy, an addiction disorder,
- 30 a stress-related somatic disorder, a peripheral neuropathy, a neuropathological disorder, a disorder related to immune enhancement or suppression, a rheumatic disease, an ophthalmic disease, acute and chronic pain or a viral disease.

A preferred embodiment of the present invention provides the use of compounds of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof in the preparation of a medicament for the treatment of a condition selected from: urological conditions or acute and chronic pain.

5

Particularly suitable urological conditions include Incontinence, hyperreflexia, benign prostatic hyperplasia, over active bladder and lower uterine tract symptoms.

10 A particularly preferred urological condition is over active bladder.

A particularly preferred pain condition is neuropathic pain.

15 The invention also provides for a method of treating or preventing a condition for which an NK₂ antagonist is efficacious which comprises administering a therapeutically effective amount of a compound of formula (I) and pharmaceutically acceptable salts and prodrugs thereof to a patient in need of treatment.

20 The compounds of the formula (I) can also be administered in combination with other active agents. Preferred agents include: compounds which modulate the action of atrial natriuretic factor (also known as atrial natriuretic peptide), such as inhibitors of neutral endopeptidase; compounds which inhibit angiotensin-converting enzyme such as enalapril, and combined inhibitors of angiotensin-
25 converting enzyme and neutral endopeptidase such as omapatrilat; angiotensin receptor antagonists such as losartan; substrates for NO-synthase, i.e. L-arginine; calcium-channel blockers such as amlodipine; antagonists of endothelin receptors and inhibitors of endothelin-converting enzyme; cholesterol lowering agents e.g. statins and fibrates; antiplatelet and antithrombotic agents, e.g. tPA,
30 uPA, warfarin, hirudin and other thrombin inhibitors, heparin, thromboplastin activating factor inhibitors; insulin sensitising agents such as rezulin and hypoglycaemic agents such as glipizide; L-DOPA and carbidopa; acetylcholinesterase inhibitors such as donezipil or steroidal; non-steroidal anti-

inflammatory agents (NSAIDs) such as aspirin and ibuprofen; cGMP PDE₅ inhibitors such as sildenafil (Viagra™), vardenafil and cialis; muscarinic antagonists such as oxybutynin, tolterodine, propiverine, trospium chloride and darifenacin; alpha-adrenoceptor antagonists such as doxazosin (Cardura™),

5 tamsulosin, 4-Amino-6,7-dimethoxy-2-(5-methanesulfonamido-1,2,3,4-tetrahydroisoquinol-2-yl)-5-(2-pyridyl)quinazoline and 5-cyclopropyl-7-methoxy-2-(2-(4-morpholinylmethyl)-7,8-dihydro[1,6]naphthryridin-6(5H)-yl)-4(3H)-quinazollnone; serotonin/noradrenalin reuptake inhibitors (SNRI) such a duloxetine, venlafaxine and milnacipran; noradrenalin reuptake inhibitors such as

10 reboxetine; NK₁ antagonists such as (αR,9R)-7-[3,5-Bis(trifluoromethyl)benzyl]-8,9,10,11-tetrahydro-9-methyl-5-(4-methylphenyl)-7H-[1,4]diazocino[2,1-g][1,7]naphthyridine-6,13-dione (TAK-637), 5-[[[(2R,3S)-2-[(1R)-1-[3,5-Bis(trifluoromethyl)phenyl]ethoxy]-3-(4-fluorophenyl)-4-morpholinyl]methyl]-1,2-dihydro-3H-1,2,4-Triazol-3-one (MK-869), lanepitant, dapitant and 3-[[2-Methoxy-

15 5-(trifluoromethoxy)phenyl]methylamino]-2-phenyl-piperidine, (2S,3S); 5-HT_{1A} agonists/antagonists such as buspirone and robalzotan; COX₂ inhibitors such as celecoxib (Celebrix™), rofecoxib (Vioxx™) and valdecoxib; non-selective COX inhibitors (preferably with 'GI protection') such as HCT-1026 (nitroflurbiprofen); opioids such as morphine, codeine; tricyclic antidepressants such as desipramine

20 and amytriptiline; anticonvulsants such as gabapentin, serotonin reuptake inhibitors such as fluoxetine and sertraline; serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, sedatives such as amobarbital and temazepam, skeletal muscle relaxants such as baclofen; NMDA receptor antagonists such as dextromorphan and ketamine; vanilloid receptor

25 agonists such as resiniferatoxin; HMG-CoA reductase inhibitors such as atorvastatin (Lipitor™), simvastatin (Zocor™), pravastatin (Pravacol™) and rosuvastatin (Crestor™); and estrogenic modulators such as hormone replacement therapy and selective estrogen receptor modulators, such as lasofoxifene, tamoxifene and raloxifene.

30

The NK₂ antagonists of this invention can also be administered in combination with other active agents in the treatment of urological conditions, particularly

incontinence, hyperreflexia, benign prostatic hyperplasia, over active bladder and lower uterine tract symptoms.

Accordingly the present invention provides for the use of a compound of formula (I) in the preparation of a medicament in combination with an agent selected from: Muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); noradrenalin reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; PDE₅ inhibitors; COX₂ inhibitors; non-selective COX inhibitors; vanilloid receptor agonists; HMG-CoA reductase inhibitors; and estrogenic modulators and selective estrogen receptor modulators for the treatment of urological conditions.

The NK₂ antagonists of this invention can also be administered in combination with other active agents in the treatment of pain.

Accordingly the present invention provides for the use of a compound of formula (I) in the preparation of a medicament in combination with an agent selected from: NSAIDs, opioids, muscarinic antagonists; cholinergic analgesics; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); COX₂ inhibitors; non-selective COX inhibitors; tricyclic antidepressants, anticonvulsants, serotonin reuptake inhibitors, serotonin receptor agonists and antagonists, sedatives, skeletal muscle relaxant and NMDA receptor antagonists for the treatment of pain.

It is to be appreciated that all references herein to treatment include curative, palliative and prophylactic treatment.

The compounds of the formula (I) can be administered alone but will generally be administered in admixture with a suitable pharmaceutical excipient, diluent or carrier selected with regard to the intended route of administration and standard pharmaceutical practice.

The present invention provides for a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and a pharmaceutically acceptable diluent or carrier.

- 5 The compounds of formula (I) may also be administered in combination with other suitable therapeutic agents. Accordingly the present invention provides for a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and an agent selected from:
- 10 compounds which modulate the action of atrial natriuretic factor (also known as atrial natriuretic peptide), such as inhibitors of neutral endopeptidase; compounds which inhibit angiotensin-converting enzyme, and combined inhibitors of angiotensin-converting enzyme and neutral endopeptidase; angiotensin receptor antagonists; substrates for NO-synthase; calcium-channel blockers; antagonists of endothelin receptors and inhibitors of endothelin-converting enzyme;
- 15 cholesterol lowering agents; antiplatelet and antithrombotic agents, thromboplastin activating factor inhibitors; insulin sensitising agents and hypoglycaemic agents; acetylcholinesterase inhibitors; non-steroidal anti-inflammatory agents (NSAIDs); cGMP PDE₅ inhibitors; muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors
- 20 (SNRI); noradrenalin reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; COX₂ inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); opioids; tricyclic antidepressants; anticonvulsants, serotonin reuptake inhibitors; serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, sedatives, skeletal muscle relaxants;
- 25 NMDA receptor antagonists; vanilloid receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier.

- In preferred embodiments the invention provides a composition comprising a
- 30 compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and an agent selected from: Muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; PDE₅ inhibitors; COX₂

inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); vanilloid receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier.

5

Another preferred embodiment provides a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and an agent selected from: NSAIDs, opioids, muscarinic antagonists; cholinergic analgesics; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); COX₂ inhibitors; non-selective COX inhibitors; tricyclic antidepressants, anticonvulsants, serotonin reuptake inhibitors, serotonin receptor agonists and antagonists, sedatives, skeletal muscle relaxant and NMDA receptor antagonists, and a pharmaceutically acceptable diluent or carrier.

15 Where other therapeutic agents are given in combination with the compounds of formula (I) they may be administered separately, simultaneously or sequentially.

The present invention provides for:

- 20 a) a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and a pharmaceutically acceptable diluent or carrier;
- b) a composition comprising an agent selected from: compounds which modulate the action of atrial natriuretic factor (also known as atrial natriuretic peptide), such as inhibitors of neutral endopeptidase; compounds which inhibit
25 angiotensin-converting enzyme, and combined inhibitors of angiotensin-converting enzyme and neutral endopeptidase; angiotensin receptor antagonists; substrates for NO-synthase; calcium-channel blockers; antagonists of endothelin receptors and inhibitors of endothelin-converting enzyme; cholesterol lowering agents; antiplatelet and antithrombotic agents,
30 thromboplastin activating factor inhibitors; insulin sensitising agents and hypoglycaemic agents; acetylcholinesterase inhibitors; non-steroidal anti-inflammatory agents (NSAIDs); cGMP PDE₅ inhibitors; muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake

- inhibitors (SNRI); noradrenalin reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; COX₂ inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); opioids; tricyclic antidepressants; anticonvulsants, serotonin reuptake inhibitors; serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, sedatives, skeletal muscle relaxants; NMDA receptor antagonists; vanilloid receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier; and
- 5 c) a container.
- 10

The invention further provides in a preferred embodiment for:

- a) a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and a pharmaceutically acceptable diluent or carrier; and
- 15 b) a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and an agent selected from: muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; PDE₅ inhibitors; COX₂ inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); vanilloid receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier; and
- 20 c) a container.
- 25

In a further preferred embodiment, the invention provides for:

- a) a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and a pharmaceutically acceptable diluent or carrier; and
- 30 b) a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and an agent selected from:

- NSAIDs, opioids, muscarinic antagonists; cholinergic analgesics; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); COX₂ inhibitors; non-selective COX inhibitors; tricyclic antidepressants, anticonvulsants, serotonin reuptake inhibitors, serotonin receptor agonists and antagonists, sedatives, skeletal muscle relaxant and NMDA receptor antagonists, and a pharmaceutically acceptable diluent or carrier; and
- 5 c) a container.
- 10 For example, the compounds of the formula (I) can be administered orally, buccally or sublingually in the form of tablets, capsules, ovules, elixirs, solutions or suspensions, which may contain flavouring or colouring agents, for immediate-, delayed-, modified-, sustained-, pulsed- or controlled-release applications.
- 15 Such tablets may contain excipients such as microcrystalline cellulose, lactose, sodium citrate, calcium carbonate, dibasic calcium phosphate and glycine, disintegrants such as starch (preferably corn, potato or tapioca starch), sodium starch glycolate, croscarmellose sodium and certain complex silicates, and granulation binders such as polyvinylpyrrolidone, hydroxypropylmethylcellulose
- 20 (HPMC), hydroxypropylcellulose (HPC), sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, stearic acid, glyceryl behenate and talc may be included.
- Solid compositions of a similar type may also be employed as fillers in gelatin capsules. Preferred excipients in this regard include lactose, starch, a cellulose,
- 25 milk sugar or high molecular weight polyethylene glycols. For aqueous suspensions and/or elixirs, the compounds of the formula (I) may be combined with various sweetening or flavouring agents, colouring matter or dyes, with emulsifying and/or suspending agents and with diluents such as water, ethanol,
- 30 propylene glycol and glycerin, and combinations thereof.

The compounds of the formula (I) can also be administered parenterally, for example, intravenously, intra-arterially, intraperitoneally, intrathecally,

intraventricularly, intraurethrally, intrasternally, intracranially, intramuscularly or subcutaneously, or they may be administered by infusion techniques. For such parenteral administration they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or
5 glucose to make the solution isotonic with blood. The aqueous solutions should be suitably buffered (preferably to a pH of from 3 to 9), if necessary. The preparation of suitable parenteral formulations under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art.

10 The compounds of formula (I) can also be administered intranasally or by inhalation and are conveniently delivered in the form of a dry powder inhaler or an aerosol spray presentation from a pressurised container, pump, spray, atomiser or nebuliser, with or without the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, a
15 hydrofluoroalkane such as 1,1,1,2-tetrafluoroethane (HFA 134A [trade mark]) or 1,1,1,2,3,3,3-heptafluoropropane (HFA 227EA [trade mark]), carbon dioxide or other suitable gas. In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurised container, pump, spray, atomiser or nebuliser may contain a solution or
20 suspension of the active compound, e.g. using a mixture of ethanol and the propellant as the solvent, which may additionally contain a lubricant, e.g. sorbitan trioleate. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated to contain a powder mix of a compound of the formula (I) and a suitable powder base such as lactose or starch.

25

They may also be administered by the ocular route. For ophthalmic use, the compounds can be formulated as micronised suspensions in isotonic, pH adjusted, sterile saline, or, preferably, as solutions in isotonic, pH adjusted, sterile saline, optionally in combination with a preservative such as a benzylalkonium
30 chloride. Alternatively, they may be formulated in an ointment such as petrolatum.

Alternatively, the compounds of the formula (I) can be administered in the form of a suppository or pessary, or they may be applied topically in the form of a gel, hydrogel, lotion, solution, cream, ointment or dusting powder. The compounds of the formula (I) may also be dermally or transdermally administered, for example, by the use of a skin patch. They may also be administered by the pulmonary or rectal routes.

For application topically to the skin, the compounds of the formula (I) can be formulated as a suitable ointment containing the active compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, they can be formulated as a suitable lotion or cream, suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, a polyethylene glycol, liquid paraffin, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

The compounds of the formula (I) may also be used in combination with a cyclodextrin. Cyclodextrins are known to form inclusion and non-inclusion complexes with drug molecules. Formation of a drug-cyclodextrin complex may modify the solubility, dissolution rate, bioavailability and/or stability property of a drug molecule. Drug-cyclodextrin complexes are generally useful for most dosage forms and administration routes. As an alternative to direct complexation with the drug the cyclodextrin may be used as an auxiliary additive, e.g. as a carrier, diluent or solubiliser. Alpha-, beta- and gamma-cyclodextrins are most commonly used and suitable examples are described in WO-A-91/11172, WO-A-94/02518 and WO-A-98/55148.


The compounds of the invention may have the advantage that they are more potent, have a longer duration of action, are more stable, have fewer side effects, are more selective (in particular for the NK₂ receptor) and have improved cardiac safety, or have other more useful properties than the compounds of the prior art.

The following examples illustrate the preparation of the compounds of the formula (I):

Example 1

(5S)-5-(3,4-Dichlorophenyl)-5-[2-(4-hydroxy-1-piperidiny)ethyl]-1-(2-pyridinyl)-2-

piperidinone



Sodium triacetoxyborohydride (262mg, 1.24mmol) was added to a solution of the aldehyde from preparation 11a (250mg, 0.62mmol) and 4-hydroxypiperidine (90mg, 0.9mmol) in dichloromethane (100ml), and the reaction stirred at room temperature for 90 minutes. The mixture was washed with water, dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol (100:0 to 95:5) to afford the title compound as a white solid, 156mg.


¹Hnmr (CDCl₃, 400MHz) δ: 1.58 (m, 2H), 1.90-2.30 (m, 11H), 2.60 (m, 2H), 2.75 (m, 2H), 3.75 (m, 1H), 3.96 (d, 1H), 4.60 (d, 1H), 7.14 (m, 1H), 7.25 (m, 1H), 7.40 (d, 1H), 7.50 (s, 1H), 7.72 (s, 2H), 8.50 (d, 1H).

LRMS : m/z (TSP⁺) 448.1, 450.1 [MH⁺]

Microanalysis found: C, 58.34; H, 6.15; N, 8.57. C₂₃H₂₇Cl₂N₃O₂·0.10CH₂Cl₂·1H₂O requires C, 58.42; H, 6.20; N, 8.85%.




Examples 2 to 10





The following examples of general structure:



were prepared from the aldehyde hydrochloride from preparation 11b and the
5 appropriate amines, following a similar procedure to that described in example 1.

Ex No.	R	Yield (%)	Data
2 ^{1a}		58 yellow gum	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.40 (s, 9H), 1.80-2.60 (m, 9H), 3.16 (m, 3H), 3.90 (d, 1H), 4.58 (d, 1H), 4.88 (s, 1H), 7.14 (m, 1H), 7.22 (d, 1H), 7.41 (d, 1H), 7.48 (d, 1H), 7.72 (d, 2H), 8.50 (d, 1H). LRMS: m/z (TSP ⁺) 507.2, 509.2 [MH ⁺] Microanalysis found: C, 57.11; H, 6.96; N, 10.42. C ₂₈ H ₃₂ Cl ₂ N ₄ O ₃ ·0.05CH ₂ Cl ₂ ·1H ₂ O requires C, 56.91; H, 6.31; N, 10.60%.
3 ^{2b}		29 white solid	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.64-1.96 (m, 2H), 2.16 (m, 2H), 2.23-2.43 (m, 7H), 2.60 (m, 1H), 2.78 (m, 2H), 2.97 (m, 1H), 3.20 (m, 4H), 3.43 (m, 2H), 3.92 (d, 1H), 4.52 (m, 3H), 7.19 (m, 2H), 7.43 (d, 1H), 7.47 (s, 1H), 7.74 (m, 2H), 8.52 (d, 1H). LRMS : m/z (TSP ⁺) 488.8, 490.7 [M-SO ₂ NH ₂] ⁺ Microanalysis Found: C, 40.29; H, 5.77; N, 11.18. C ₂₅ H ₃₂ Cl ₂ N ₆ O ₃ S·3HCl·4H ₂ O requires C, 40.09; H, 5.79; N, 11.22%

4 ^{1c}		66	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.50 (bs, 3H), 1.80-2.35 (m, 8H), 2.56 (m, 1H), 3.60 (bs, 4H), 3.95 (d, 1H), 4.60 (d, 1H), 7.08 (dd, 1H), 7.24 (m, 1H), 7.38 (d, 1H), 7.44 (d, 1H), 7.66 (d, 2H), 8.45 (d, 1H). Microanalysis, Found: C, 59.72; H, 5.71; N, 9.56. C ₂₂ H ₂₅ Cl ₂ N ₃ O ₂ ; 0.1CH ₂ Cl ₂ ; 1H ₂ O Calc. C, 59.70; H, 5.76; N, 9.45%.
5 ^d		38 yellow solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.28 (s, 3H), 1.60 (m, 2H), 2.24 (m, 10H), 2.55-2.82 (m, 4H), 3.80 (m, 1H), 3.94 (d, 1H), 4.60 (d, 1H), 7.15 (dd, 1H), 7.34 (d, 1H), 7.44 (d, 1H), 7.52 (s, 1H), 7.74 (d, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 462.2, 464.2 [MH ⁺] Microanalysis found : C, 56.46; H, 6.51; N, 8.07. C ₂₄ H ₂₈ Cl ₂ N ₃ O ₂ ; 0.5CH ₂ Cl ₂ ; H ₂ O requires C, 56.28; H, 6.17; N, 8.04%.
6 ^e		57 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.62 (m, 2H), 1.74 (m, 2H), 1.90 (m, 4H), 2.01 (m, 1H), 2.15 (m, 2H), 2.30 (m, 2H), 2.44 (m, 1H), 2.54 (m, 1H), 2.62 (d, 3H), 2.83 (m, 2H), 4.00 (d, 1H), 4.20 (m, 1H), 4.65 (d, 1H), 7.10 (m, 1H), 7.20 (m, 1H), 7.30 (d, 2H), 7.39 (d, 1H), 7.46 (s, 1H), 7.66 (m, 2H), 7.74 (d, 2H), 8.45 (d, 1H). LRMS : m/z (TSP ⁺) 601.4 [M ⁺]

7 ⁱ		39 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.18 (t, 3H), 1.82-2.02 (m, 4H), 2.07-2.37 (m, 7H), 2.58 (m, 1H), 3.36 (m, 4H), 3.90 (d, 1H), 4.07 (q, 2H), 4.61 (d, 1H), 7.10 (dd, 1H), 7.19 (d, 1H), 7.38 (d, 1H), 7.43 (s, 1H), 7.67 (m, 2H), 8.44 (d, 1H). LRMS : m/z (TSP ⁺) 505.3, 507.2 [MH ⁺] Microanalysis found: C, 58.64; H, 6.16; N, 10.52. C ₂₅ H ₃₀ Cl ₂ N ₄ O ₃ ·0.45H ₂ O
8 ^g		4 white solid	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.92 (m, 4H), 2.10-2.40 (m, 6H), 2.50 (s, 3H), 2.54-2.88 (m, 8H), 3.92 (d, 1H), 4.64 (d, 1H), 7.16 (m, 1H), 7.20 (d, 1H), 7.40 (d, 1H), 7.46 (s, 1H), 7.72 (d, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 461.2, 463.2 [MH ⁺]
9 ^h		19	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.78 (m, 2H), 1.85 (m, 2H), 2.18 (m, 2H), 2.24 (m, 3H), 2.54 (m, 5H), 2.77 (s, 3H), 3.28 (m, 2H), 3.35 (m, 2H), 3.88 (d, 1H), 4.64 (d, 1H), 7.12 (m, 1H), 7.18 (d, 1H), 7.38 (d, 1H), 7.44 (d, 1H), 7.68 (d, 2H), 8.46 (d, 1H). LRMS : m/z (TSP ⁺) 525.1, 527.1 [MH ⁺]
10 ^h		43 White solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.40 (s, 9H), 1.70-2.60 (m, 14H), 3.40 (bs, 4H), 3.90 (d, 1H), 4.62 (d, 1H), 7.12 (m, 1H), 7.22 (d, 1H), 7.40 (d, 1H), 7.44 (s, 1H), 7.70 (s, 2H), 8.44 (d, 1H). LRMS : m/z (TSP ⁺) 547.2, 549.2 [MH ⁺] Microanalysis found: C, 58.52; H, 6.32; N, 9.13. C ₂₈ H ₃₆ Cl ₂ N ₄ O ₃ ·0.2CH ₂ Cl ₂ ·H ₂ O requires C, 58.15; H, 6.64; N, 9.62%.

1 = the aldehyde from preparation 11a was used

2 = prepared as the HCl salt

Starting amines:

a = *t*-butyl N-(2-aminoethyl)carbamate

b = 4-(3-azetidiny)-1-piperazine sulphonamide trifluoroacetate as prepared in WO 9725322

c = morpholine

5 d = 4-methyl-4-piperidinol from preparation 26

e = N-methyl-4-(4-piperidinyl)benzenesulphonamide as prepared in EP291210

f = ethyl 1-piperazinecarboxylate


g = 1-methyl-1,4-diazepine as prepared in J.A.C.S., 76, 5805

h = 1-(methylsulphonyl)-1,4-diazepine from preparation 82

10 i = *t*-butyl 1-homopiperazinecarboxylate




Examples 11 to 16




The following examples of general structure:



15

were prepared from the aldehyde from preparation 12a and the appropriate amines, following the procedure described in example 1.

Example	R	Yield (%)	Data
11 ^a		74 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.54 (d, 2H), 1.84-2.30 (m, 12H), 2.50-2.74 (m, 6H), 3.66 (m, 1H), 3.94 (d, 1H), 4.60 (d, 1H), 7.00 (d, 1H), 7.24 (d, 1H), 7.42 (d, 2H), 7.62 (m, 2H). LRMS : m/z (TSP ⁺) 462.1, 464.1 [MH ⁺] Microanalysis found: C, 59.81; H, 6.34; N, 8.57. C ₂₄ H ₂₈ Cl ₂ N ₃ O ₂ ·0.1CH ₂ Cl ₂ ·0.7H ₂ O requires C, 59.87; H, 6.38; N, 8.69%.
12 ^b		64 yellow solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.20 (s, 3H), 1.55 (d, 2H), 1.70-2.65 (m, 18H), 3.90 (d, 1H), 4.60 (d, 1H), 7.00 (d, 1H), 7.25 (d, 1H), 7.40 (dd, 2H), 7.60 (m, 2H). LRMS : m/z (TSP ⁺) 476.2, 478.2 [MH ⁺] Microanalysis found: C, 59.36; H, 6.63; N, 8.08. C ₂₅ H ₃₁ Cl ₂ N ₃ O ₂ ·0.05CH ₂ Cl ₂ ·1.3H ₂ O requires C, 59.68; H, 6.74; N, 8.34%.
13 ^c		70	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.10 (s, 6H), 1.22 (m, 2H), 1.57-1.80 (m, 6H), 1.92 (m, 3H), 2.10 (m, 2H), 2.25 (m, 2H), 2.50 (m, 4H), 2.82 (m, 2H), 4.88 (d, 1H), 4.50 (d, 1H), 6.95 (d, 1H), 7.20 (d, 1H), 7.38 (d, 2H), 7.56 (m, 2H). LRMS : m/z (ES ⁺) 504, 506 [MH ⁺]

14 ^d		51 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.90-2.20 (m, 12H), 2.30 (m, 4H), 2.52 (s, 3H), 2.58 (m, 2H), 2.70 (d, 1H), 3.92 (d, 1H), 4.65 (d, 1H), 5.40 (d, 1H), 7.00 (d, 1H), 7.18-7.36 (m, 6H), 7.40 (d, 2H), 7.60 (dd, 2H). LRMS : m/z (TSP ⁺) 580.2, 582.2 [MH ⁺]
15 ^a		39 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.70 (m, 3H), 1.95-2.05 (m, 4H), 2.18 (m, 2H), 2.30 (m, 5H), 2.54 (s, 3H), 2.54 (m, 1H), 2.65 (m, 2H), 3.95 (d, 1H), 4.62 (d, 1H), 7.00 (m, 3H), 7.30 (m, 1H), 7.40 (dd, 4H), 7.60 (dd, 2H). LRMS : m/z (ES ⁺) 556, 558 [MH ⁺] Microanalysis found: C, 63.60; H, 5.76; N, 7.41. C ₃₀ H ₃₂ Cl ₂ FN ₃ O ₂ ; 0.1CH ₂ Cl ₂ ; 0.1H ₂ O requires C, 63.78; H, 5.76; N, 7.41%.
16 ^c		62 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.95-2.40 (m, 11H), 2.58 (s, 3H), 2.58 (m, 1H), 3.65 (m, 4H), 3.90 (d, 1H), 4.62 (d, 1H), 7.00 (d, 1H), 7.28 (d, 1H), 7.40 (d, 2H), 7.60 (dd, 2H). LRMS : m/z (TSP ⁺) 448.2, 450.2 [MH ⁺] Microanalysis found: C, 60.64; H, 5.98; N, 9.16. C ₂₃ H ₂₇ Cl ₂ N ₃ O ₂ ; 0.1CH ₂ Cl ₂ requires C, 60.73; H, 6.00; N, 9.20%.

Starting amines:

a= 4-hydroxypiperidine

b= 4-methyl-4-piperidinol from preparation 26

c = 2-(4-piperidinyl)-2-propanol as prepared in EP 625509

35

d = N-(4-phenyl-4-piperidinyl)acetamide from preparation 32


e = 4-(4-fluorophenyl)-4-piperidinol

f = morpholine

5

Example 17

(5S)-5-(3,4-Dichlorophenyl)-5-[2-(4-phenyl-1-piperidinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone



- 10 Sodium triacetoxyborohydride (267mg, 1.26mmol) was added to a solution of the aldehyde hydrochloride from preparation 11b (250mg, 0.63mmol) and 4-phenylpiperidine (151mg, 0.94mmol) in dichloromethane (200ml), and the reaction stirred at room temperature for 2 hours. The mixture was washed with 2N sodium hydroxide solution (200ml), the aqueous layer extracted with
- 15 dichloromethane (2x200ml), the combined organic extracts were dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as eluant to afford the title compound as a white foam.


- ¹Hnmr (CDCl₃, 400MHz) δ: 1.74 (m, 6H), 1.83-2.08 (m, 5H), 2.18 (m, 2H), 2.32 (m, 2H), 2.40 (m, 1H), 2.60 (m, 1H), 3.90 (d, 1H), 4.68 (d, 1H), 7.18 (m, 4H), 7.28 (m, 3H), 7.41 (d, 1H), 7.50 (s, 1H), 7.73 (m, 2H), 8.50 (d, 1H).
- 20

Microanalysis found: C, 66.65; H, 6.16; N, 8.06. C₂₉H₃₁Cl₂N₃O;0.78H₂O requires C, 66.66; H, 6.28; N, 8.04%.

25



Examples 18 to 25




The following examples of general structure:





were prepared from the aldehyde hydrochloride from preparation 11b and the
 5 appropriate amine, following a similar procedure to that described in example 17.

Example	R	Yield (%)	Data
18 ¹⁸		50 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 2.24 (m, 2H), 2.41-2.75 (m, 5H), 2.91 (m, 1H), 3.80-4.00 (m, 2H), 4.40 (d, 1H), 4.54 (d, 1H), 7.30 (m, 4H), 7.39 (d, 1H), 7.43 (s, 1H), 7.54 (m, 3H), 7.88 (d, 1H), 8.15 (dd, 1H), 8.61 (d, 1H), 9.90 (bs, 1H), 10.05 (bs, 1H). LRMS : m/z (TSP ⁺) 454.0, 456.0 [MH ⁺]

19 ^{1b}		65 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 2.25 (m, 2H), 2.42 (m, 1H), 2.57 (m, 3H), 2.75 (m, 1H), 2.90 (m, 1H), 3.01 (m, 2H), 3.16 (m, 2H), 4.35 (m, 2H), 4.58 (d, 1H), 7.21 (m, 5H), 7.38 (d, 1H), 7.42 (d, 2H), 7.50 (s, 1H), 7.84 (d, 1H), 8.05 (dd, 1H), 8.62 (d, 1H), 9.82 (bs, 1H), 9.96 (bs, 1H). LRMS : m/z (TSP ⁺) 468.0, 470.0 [MH ⁺] Microanalysis found: C, 57.32; H, 5.50; N, 7.59. C ₂₆ H ₂₇ Cl ₂ N ₃ O; 2HCl; 0.05CH ₂ Cl ₂ requires C, 57.35; H, 5.38; N, 7.70%.
20 ^c		69 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.63-1.88 (m, 6H), 2.08-2.40 (m, 5H), 2.55-2.64 (m, 1H), 2.89 (m, 4H), 3.57 (m, 4H), 3.93 (d, 1H), 4.62 (d, 1H), 7.16 (m, 1H), 7.23 (d, 1H), 7.42 (d, 1H), 7.48 (s, 1H), 7.74 (d, 2H), 8.49 (m, 1H). LRMS : m/z (TSP ⁺) 473, 475 [MH ⁺] Microanalysis found: C, 62.66; H, 6.21; N, 8.80. C ₂₅ H ₂₉ Cl ₂ N ₃ O ₂ ; 0.1CH ₂ Cl ₂ requires C, 62.43; H, 6.09; N, 8.70%.

21 ^{2d}		15 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.68 (m, 2H), 1.91-2.16 (m, 5H), 2.16-2.24 (m, 2H), 2.24-2.40 (m, 4H), 2.53-2.73 (m, 3H), 3.95 (d, 1H), 4.70 (d, 1H), 7.14 (m, 1H), 7.27 (m, 2H), 7.36 (m, 2H), 7.41 (d, 1H), 7.45 (d, 2H), 7.50 (s, 1H), 7.71 (m, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 425.1, 526.1 [MH ⁺]
22 ^{3a}		64 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.38-1.78 (m, 3H), 1.85-2.43 (m, 11H), 2.51-2.78 (m, 3H), 3.98 (d, 1H), 4.72 (d, 1H), 7.02 (m, 2H), 7.15 (dd, 1H), 7.25 (d, 1H), 7.42 (m, 3H), 7.51 (s, 1H), 7.73 (m, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 542.1, 544.1 [MH ⁺] Microanalysis found: C, 63.55; H, 5.87; N, 7.48. C ₂₉ H ₃₀ Cl ₂ FN ₃ O ₂ ·0.3H ₂ O requires C, 63.58; H, 5.63; N, 7.67%.
23 ¹		77 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 0.80 (t, 3H), 1.42 (m, 4H), 1.58 (m, 3H), 1.95-2.02 (m, 3H), 2.17 (m, 4H), 2.25 (m, 2H), 2.50 (m, 3H), 3.88 (d, 1H), 4.60 (d, 1H), 7.10 (dd, 1H), 7.18 (d, 1H), 7.38 (d, 1H), 7.42 (s, 1H), 7.64 (m, 2H), 8.42 (d, 1H). LRMS : m/z (TSP ⁺) 476.2, 478.2 [MH ⁺]

24 ^g		46 clear oil	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.65 (m, 2H), 1.82 (m, 2H), 1.92-2.36 (m, 9H), 2.58 (m, 1H), 2.63 (m, 2H), 3.90 (d, 1H), 4.60 (d, 1H), 4.98 (s, 2H), 7.04 (m, 1H), 7.14 (m, 2H), 7.20 (m, 3H), 7.39 (d, 1H), 7.45 (s, 1H), 7.66 (s, 2H), 8.45 (d, 1H). LRMS : m/z (TSP ⁺) 536.2, 538.2 [MH ⁺]
25 ^h		20 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.60 (d, 2H), 1.99 (m, 4H), 2.16 (m, 2H), 2.20-2.40 (m, 5H), 2.58 (m, 1H), 2.72 (m, 2H), 3.95 (d, 1H), 4.61 (d, 1H), 7.14 (m, 1H), 7.21 (m, 1H), 7.32 (d, 1H), 7.40 (d, 1H), 7.47 (m, 2H), 7.61 (dd, 1H), 7.66 (m, 2H), 7.81 (d, 1H), 8.46 (d, 1H). LRMS : m/z (TSP ⁺) 550.2, 552.2 [MH ⁺]

1 = Isolated as the dihydrochloride salt

2 = Free base of the aldehyde used

3 = Tetrahydrofuran was used as the reaction solvent

5 Starting amines:

a = benzylamine

b = phenethylamine

c = 7-oxa-2-azaspiro[3.5]nonane p-toluenesulphinate from preparation 38

d = 4-phenyl-4-piperidinol

10 e = 4-(4-fluorophenyl)-4-piperidinol


f = 4-ethyl-4-piperidinol from preparation 56

g = spiro[isobenzofuran-1(3H),4'-piperidine] prepared as in EP 630887

h = 3-oxaspiro[isobenzofuran-1(3H),4'-piperidine] prepared as in EP 630887




Examples 26 to 31



The following examples of general structure:



- 5 were prepared from the aldehyde hydrochloride from preparation 12b and the appropriate amine, following the procedure described in example 17.

Example	R	Yield (%)	Data
26 ^a		29 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.14 (dq, 2H), 1.45 (m, 2H), 1.57 (m, 1H), 1.63-1.82 (m, 4H), 2.00-2.18 (m, 2H), 2.18-2.34 (m, 2H), 2.56 (m, 4H), 2.66 (m, 2H), 3.26 (m, 4H), 3.85 (d, 1H), 3.94 (m, 2H), 4.52 (d, 1H), 7.00 (d, 1H), 7.21 (d, 1H), 7.40 (d, 2H), 7.59 (m, 2H). LRMS : m/z (TSP ⁺) 502.1, 504.1 [MH ⁺] Microanalysis found: C, 63.30; H, 6.68; N, 8.16. C ₂₇ H ₃₃ Cl ₂ N ₃ O ₂ ·0.15CH ₂ Cl ₂ requires C, 63.29; H, 6.51; N, 8.16%.

27 ^{1b}		50 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.62-1.88 (m, 6H), 2.07-2.39 (m, 5H), 2.58 (m, 4H), 2.90 (m, 4H), 3.58 (m, 4H), 3.90 (d, 1H), 4.58 (d, 1H), 7.01 (d, 1H), 7.24 (d, 1H), 7.42 (d, 2H), 7.60 (m, 2H). LRMS : m/z (ES ⁺) 488, 490 [MH ⁺] Microanalysis found: C, 62.96; H, 6.50; N, 8.45. C ₂₆ H ₃₁ N ₃ O ₂ Cl ₂ ; 0.1CH ₂ Cl ₂ requires C, 63.08; H, 6.33; N, 8.46%
28 ^c		51 clear oil	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.68 (m, 2H), 1.82-2.04 (m, 8H), 2.14 (m, 2H), 2.26 (m, 2H), 2.50 (s, 3H), 2.58 (m, 1H), 2.84 (m, 2H), 3.90 (d, 1H), 4.62 (d, 1H), 6.96 (d, 1H), 7.08 (m, 2H), 7.22 (m, 1H), 7.38 (m, 2H), 7.58 (m, 3H), 8.46 (d, 1H). LRMS : m/z (TSP ⁺) 523.9, 525.9 [MH ⁺]
29 ^d		42 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.65 (m, 4H), 1.82-2.08 (m, 5H), 2.18 (m, 2H), 2.30 (m, 2H), 2.40 (m, 1H), 2.60 (m, 1H), 2.90 (m, 2H), 4.00 (d, 1H), 4.66 (d, 1H), 7.18 (m, 4H), 7.28 (m, 3H), 7.41 (d, 1H), 7.52 (s, 1H), 7.72 (m, 2H), 8.50 (d, 1H). LRMS : m/z (ES ⁺) 552, 554 [MH ⁺]

30 ^a		18 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.86-2.19 (m, 8H), 2.34 (m, 6H), 2.48 (m, 5H), 3.86 (m, 1H), 4.62 (m, 1H), 5.15 (bs, 2H), 6.98 (m, 1H), 7.22 (m, 2H), 7.34 (m, 4H), 7.39 (m, 2H), 7.57 (m, 2H). LRMS : m/z (TSP ⁺) 566.1, 568.1 [MH ⁺]
31 ^f		33	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.96 (m, 2H), 2.05 (m, 1H), 2.16 (m, 2H), 2.40 (m, 3H), 2.50 (m, 5H), 3.45 (m, 2H), 3.90 (d, 1H), 4.68 (d, 1H), 6.58 (d, 2H), 6.96 (d, 1H), 7.22 (s, 1H), 7.40 (m, 3H), 7.58 (d, 2H), 8.15 (d, 1H). LRMS : m/z (TSP ⁺) 524.9, 526.9 [MH ⁺]

1 = the aldehyde from preparation 12a was used

Starting amines:

a = 3-tetrahydro-2H-pyran-4-ylazetidine from preparation 31

5 b = 7-oxa-2-azaspiro[3.5]nonane p-toluenesulphinate from preparation 38

c = 2-(4'-piperidinyl)pyridine as prepared in EP 630887


d = 4-phenylpiperidine

e = 4-phenyl-4-piperidinecarboxamide as prepared in WO 9426735

f = 1 (2-pyridinyl)piperazine

Example 32

(5S)-5-(3,4-Dichlorophenyl)-1-(2-pyridinyl)-5-(2-{[2-(4-pyridinyloxy)ethyl]amino}ethyl)-2-piperidinone trihydrochloride



- 5 A solution of 2-(4-pyridinyloxy)ethylamine (EP 982322) (90.6mg, 0.66mmol) in dichloromethane (0.5ml) was added to a solution of the aldehyde from preparation 11a (238.2mg, 0.66mmol) in dichloromethane (10ml), and the solution stirred for 5 minutes. Sodium triacetoxyborohydride (139mg, 0.66mmol) was added and the reaction stirred for 1 hour. The reaction was washed with
- 10 aqueous sodium bicarbonate solution (20ml), brine (20ml), then dried (MgSO₄) and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant. The product was dissolved in a minimum volume of dichloromethane, 1N ethereal hydrochloric acid added, and the mixture
- 15 evaporated under reduced pressure to afford the title compound as a white foam, 170mg.

¹Hnmr (CD₃OD, 400MHz) δ: 2.24-2.47 (m, 3H), 2.47-2.60 (m, 2H), 2.77-2.92 (m, 2H), 3.01 (m, 1H), 3.73 (s, 2H), 4.29 (d, 1H), 4.45 (d, 1H), 4.64 (s, 2H), 7.45 (d, 1H), 7.60 (m, 3H), 7.73 (s, 1H), 7.80 (dd, 1H), 8.07 (d, 1H), 8.61 (d, 2H), 8.72


20 (d, 2H).

LRMS : m/z (TSP⁺) 485.1, 487.2 [MH⁺]

Microanalysis found: C, 45.06; H, 5.32; N, 8.19. C₂₅H₂₆Cl₂N₃O₂·3HCl·4H₂O requires C, 45.03; H, 5.59; N, 8.40%.



Examples 33 to 40



The following examples of general structure:





were prepared from the aldehyde from preparation 11a, and the appropriate
 5 amines, following a similar procedure to that described in example 32.


Example	R	Yield (%)	Data
33 ^{1a}		53 white solid	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.19-2.49 (m, 3H), 2.49-2.65 (m, 2H), 2.88-2.97 (m, 2H), 2.97-3.17 (m, 1H), 3.52 (s, 2H), 4.28 (d, 1H), 4.46 (d, 1H), 4.57 (m, 2H), 7.47 (d, 1H), 7.60 (d, 1H), 7.73 (s, 1H), 7.80 (d, 1H), 8.00-8.17 (m, 2H), 8.32 (d, 1H), 8.59-8.67 (m, 3H), 8.70 (s, 1H). Microanalysis found: C, 46.30; H, 5.29; N, 8.39. C ₂₅ H ₂₆ Cl ₂ N ₃ O ₂ ·3HCl·3H ₂ O requires C, 46.28; H, 5.44; N, 8.64%.

34 ^{1b}		48 white solid	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.23-2.45 (m, 3H), 2.45-2.62 (m, 2H), 2.79-2.95 (m, 2H), 3.00 (t, 1H), 3.50 (m, 2H), 4.28 (d, 1H), 4.46 (d, 1H), 4.69 (m, 2H), 7.29-7.41 (m, 2H), 7.46 (d, 1H), 7.60 (d, 1H), 7.73 (s, 1H), 7.80 (dd, 1H), 8.06 (d, 1H), 8.23 (dd, 1H), 8.35 (d, 1H), 8.52-8.69 (m, 2H). LRMS : m/z (TSP ⁺) 485.2, 487.2 [MH ⁺] Microanalysis found: C, 45.82; H, 5.33; N, 8.25. C ₂₅ H ₂₆ Cl ₂ N ₃ O ₂ ·3HCl·3.5H ₂ O requires C, 45.64; H, 5.52; N, 8.52%.
35 ^{2c}		39 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.80-2.02 (m, 2H), 2.12 (m, 1H), 2.20-2.36 (m, 3H), 2.40 (m, 1H), 2.57 (m, 1H), 3.35-3.55 (m, 5H), 3.91 (d, 1H), 4.59 (d, 1H), 6.02 (d, 1H), 6.38 (s, 1H), 7.06-7.23 (m, 3H), 7.38 (d, 1H), 7.43 (s, 1H), 7.61-7.77 (m, 2H), 8.47 (d, 1H). LRMS : m/z (TSP ⁺) 485.3, 487.3 [MH ⁺]

36 ^{1d}		62 white foam	¹ Hnmr (CD ₃ OD, 400MHz) δ : 2.17 (m, 2H), 2.21-2.40 (m, 4H), 2.53 (m, 2H), 2.78 (m, 2H), 2.92 (m, 1H), 3.06 (t, 1H), 3.17 (t, 2H), 3.26 (m, 4H), 3.83 (m, 2H), 4.04 (m, 2H), 4.20 (m, 1H), 4.48 (m, 1H), 7.44 (d, 1H), 7.60 (d, 1H), 7.68 (dd, 1H), 7.71 (s, 1H), 7.93 (d, 1H), 8.40 (dd, 1H), 8.59 (d, 1H). LRMS : m/z (TSP ⁺) 491.2, 493.2 [MH ⁺]
37 ^{1e}		52 white solid	¹ Hnmr (CD ₃ OD, 400MHz) δ : 1.60 (d, 3H), 2.04-2.33 (m, 3H), 2.37-2.43 (m, 3H), 2.61-2.83 (m, 2H), 4.15 (d, 1H), 4.28 (q, 1H), 4.40 (d, 1H), 7.24-7.40 (m, 6H), 7.46 (d, 1H), 7.58 (s, 1H), 7.61 (dd, 1H), 7.82 (d, 1H), 8.32 (dd, 1H), 8.57 (d, 1H). LRMS : m/z (TSP ⁺) 468.1, 470.1 [MH ⁺]

38 ^{II}		71 white solid	¹ Hnmr (CD ₃ OD, 400MHz) δ: 1.60 (d, 3H), 2.05-2.20 (m, 1H), 2.20-2.40 (m, 2H), 2.40-2.57 (m, 2H), 2.57-2.66 (m, 2H), 2.70-2.81 (m, 1H), 4.18 (d, 1H), 4.35 (m, 2H), 7.23-7.42 (m, 6H), 7.46 (d, 1H), 7.60 (s, 1H), 7.75 (dd, 1H), 7.65 (d, 1H), 8.50 (dd, 1H), 8.57 (d, 1H). LRMS : m/z (TSP ⁺) 468.1, 470.1 [MH ⁺] Microanalysis found: C, 53.31; H, 5.69; N, 6.82. C ₂₆ H ₂₇ Cl ₂ N ₃ O; 2HCl; 2.3H ₂ O requires C, 53.58; H, 5.81; N, 7.21%.
39 ^a			¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.00-2.27 (m, 6H), 2.37 (m, 1H), 2.35 (m, 1H), 2.52 (m, 2H), 2.70-3.24 (m, 6H), 3.55 (m, 1H), 3.68 (m, 1H), 3.81 (m, 2H), 4.03 (m, 1H), 4.21 (m, 2H), 4.42 (m, 1H), 7.43 (m, 1H), 7.60 (m, 1H), 7.70 (m, 1H), 7.81 (m, 1H), 8.07 (m, 1H), 8.60 (m, 1H). LRMS: m/z (TSP ⁺) 489.1, 491.2 [MH ⁺]

48

40 ^h		46 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.25 (m, 3H), 2.04-2.72 (m, 14H), 2.80 (m, 2H), 3.35 (m, 1H), 4.20 (m, 3H), 4.80 (m, 1H), 7.20 (m, 1H), 7.39 (m, 1H), 7.50 (m, 2H), 7.80 (m, 2H), 8.50 (m, 1H). LRMS : m/z (TSP ⁺) 504.1, 506.1 [MH ⁺]
-----------------	---	----------------------	---

1 = aldehyde hydrochloride was used


2 = isolated as the free base

Starting amines:

- 5 a = 2-(3-pyridinyloxy)ethylamine as prepared in WO 0071520
 b = 2-(2-pyridinyloxy)ethylamine as prepared in Tetrahedron, 44, 1988, 91
 c = 4-(aminomethyl)-1-methyl-2(1H)-pyridinone from preparation 21
 d = N-(3-aminopropyl)morpholine
 e = L-(-)-alpha-methylbenzylamine
- 10 f = R-(+)-1-phenylethylamine
 g = N-(3-azetidinylmethyl)-N-methylacetamide from preparation 25
 h = ethyl 4-piperidinecarboxylate



Examples 41 to 47



The following examples of general structure:





were prepared from the aldehyde from preparation 12a, and the appropriate
 5 amines, following a similar procedure to that described in example 32.

Example	R	Yield (%)	Data
41 ^a		53 white powder	¹ Hnmr (CD ₃ OD, 400MHz) δ: 1.58 (d, 3H), 2.07-2.28 (m, 3H), 2.35-2.50 (m, 3H), 2.60-2.75 (m, 5H), 4.17 (d, 1H), 4.20-4.32 (m, 2H), 7.22-7.39 (m, 6H), 7.48 (d, 1H), 7.53 (s, 1H), 7.67 (d, 1H), 7.76 (d, 1H), 8.44 (dd, 1H). LRMS : m/z (TSP ⁺) 482.1, 484.1 [MH ⁺] Microanalysis found: C, 53.87; H, 5.68; N, 6.82. C ₂₇ H ₂₉ Cl ₂ N ₃ O; 2HCl; 2.35H ₂ O requires C, 54.25; H, 6.02; N, 7.03%.

42 ^b		24 white foam	¹ Hnmr (CD ₃ OD, 400MHz) δ: 1.58 (m, 3H), 2.08 (m, 1H), 2.18-2.32 (m, 2H), 2.37-2.61 (m, 4H), 2.61-2.75 (m, 4H), 4.13 (d, 1H), 4.20-4.35 (m, 2H), 7.23-7.39 (m, 6H), 7.47 (d, 1H), 7.57 (s, 1H), 7.62 (d, 1H), 7.69 (d, 1H), 8.39 (dd, 1H). LRMS : m/z (TSP ⁺) 482.1, 484.0 [MH ⁺] Microanalysis found: C, 53.36; H, 5.95; N, 6.55. C ₂₇ H ₂₉ Cl ₂ N ₃ O; 2HCl; 3.0H ₂ O requires C, 53.24; H, 6.12; N, 6.90%.
43 ^c		26 white solid	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.11 (m, 2H), 2.20-2.37 (m, 4H), 2.41-2.56 (m, 2H), 2.68- 2.78 (m, 4H), 2.86 (m, 1H), 3.03 (t, 2H), 3.10 (t, 2H), 3.21 (m, 2H), 3.46 (m, 2H), 3.79 (t, 2H), 4.01 (m, 2H), 4.17 (d, 1H), 4.37 (d, 1H), 7.42 (d, 1H), 7.55-7.61 (m, 2H), 7.72 (m, 2H), 8.55 (dd, 1H). LRMS : m/z (ES ⁺) 505, 507 [MH ⁺]

44 ^{1d}		47 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.52 (m, 2H), 1.88 (s, 3H), 1.92-2.36 (m, 10H), 2.51-2.60 (m, 5H), 2.79 (m, 1H), 3.86 (d, 1H), 4.35 (bs, 1H), 4.68 (d, 1H), 7.00 (d, 1H), 7.21 (d, 1H), 7.41 (m, 2H), 7.56-7.63 (m, 2H). LRMS : m/z (TSP ⁺) 489.1, 491.1 [MH ⁺] Microanalysis found: C, 60.17; H, 6.19; N, 11.11. C ₂₅ H ₃₀ Cl ₂ N ₄ O ₂ requires C, 60.24; H, 6.27; N, 11.24%.
45 ^{1e}		55 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.53 (m, 2H), 1.91 (s, 3H), 1.98-2.43 (m, 10H), 2.48-2.66 (m, 5H), 2.84 (m, 1H), 3.92 (d, 1H), 4.38 (bs, 1H), 4.60 (d, 1H), 7.02 (d, 1H), 7.24 (d, 1H), 7.44 (m, 2H), 7.57-7.66 (m, 2H). LRMS : m/z (TSP ⁺) 489.2, 491.2 [MH ⁺] Microanalysis found: C, 59.52; H, 6.23; N, 10.82. C ₂₅ H ₃₀ Cl ₂ N ₄ O ₂ ·0.8H ₂ O requires C, 59.60; H, 6.32; N, 11.12%.

46 ^{1f}		51 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ (mixture of diastereoisomers): 1.55 (m, 2H), 1.63-2.33 (m, 12H), 2.37- 2.59 (m, 5H), 2.77, 2.86 (d, 3H, 60:40), 3.86 (2xd, 1H), 4.30, 5.16 (m, 1H, 40:60), 4.57 (2xd, 1H), 6.96 (d, 1H), 7.19 (d, 1H), 7.37 (d, 2H), 7.50-7.60 (m, 2H). LRMS : m/z (ES ⁺) 503, 505 [MH ⁺]
47 ^{1g}		66 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.58 (m, 2H), 1.95 (d, 2H), 2.08 (s, 3H), 2.12-2.40 (m, 8H), 2.58 (s, 3H), 3.40 (t, 2H), 3.56 (t, 2H), 3.92 (d, 1H), 4.71 (d, 1H), 7.01 (d, 1H), 7.23 (m, 1H), 7.43 (t, 2H), 7.61 (m, 2H). LRMS : m/z (TSP ⁺) 489.2, 491.2 [MH ⁺] Microanalysis found: C, 59.66; H, 6.29; N, 10.95. C ₂₅ H ₃₀ Cl ₂ N ₄ O ₂ ·0.7H ₂ O requires C, 59.81; H, 6.30; N, 11.16%.

1 = product isolated as the free base

Starting amines:

a = L-(-)-α-methylbenzylamine

b = R-(+)-1-phenylethylamine


c = N-(3-aminopropyl)morpholine

d = (3S)-(-)-3-acetamidopyrrolidine

e = (3R)-(+)-3-acetamidopyrrolidine

f = 3-(N-acetyl-N-methylamino)pyrrolidine

g = 1-acetypiperazine

Example 48**(5S)-5-(3,4-Dichlorophenyl)-5-(2-[(2-fluorobenzyl)amino]ethyl)-1-(2-pyridinyl)-2-piperidinone dihydrochloride**

- 5 2-Fluorobenzylamine (90 μ l, 0.79mmol) was added to a solution of the aldehyde hydrochloride from preparation 11b (250mg, 0.63mmol), in dichloromethane (5ml), and the solution stirred for 5 minutes. Sodium triacetoxyborohydride (132.6mg, 0.63mmol) was added and the reaction stirred for a further 10 minutes. Saturated aqueous sodium bicarbonate solution (3ml) was added, the mixture stirred for 10 minutes, then filtered through a phase separation filter, and the organic filtrate evaporated. The residue was purified by Biotage® column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as the eluant. The product was redissolved in dichloromethane, 1N ethereal hydrochloric acid added, and the solvents
- 10
- 15 evaporated under reduced pressure to afford the title compound as a white solid, 77.4mg.

$^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 2.10-2.36 (m, 3H), 2.36-2.53 (m, 2H), 2.60-2.80 (m, 2H), 2.92 (m, 1H), 4.13 (d, 1H), 4.18 (s, 2H), 4.42 (d, 1H), 7.12-7.23 (m, 2H), 7.37-7.50 (m, 3H), 7.54 (m, 2H), 7.67 (s, 1H), 7.79 (d, 1H), 8.20 (m, 1H), 8.55 (m, 1H).


20

LRMS : m/z (TSP $^+$) 472.1, 474.1 [MH^+]

Microanalysis found: C, 51.92; H, 5.17; N, 7.00. $\text{C}_{25}\text{H}_{24}\text{Cl}_2\text{FN}_3\text{O} \cdot 2\text{HCl} \cdot 1.6\text{H}_2\text{O}$ requires C, 52.30; H, 4.78; N, 7.32%.



Examples 49 to 53



The following compounds of general structure:



- 5 were prepared from the aldehyde hydrochloride from preparation 11b and the corresponding commercially available amine, following the procedure described in example 48.

Example	R	Yield (%)	Data
49 ^a		25	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.12-2.36 (m, 3H), 2.38-2.55 (m, 2H), 2.64-2.80 (m, 2H), 2.85 (m, 1H), 4.12 (d, 1H), 4.19 (s, 2H), 4.50 (d, 1H), 7.16-7.23 (m, 3H), 7.40 (m, 2H), 7.50 (m, 1H), 7.59 (d, 1H), 7.70 (d, 1H), 7.74 (d, 1H), 8.16 (m, 1H), 8.50 (m, 1H). LRMS : m/z (TSP ⁺) 472.1, 474.1 [MH ⁺] Microanalysis found: C, 52.42; H, 5.22; N, 7.03. C ₂₅ H ₂₄ Cl ₂ FN ₃ O; 2HCl; 1.5H ₂ O requires C, 52.46; H, 4.76; N, 7.34%

50 ^b		<p>31</p> <p>¹Hnmr (CD₃OD, 400MHz) δ: 2.17-2.40 (m, 3H), 2.40-2.57 (m, 2H), 2.65-2.82 (m, 2H), 2.91 (m, 1H), 4.09 (s, 2H), 4.20 (d, 1H), 4.41 (d, 1H), 7.10 (m, 2H), 7.32-7.49 (m, 3H), 7.57 (d, 1H), 7.66 (m, 2H), 7.91 (d, 1H), 8.41 (dd, 1H), 8.58 (d, 1H).</p> <p>LRMS : m/z (TSP⁺) 472.1, 474.2 [MH⁺]</p> <p>Microanalysis found: C, 51.12; H, 5.04; N, 6.76.</p> <p>C₂₅H₂₄Cl₂FN₃O.2HCl;2.5H₂O requires C, 50.86; H, 4.95; N, 7.12%</p>
51 ^c		<p>16</p> <p>¹Hnmr (CD₃OD, 400MHz) δ: 2.19-2.40 (m, 3H), 2.43-2.60 (m, 2H), 2.68-2.83 (m, 2H), 2.83-2.98 (m, 1H), 3.85 (s, 3H), 4.15 (s, 2H), 4.24 (d, 1H), 4.43 (d, 1H), 6.90-7.13 (m, 2H), 7.31 (d, 1H), 7.39-7.50 (m, 2H), 7.59 (d, 1H), 7.70 (s, 1H), 7.77 (dd, 1H), 8.00 (d, 1H), 8.52 (dd, 1H), 8.60 (d, 1H).</p> <p>LRMS : m/z (TSP⁺) 484.2, 486.2 [MH⁺]</p> <p>Microanalysis found: C, 51.60; H, 5.56; N, 6.72.</p> <p>C₂₆H₂₇Cl₂N₃O₂;2HCl;2.5H₂O requires C, 51.84; H, 5.69; N, 6.97%</p>

52 ^d	 <chem>CNCC1=CC=C(OC)C=C1</chem>	11 ¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.19-2.40 (m, 3H), 2.42-2.60 (m, 2H), 2.71-2.86 (m, 2H), 2.92 (m, 1H), 3.81 (s, 3H), 4.10 (s, 2H), 4.22 (d, 1H), 4.43 (d, 1H), 6.90-7.11 (m, 3H), 7.33 (dd, 1H), 7.42 (d, 1H), 7.59 (d, 1H), 7.69 (s, 1H), 7.75 (dd, 1H), 7.99 (d, 1H), 8.48 (dd, 1H), 8.60 (d, 1H). LRMS : m/z (TSP ⁺) 484.2, 486.1 [MH ⁺] Microanalysis found: C, 52.10; H, 5.46; N, 6.97. C ₂₈ H ₂₇ Cl ₂ N ₃ O ₂ ·2HCl·2.5H ₂ O requires C, 51.84; H, 5.69; N, 6.97%
53 ^e	 <chem>CNCC1=CC=C(OC)C=C1</chem>	8 ¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.13-2.40 (m, 3H), 2.40-2.58 (m, 2H), 2.65-2.82 (m, 2H), 2.82-2.95 (m, 1H), 3.80 (s, 3H), 4.05 (s, 2H), 4.20 (d, 1H), 4.44 (d, 1H), 6.93 (d, 2H), 7.32 (d, 2H), 7.42 (d, 1H), 7.59 (d, 1H), 7.63-7.76 (m, 2H), 7.92 (d, 1H), 8.40 (dd, 1H), 8.59 (d, 1H). LRMS : m/z (TSP ⁺) 484.2, 486.2 [MH ⁺] Microanalysis found: C, 51.14; H, 5.70; N, 6.65. C ₂₆ H ₂₇ Cl ₂ N ₃ O ₂ ·2HCl·3H ₂ O requires C, 51.07; H, 5.77; N, 6.91%

Starting amines:

a = 3-fluorobenzylamine

b = 4-fluorobenzylamine


c = 2-methoxybenzylamine

5 d = 3-methoxybenzylamine

e = 4-methoxybenzylamine

Example 54

10 *tert*-butyl 1-{2-[3-(3,4-dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-piperidinyldicarbamate



Triethylamine (0.15ml, 1.1mmol) was added to a solution of the aldehyde hydrochloride from preparation 11b (400mg, 1.0mmol) in dichloromethane (5ml), and the solution stirred for 5 minutes. *Tert*-butyl 4-piperidinyldicarbamate (240mg, 1.2mmol) and sodium triacetoxyborohydride (295mg, 1.4mmol) were then added, and the reaction stirred at room temperature for 3 days. Methanol was added, the mixture stirred for 15 minutes, then evaporated under reduced pressure. The residue was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant to give the title compound as a white foam, 580mg.


¹Hnmr (CDCl₃, 400MHz) δ : 1.28 (m, 2H), 1.39 (s, 9H), 1.78-2.00 (m, 7H), 2.11 (m, 2H), 2.27 (m, 2H), 2.56 (m, 1H), 2.62 (m, 2H), 3.36 (m, 1H), 3.88 (d, 1H), 4.31 (bs, 1H), 4.58 (d, 1H), 7.10 (dd, 1H), 7.17 (d, 1H), 7.36 (d, 1H), 7.43 (d, 1H), 7.67 (s, 2H), 8.45 (d, 1H).

LRMS : m/z (ES⁺) 547, 549 [MH⁺]

Microanalysis found: C, 71.17; H, 6.63; N, 10.23. C₂₈H₃₆Cl₂N₄O₃ requires C, 61.42; H, 6.63; N, 10.23%.

Example 55

(5S)-5-(3,4-Dichlorophenyl)-1-(2-pyridinyl)-5-[2-[(4-pyridinylmethyl)amino]ethyl]-2-piperidinone hydrochloride



- 5 The aldehyde from preparation 11a (192mg, 0.53mmol) and 4-(aminomethyl)-pyridine (54μl, 0.53mmol) in dichloromethane (10ml) were stirred together for 5 minutes, then acetic acid (61μl, 1.06mmol) and sodium triacetoxymethylborohydride (224mg, 1.06mmol) were added, and the reaction stirred at room temperature for 18 hours. The mixture was washed with aqueous 1N sodium hydroxide solution (5ml), dried (MgSO₄) and concentrated under reduced pressure. The residual oil was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as eluant, to give a sticky foam. This product was dissolved in dichloromethane (2ml), treated with ethereal hydrochloric acid and the mixture evaporated under reduced pressure to give the title compound as a white powder, 208mg.


¹Hnmr (CD₃OD, 400MHz) δ: 2.30-2.55 (m, 5H), 2.80 (m, 1H), 2.92 (m, 1H), 3.09 (m, 1H), 4.21 (d, 1H), 4.50 (d, 1H), 4.52 (s, 2H), 7.49 (dd, 1H), 7.61 (d, 1H), 7.67 (dd, 1H), 7.75 (s, 1H), 7.92 (d, 1H), 8.22 (d, 2H), 8.38 (dd, 1H), 8.59 (d, 1H), 8.95 (d, 2H).

- 20 LRMS : m/z (TSP⁺) 455.0, 457.1 [MH⁺]

Microanalysis found: C, 47.68; H, 5.07; N, 9.17.
C₂₄H₂₄Cl₂N₄O;3HCl;0.5H₂O;0.5CH₂Cl₂ requires C, 47.68; H, 5.07; N, 9.17%.

Examples 56 to 57


The compounds of the following general structure:



were prepared from the aldehyde from preparation 11a, and the appropriate
 5 amine, following a similar procedure to that described in example 55.

	R	Yield (%)	Data
56 ^{1a}		59 white solid	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.21-2.60 (m, 5H), 2.75-2.95 (m, 2H), 3.05 (m, 1H), 4.26 (d, 1H), 4.40 (s, 2H), 4.43 (d, 1H), 7.42 (d, 1H), 7.58 (d, 1H), 7.65 (d, 1H), 7.71 (m, 2H), 7.76 (m, 1H), 8.01 (d, 1H), 8.13 (dd, 1H), 8.56 (dd, 1H), 8.60 (d, 1H), 8.70 (d, 1H). LRMS : m/z (TSP ⁺) 455.0, 457.1 [MH ⁺] Microanalysis found: C, 46.29; H, 5.27; N, 8.80. C ₂₄ H ₂₄ Cl ₂ N ₄ O;3HCl;2H ₂ O requires C, 46.58; H, 5.05; N, 9.05%

60

57 ^b		58 yellow solid	¹ Hnmr (CD ₃ OD, 300MHz) δ: 2.20-2.40 (m, 5H), 2.79 (m, 2H), 2.97 (m, 1H), 3.59 (m, 4H), 4.23 (m, 1H), 4.48 (m, 1H), 7.48 (m, 1H), 7.81 (m, 1H), 7.70 (m, 1H), 7.76 (m, 1H), 7.96 (m, 1H), 8.10 (m, 2H), 8.41 (m, 1H), 8.60 (m, 1H), 8.82 (m, 2H). Microanalysis found: C, 46.46; H, 5.71; N, 8.78. C ₂₅ H ₂₈ Cl ₂ N ₄ O; 3HCl; 3H ₂ O; 0.2CH ₂ Cl ₂ requires C, 46.58; H, 5.49; N, 8.62%.
-----------------	---	-----------------------	---

1 = aldehyde hydrochloride from preparation 11b was used as the starting material

Starting amines:


a = 2-(aminoethyl)pyridine

b = 4-(2-aminoethyl)pyridine

5

Example 58

1-(2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl)-4-piperidinecarboxamide



10

Isonipecotamide (128mg, 1mmol) and glacial acetic acid (120mg, 2mmol) were added to a solution of the aldehyde from preparation 11a (400mg, 1mmol) in dichloromethane (10ml) and the solution stirred at room temperature for 30 minutes. Sodium triacetoxyborohydride (212mg, 2mmol) was added and the reaction stirred at room temperature for 4 hours. The reaction was quenched with methanol (10ml), the solution stirred for 10 minutes, then concentrated

15

under reduced pressure. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant to afford the title compound as a colourless foam, 162mg.


¹Hnmr (CDCl₃, 400MHz) δ: 1.74-2.17 (m, 11H), 2.26 (m, 2H), 2.55 (m, 1H), 2.75 (m, 2H), 3.42 (m, 1H), 3.86 (dd, 1H), 4.61 (dd, 1H), 5.21 (bs, 1H), 5.38 (bs, 1H), 7.08 (m, 1H), 7.19 (m, 1H), 7.36 (m, 1H), 7.42 (s, 1H), 7.62 (m, 2H), 8.42 (m, 1H).

LRMS : m/z (ES⁺) 475, 477 (MH⁺)

Microanalysis found: C, 57.95; H, 6.06; N, 11.02. C₂₄H₂₈Cl₂N₄O₂·H₂O requires C, 58.08; H, 6.06; N, 11.01%.

Example 59

(5S)-5-{2-[(1-Acetyl-4-piperidiny)amino]ethyl}-5-(3,4-dichlorophenyl)-1-(2-pyridinyl)-2-piperidinone



The title compound was prepared as a white foam in 71% yield, from the aldehyde from preparation 11a and the amine from preparation 22, following a similar procedure to that described in example 58.


¹Hnmr (CDCl₃, 400MHz) δ: 1.10 (t, 2H), 1.70 (s, 2H), 1.85 (m, 1H), 1.92 (m, 1H), 2.00 (s, 3H), 2.10 (m, 1H), 2.25 (m, 3H), 2.40-2.70 (m, 4H), 2.95 (t, 1H), 3.65 (d, 1H), 3.90 (d, 1H), 4.30 (t, 1H), 4.55 (dd, 1H), 7.10 (dd, 1H), 7.19 (d, 1H), 7.38 (d, 1H), 7.45 (d, 1H), 7.68 (d, 2H), 8.45 (d, 1H).

LRMS : m/z (TSP⁺) 489.1, 491.1 [MH⁺]

Microanalysis found: C, 58.91; H, 6.27; N, 10.85. C₂₅H₃₀Cl₂N₄O₂·0.3CH₂Cl₂ requires C, 59.07; H, 5.99; N, 10.88%.

Example 60

(5S)-5-(3,4-Dichlorophenyl)-5-(2-([2-(4-morpholinyl)ethyl]amino)ethyl)-1-(2-pyridinyl)-2-piperidinone trihydrochloride



- 5 2-(4-Morpholinyl)ethylamine (970mg, 7.5mmol), acetic acid (10 drops), and sodium triacetoxyborohydride (500mg, 2.4mmol) were added consecutively to a solution of the aldehyde from preparation 11a (270mg, 0.68mmol), and the reaction stirred at room temperature for an hour. The mixture was washed with 1N sodium hydroxide solution, then brine, dried (MgSO₄) and concentrated
- 10 under reduced pressure. The product was redissolved in dichloromethane, treated with 1N ethereal hydrochloric acid and the solution evaporated under reduced pressure to afford the title compound as a yellow solid, 220mg.

¹Hnmr (CD₃OD, 300MHz) δ: 2.35 (m, 3H), 2.54 (m, 2H), 2.83 (m, 2H), 2.98 (m, 1H), 3.23 (m, 2H), 3.42-3.66 (m, 6H), 3.87-4.12 (m, 4H), 4.23 (m, 1H), 4.48 (m, 1H), 7.48 (m, 1H), 7.61 (m, 1H), 7.75 (m, 2H), 7.98 (m, 1H), 8.47 (m, 1H), 8.61 (m, 1H).


LRMS : m/z (ES⁺) 477, 479 [MH⁺]

Microanalysis found: C, 44.94; H, 6.17; N, 8.72.

C₂₄H₃₀Cl₂N₄O₂·3HCl·2H₂O·0.25CH₂Cl₂ requires C, 45.22; H, 5.87; N, 8.70%.

Example 61


(5S)-5-(3,4-Dichlorophenyl)-5-[2-(4-methoxy-1-piperidinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone



- 5 4-Methoxypiperidine (WO 9847876) (81mg, 0.7mmol) was added to a solution of the aldehyde hydrochloride from preparation 11b (200mg, 0.5mmol) in dichloromethane (4ml) and triethylamine (100 μ l, 0.7mmol), and the solution stirred at room temperature for 10 minutes. Sodium triacetoxyborohydride (158mg, 1.5mmol) and acetic acid (6 μ l, 2mmol) were added and the reaction
- 10 stirred at room temperature for 2 hours. The reaction was quenched with methanol, and the solution concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (95:5:0 to 90:10:1) to give the title compound as a white foam, 182mg.
- 15 ¹Hnmr (CDCl₃, 400MHz) δ : 1.57 (s, 2H), 1.85-2.11 (m, 7H), 2.24 (m, 4H), 2.55 (m, 3H), 3.19 (s, 1H), 3.24 (s, 3H), 3.85 (d, 1H), 4.58 (d, 1H), 7.06 (m, 1H), 7.18 (d, 1H), 7.36 (d, 1H), 7.41 (s, 1H), 7.62 (s, 2H), 7.42 (d, 1H).
LRMS : m/z (ES⁺) 462, 464 [MH⁺]




Examples 62 to 72




The following examples of general structure:






were prepared from the aldehyde hydrochloride from preparation 11b and the appropriate amines, following a similar procedure to that described in example 61.

	R	Yield (%)	Data
62 ^a		39 foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.53 (m, 2H), 1.81 (m, 2H), 1.95 (m, 5H), 2.11 (m, 2H), 2.27 (m, 2H), 2.57 (m, 3H), 3.25 (m, 1H), 3.32 (s, 3H), 3.47 (d, 2H), 3.51 (d, 2H), 3.87 (d, 1H), 4.58 (d, 1H), 7.09 (dd, 1H), 7.18 (d, 1H), 7.35 (d, 1H), 7.43 (s, 1H), 7.65 (s, 2H), 8.44 (dd, 1H). LRMS : m/z (ES ⁺) 506, 508 [MH ⁺]
63 ^b		78 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.66 (m, 2H), 1.80 (m, 2H), 1.88-1.98 (m, 3H), 2.07 (m, 4H), 2.27 (m, 2H), 2.51 (m, 3H), 2.72 (s, 3H), 3.88 (d, 1H), 4.51 (s, 1H), 4.61 (d, 2H), 7.09 (dd, 1H), 7.18 (d, 1H), 7.36 (d, 1H), 7.44 (s, 1H), 7.66 (s, 2H), 8.44 (dd, 1H). Microanalysis found: C, 58.27; H, 6.02; N, 10.61. C ₂₅ H ₃₀ Cl ₂ N ₄ O ₃ ·0.15CH ₂ Cl ₂ requires C, 58.30; H, 5.89; N, 10.81%

64 ^c		49 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.30 (m, 2H), 1.80-2.00 (m, 10H), 2.14 (m, 2H), 2.30 (m, 2H), 2.58 (m, 1H), 2.69 (m, 2H), 3.73 (m, 1H), 3.92 (d, 1H), 4.59 (d, 1H), 5.26 (d, 1H), 7.15 (dd, 1H), 7.20 (d, 1H), 7.40 (d, 1H), 7.47 (s, 1H), 7.70 (s, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 489.1, 491.2 [MH ⁺]
65 ^d		48 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.20 (t, 3H), 1.52 (s, 4H), 1.88 (m, 4H), 1.98 (m, 1H), 2.10 (m, 2H), 2.26 (m, 2H), 2.54 (m, 1H), 2.69 (s, 3H), 2.76 (m, 2H), 3.76 (bs, 1H), 3.90 (d, 1H), 4.08 (q, 2H), 4.60 (d, 1H), 7.09 (dd, 1H), 7.18 (d, 1H), 7.36 (d, 1H), 7.44 (s, 1H), 7.67 (s, 2H), 8.45 (dd, 1H). Microanalysis found: C, 60.08; H, 6.48; N, 10.21. C ₂₇ H ₃₄ Cl ₂ N ₄ O ₃ ·0.1CH ₂ Cl ₂ requires C, 60.06; H, 6.36; N, 10.34%.
66 ^a		48 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.46 (m, 2H), 1.64 (m, 2H), 1.77 (m, 7H), 1.97 (m, 4H), 2.13 (m, 2H), 2.30 (m, 2H), 2.54 (m, 4H), 2.76 (dd, 2H), 3.92 (d, 1H), 4.56 (d, 1H), 7.13 (dd, 1H), 7.26 (d, 1H), 7.39 (d, 1H), 7.48 (s, 1H), 7.68 (m, 2H), 8.48 (d, 1H). LRMS : m/z (ES ⁺) 501, 503 [MH ⁺]

67 ^f		42 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.57 (m, 3H), 1.90-1.98 (m, 8H), 2.13 (m, 2H), 2.31 (m, 2H), 2.37 (m, 2H), 2.58 (m, 1H), 2.79 (m, 2H), 3.30 (t, 2H), 3.90 (m, 1H), 3.93 (d, 1H), 4.60 (d, 1H), 7.13 (m, 1H), 7.23 (d, 1H), 7.40 (d, 1H), 7.47 (s, 1H), 7.70 (s, 2H), 8.48 (d, 1H). LRMS : m/z (ES ⁺) 515, 517 [MH ⁺]
68 ^g		51 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.50 (m, 2H), 1.80-2.00 (m, 5H), 2.10 (m, 2H), 2.20-2.40 (m, 4H), 2.56 (m, 1H), 2.80 (m, 2H), 3.80 (m, 1H), 3.86 (s, 2H), 3.90 (d, 1H), 4.60 (d, 1H), 5.15 (s, 1H), 7.10 (m, 1H), 7.20 (d, 1H), 7.36 (d, 1H), 7.45 (s, 1H), 7.67 (d, 2H), 8.46 (d, 1H). LRMS : m/z (ES ⁺) 530, 532 [MH ⁺]
69 ^h		63 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.63 (t, 4H), 1.88 (m, 2H), 2.00 (m, 1H), 2.11 (m, 2H), 2.22-2.40 (m, 6H), 2.53 (m, 1H), 3.88 (m, 5H), 4.62 (d, 1H), 7.09 (dd, 1H), 7.18 (d, 1H), 7.37 (d, 1H), 7.43 (s, 1H), 7.65 (s, 2H), 8.43 (d, 1H). LRMS : m/z (TSP ⁺) 490.2, 492.2 [MH ⁺]

70 ⁱ		30 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.61 (m, 2H), 1.66 (m, 2H), 1.96 (m, 5H), 2.12 (m, 2H), 2.27 (m, 2H), 2.57 (m, 1H), 2.88 (m, 3H), 3.75 (s, 3H), 3.94 (d, 1H), 4.64 (d, 1H), 6.80 (d, 1H), 6.91 (m, 1H), 7.10 (m, 3H), 7.23 (m, 1H), 7.38 (d, 1H), 7.47 (s, 1H), 7.68 (d, 2H), 8.47 (m, 1H). LRMS : m/z (ES ⁺) 538, 540 [MH ⁺] Microanalysis found: C, 66.09; H, 6.20; N, 7.67. C ₃₀ H ₃₃ Cl ₂ N ₃ O ₂ ·0.1CH ₂ Cl ₂ requires C, 66.09; H, 6.12; N, 7.68%.
71 ⁱ		84 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.62 (m, 4H), 1.97 (m, 4H), 2.15 (m, 2H), 2.29 (m, 3H), 2.56 (dd, 1H), 2.73 (bs, 1H), 3.07 (m, 2H), 3.94 (d, 1H), 4.63 (d, 1H), 7.12 (m, 3H), 7.23 (d, 1H), 7.41 (d, 1H), 7.48 (s, 1H), 7.59 (dd, 1H), 7.69 (m, 2H), 8.45 (m, 2H). LRMS : m/z (ES ⁺) 509, 511 [MH ⁺]
72 ^k		97 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.95 (m, 2H), 2.04 (m, 1H), 2.15 (m, 2H), 2.28 (m, 2H), 2.38 (m, 4H), 2.58 (m, 1H), 3.44 (m, 4H), 3.92 (d, 1H), 4.63 (d, 1H), 6.58 (d, 2H), 7.09 (m, 1H), 7.20 (d, 1H), 7.38 (d, 1H), 7.42 (m, 1H), 7.46 (s, 1H), 7.66 (s, 2H), 8.12 (d, 1H), 8.43 (d, 1H). LRMS : m/z (ES ⁺) 510, 512 [MH ⁺]

Starting amines:


a = 4-(2-methoxyethoxy)piperidine hydrochloride as prepared in US 4237138

68

- b = 4-piperidinyl methylcarbamate from preparation 62
 c = N-(4-piperidinyl)acetamide as prepared in EP 908452
 d = Ethyl methyl(4-piperidinyl)carbamate as prepared in FR 2321890
 e = 4-(1-pyrrolidinyl)piperidine
 5 f = 1-(4-piperidinyl)-2-pyrrolidinone as prepared in WO 9410146
 g = 3-(4-piperidinyl)-2,4-imidazolidinedione from preparation 67
 h = 1,4-dioxo-8-azaspiro[4.5]decane
 i = 4-(2-methoxyphenyl)piperidine
 j = 2-(4'-piperidinyl)pyridine as prepared in EP 630887
 10 k = 1 (2-pyridinyl)piperazine

Example 73

(5S)-5-(3,4-Dichlorophenyl)-5-{2-[3-(4-hydroxy-1-piperidinyl)-1-azetidiny]ethyl}-
1-(2-pyridinyl)-2-piperidinone




15

Triethylamine (1ml, 7.2mmol) and acetic acid (1.1ml, 18.3mmol) were added to a solution of 1-(3-azetidiny)-4-piperidinol trifluoroacetate (WO 9605193) (250mg, 0.93mmol), the aldehyde from preparation 11a (250mg, 0.62mmol) and sodium triacetoxyborohydride (250mg, 1.2mmol) in dichloromethane (100ml) and the reaction stirred at room temperature for 90 minutes. The solution was washed with water, dried (MgSO₄), and concentrated under reduced pressure. The residual gum was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol (100:0 to 95:5) to afford the title compound as a yellow solid, 82mg.

25 ¹Hnmr (CDCl₃, 400MHz) δ: 1.55-2.40 (m, 14H), 2.60 (m, 3H), 2.80 (m, 2H), 2.95 (m, 1H), 3.52 (m, 2H), 3.75 (m, 1H), 3.95 (d, 1H), 4.58 (d, 1H), 7.20 (m, 2H), 7.40 (dd, 2H), 7.74 (m, 2H), 8.50 (d, 1H).
 LRMS : m/z (TSP⁺) 503.1, 505.1 [MH⁺]




Examples 74 to 85




The following examples of general structure:








were prepared from the aldehyde from preparation 11a and the appropriate
 5 amines, following a similar procedure to that described in example 73.


Example	R	Yield (%)	Data
74 ^a		43 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.25 (s, 3H), 1.50-2.00 (m, 8H), 2.18 (m, 4H), 2.34 (m, 4H), 2.60 (m, 1H), 2.75 (m, 2H), 2.95 (m, 1H), 3.44 (t, 2H), 3.90 (d, 1H), 4.48 (d, 1H), 6.55 (d, 1H), 7.20 (m, 2H), 7.42 (d, 1H), 7.45 (d, 1H), 7.72 (d, 1H), 8.50 (d, 1H). LRMS : m/z (TSP ⁺) 517.9, 519.9 [MH ⁺] Microanalysis found: C, 60.41; H, 6.75; N, 10.20. C ₂₇ H ₃₄ Cl ₂ N ₄ O ₂ ·0.2CH ₂ Cl ₂ ·0.4H ₂ O requires C, 60.31; H, 6.55; N, 10.34%.

75 ^b		60 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.70 (m, 1H), 1.82 (m, 1H), 2.14 (m, 2H), 2.30 (m, 7H), 2.58 (m, 1H), 2.72 (m, 2H), 2.90 (m, 1H), 3.40 (t, 2H), 3.68 (m, 4H), 3.90 (d, 1H), 4.56 (d, 1H), 7.16 (dd, 1H), 7.20 (dd, 1H), 7.40 (d, 1H), 7.44 (s, 1H), 7.70 (m, 2H), 8.50 (d, 1H). LRMS : m/z (TSP ⁺) 489.1, 491.1 [MH ⁺]
76 ^{1c}		78 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.78 (m, 1H), 1.86 (m, 1H), 2.10-2.40 (m, 5H), 2.57 (m, 1H), 3.25 (t, 2H), 3.50 (t, 2H), 3.95 (d, 1H), 4.63 (d, 1H), 7.15 (m, 1H), 7.25 (m, 2H), 7.35 (m, 2H), 7.40 (m, 1H), 7.45 (m, 3H), 7.70 (d, 2H), 8.48 (d, 1H).
77 ^{1d}		32 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.80 (m, 1H), 1.92 (m, 1H), 2.08-2.45 (m, 5H), 2.59 (m, 1H), 3.42 (m, 2H), 3.56 (m, 2H), 3.84 (s, 3H), 3.90 (d, 1H), 4.58 (d, 1H), 6.88 (d, 1H), 6.95 (dd, 1H), 7.14 (m, 1H), 7.20 (d, 2H), 7.24 (m, 2H), 7.39 (d, 1H), 7.44 (s, 1H), 7.70 (m, 2H), 8.45 (d, 1H). LRMS : m/z (TSP ⁺) 526.1, 528.1 [MH ⁺]

78 ^{1e}		29 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.70 (m, 1H), 1.81 (m, 1H), 2.10 (m, 2H), 2.27 (m, 4H), 2.38 (m, 4H), 2.58 (m, 1H), 2.70 (m, 3H), 2.77 (s, 3H), 2.96 (m, 1H), 3.21 (m, 2H), 3.38 (m, 2H), 3.90 (d, 1H), 4.57 (d, 1H), 7.16 (dd, 1H), 7.19 (d, 1H), 7.41 (d, 1H), 7.42 (s, 1H), 7.70 (m, 2H), 8.49 (d, 1H). LRMS : m/z (TSP ⁺) 566.3 [MH ⁺]
79 ⁱ		25 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.80 (m, 1H), 2.0-2.65 (m, 12H), 2.80 (m, 1H), 3.04 (m, 1H), 3.96 (d, 1H), 4.36 (s, 1H), 4.72 (d, 1H), 7.16 (dd, 1H), 7.24 (d, 1H), 7.44 (d, 1H), 7.50 (s, 1H), 7.75 (m, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 434.1, 436.1 [MH ⁺] Microanalysis found: C, 57.10; H, 5.73; N, 8.64. C ₂₂ H ₂₅ Cl ₂ N ₃ O ₂ ·0.4CH ₂ Cl ₂ ·0.25H ₂ O requires C, 56.90; H, 5.61; N, 8.89%.
80 ^g		8	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.62 (d, 2H), 1.93 (m, 5H), 2.18 (m, 5H), 2.30 (m, 2H), 2.62 (m, 3H), 3.94 (d, 1H), 4.74 (d, 1H), 7.16 (m, 1H), 7.22 (d, 1H), 7.41 (d, 1H), 7.48 (s, 1H), 7.72 (s, 2H), 8.48 (d, 1H). LRMS : m/z (ES ⁺) 516, 518 [MH ⁺] Microanalysis found: C, 55.52; H, 5.13; N, 7.88. C ₂₄ H ₂₆ Cl ₂ F ₃ N ₃ O ₂ requires C, 55.45; H, 5.11; N, 7.92%.

81 ⁿ		18 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.50-2.70 (m, 24H), 2.88 (m, 4H), 3.75 (m, 1H), 3.91 (d, 1H), 4.64 (d, 1H), 7.16 (m, 1H), 7.24 (dd, 1H), 7.42 (d, 1H), 7.50 (s, 1H), 7.72 (s, 2H), 8.50 (d, 1H). LRMS : m/z (TSP ⁺) 531.2, 533.2 [MH ⁺]
82 ⁱ		53 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.26 (m, 2H), 1.44 (m, 2H), 1.85-2.08 (m, 5H), 2.15 (m, 2H), 2.26 (m, 2H), 2.58 (m, 1H), 2.85 (m, 2H), 3.41 (m, 1H), 3.95 (d, 1H), 4.61 (d, 1H), 7.08 (m, 2H), 7.18 (m, 3H), 7.38 (d, 1H), 7.46 (s, 1H), 7.64 (m, 2H), 8.19 (d, 1H), 8.42 (d, 1H). LRMS : m/z (TSP ⁺) 525.1, 527.2 [MH ⁺]
83 ^{II}		16 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.58 (m, 2H), 1.75 (m, 2H), 1.81-2.08 (m, 5H), 2.15 (m, 2H), 2.24-2.40 (m, 3H), 2.58 (m, 1H), 2.84 (m, 2H), 3.90 (d, 1H), 4.60 (d, 1H), 7.02-7.20 (m, 4H), 7.38 (d, 1H), 7.45 (m, 1H), 7.68 (m, 2H), 8.02 (m, 2H), 8.45 (m, 1H).

84 ^{1k}		33 white crystal	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.58 (m, 2H), 1.72 (m, 2H), 1.83-2.08 (m, 5H), 2.16 (m, 2H), 2.28 (m, 2H), 2.40 (m, 1H), 2.55 (m, 1H), 2.85 (m, 2H), 3.90 (d, 1H), 4.60 (d, 1H), 7.02 (d, 2H), 7.10 (m, 1H), 7.20 (m, 1H), 7.38 (d, 1H), 7.42 (m, 1H), 7.66 (m, 2H), 8.08 (d, 2H), 8.42 (m, 1H). LRMS : m/z (TSP ⁺) 525.3, 527.3 [MH ⁺]
85 ^{1l}		37 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.94 (m, 2H), 2.04 (m, 1H), 2.14 (m, 2H), 2.30 (m, 2H), 2.42 (m, 4H), 2.58 (m, 1H), 3.17 (m, 4H), 2.90 (d, 1H), 4.69 (d, 1H), 5.62 (s, 1H), 7.12 (m, 1H), 7.10 (m, 1H), 7.20 (m, 1H), 7.39 (d, 1H), 7.45 (s, 1H), 7.68 (m, 2H), 8.24 (d, 1H), 8.34 (m, 2H), 8.45 (d, 1H). LRMS : m/z (TSP ⁺) 553.2, 555.2 [MH ⁺] Microanalysis found: C, 59.34; H, 5.58; N, 14.52. C ₂₈ H ₃₀ Cl ₂ N ₆ O ₂ ·0.73H ₂ O requires C, 59.35; H, 5.60; N, 14.83%.

86 ^m		30 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.73-1.98 (m, 6H), 2.04 (2xs, 3H), 2.17 (m, 2H), 2.28 (m, 2H), 2.38-2.62 (m, 4H), 3.38-3.58 (m, 4H), 3.92 (d, 1H), 4.66 (m, 1H), 7.17 (m, 1H), 7.22 (m, 1H), 7.41 (m, 1H), 7.46 (s, 1H), 7.72 (m, 2H), 8.46 (d, 1H). LRMS : m/z (TSP ⁺) 489.1, 491.1 [MH ⁺]
-----------------	---	----------------------	--

1 = the aldehyde hydrochloride from preparation 11b was used


2 = no triethylamine was used in the reaction

Starting amines:

- 5 a = 1-(3-azetidiny)-4-methyl-4-piperidinol trifluoroacetate from preparation 28
b = 3-morpholinoazetidine dihydrochloride as prepared in WO 9725322
c = 3-hydroxy-3-phenylazetidine hydrochloride as prepared in J.A.C.S; 1972; 94(8); 2758
d = 3-hydroxy-3-(2-methoxyphenyl)azetidine hydrochloride from preparation 30
e = 1-(3-azetidiny)-4-(methylsulphonyl)piperazine trifluoroacetate as prepared in WO 9725322
- 10 f = (S)-3-pyrrolidinol
g = 4-trifluoromethylpiperidinol trifluoroacetate from preparation 58
h = 1-(piperidin-4-yl)-4-piperidinol from preparation 64
i = 2-(4-piperidinyl)pyridine 1-oxide as prepared in WO 0037026
j = 3-(4-piperidinyl)pyridine 1-oxide dihydrochloride as prepared in preparation 74
- 45 k = 4-(4-piperidinyl)pyridine 1-oxide dihydrochloride as prepared in preparation 73
l = 2-(1-piperazinyl)nicotinamide hydrochloride as prepared in J.Med.Chem. 1983; 26(12);1696
m = N-acetylhomopiperazine

Example 87

(5S)-5-(3,4-Dichlorophenyl)-5-[2-[4-hydroxy-4-(trifluoromethyl)-1-piperidinyl]ethyl]-1-(6-methyl-2-pyridinyl)-2-piperidinone



- 5 The title compound was obtained as a solid, from the aldehyde from preparation 12a, and the amine from preparation 58, following a similar procedure to that described in example 73.

¹Hnmr (CDCl₃, 400MHz) δ: 1.62 (m, 4H), 1.82 (m, 2H), 1.94 (t, 2H), 2.02-2.18 (m, 5H), 2.30 (m, 2H), 2.50 (s, 3H), 2.58 (d, 1H), 2.68 (d, 1H), 3.90 (d, 1H), 4.42 (d, 1H), 6.98 (d, 1H), 7.22 (d, 1H), 7.41 (d, 2H), 7.58 (d, 2H).


LRMS : m/z (TSP⁺) 530.1, 532.1 [MH⁺]

Microanalysis found: C, 55.39; H, 5.48; N, 7.58. C₂₅H₂₈Cl₂F₃N₃O₂·0.2CH₂Cl₂ requires C, 55.29; H, 5.23; N, 7.68%.

15

Example 88

N-(1-[2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinylethyl]-4-phenyl-4-piperidinyl]-N-methylacetamide



Triethylamine (2ml) and glacial acetic acid (2ml) were added to a solution of N-methyl-N-(4-phenyl-4-piperidiny)acetamide (WO 9805640) (97.9mg, 0.36mmol) and the aldehyde hydrochloride from preparation 11b (145mg, 0.36mmol) in dichloromethane (20ml). Sodium triacetoxymorohydride (79mg, 0.37mmol) was added and the reaction stirred at room temperature for 18 hours. The reaction was neutralised using saturated sodium bicarbonate solution, the phases separated and the organic layer washed with brine, dried (MgSO₄), and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as eluant, and azeotroped with diethyl ether to afford the title compound as a white foam, 92mg.


¹Hnmr (CDCl₃, 400MHz) δ: 1.86-2.21 (m, 15H), 2.21-2.37 (m, 2H), 2.60 (m, 3H), 2.81 (s, 2H), 3.93 (d, 1H), 4.68 (d, 1H), 7.13 (m, 1H), 7.20-7.38 (m, 6H), 7.40 (d, 1H), 7.49 (s, 1H), 7.71 (d, 2H), 8.48 (d, 1H).

LRMS : m/z (TSP⁺) 579.1, 581.1 [MH⁺]



Microanalysis found: C, 63.57; H, 6.72; N, 9.10. C₃₂H₃₆Cl₂N₄O₂·1.3H₂O requires C, 63.74; H, 6.45; N, 9.29%.



20 Examples 89 to 94



The following examples of general structure:



were prepared from the aldehyde hydrochloride from preparation 11b and the appropriate amine, following a similar procedure to that described in example 88.

Example	R	Yield (%)	Data
89 ^a		62 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.43 (m, 2H), 1.77 (m, 4H), 1.97 (m, 3H), 2.12 (m, 3H), 2.34 (m, 2H), 2.52 (m, 4H), 2.60 (m, 1H), 2.81 (m, 2H), 3.71 (m, 4H), 3.95 (d, 1H), 4.63 (d, 1H), 7.14 (m, 1H), 7.21 (d, 1H), 7.41 (d, 1H), 7.50 (s, 1H), 7.72 (m, 2H), 8.51 (d, 1H). Microanalysis found: C, 61.67; H, 6.70; N, 10.63. C ₂₇ H ₃₄ Cl ₂ N ₄ O ₂ ·0.5H ₂ O requires C, 61.59; H, 6.70; N, 10.64%.
90 ^{1b}		19 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.83-2.18 (m, 12H), 2.21-2.36 (m, 4H), 2.49-2.67 (m, 3H), 3.91 (d, 1H), 4.68 (d, 1H), 5.40 (bs, 1H), 7.10 (m, 1H), 7.15-7.32 (m, 6H), 7.39 (d, 1H), 7.45 (s, 1H), 7.66 (m, 2H), 8.42 (d, 1H). LRMS : m/z (TSP ⁺) 565.2, 567.2 [MH ⁺] Microanalysis found: C, 63.41; H, 6.25; N, 9.54. C ₃₁ H ₃₄ Cl ₂ N ₄ O ₂ ·1.2H ₂ O requires C, 63.24; H, 6.01; N, 9.27%.

91 ^{2c}	 <chem>NC(=O)C1(CCN1)c2ccccc2</chem>	62 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.59 (m, 2H), 1.80-2.70 (m, 14H), 3.92 (d, 1H), 4.72 (d, 1H), 5.20 (bs, 2H), 7.15 (m, 1H), 7.24 (m, 3H), 7.33-7.46 (m, 4H), 7.49 (s, 1H), 7.70 (m, 2H), 8.44 (d, 1H). LRMS : m/z (TSP*) 551.1, 553.1 [MH ⁺] Microanalysis found: C, 63.28; H, 6.04; N, 9.41. C ₃₀ H ₃₂ Cl ₂ N ₄ O ₂ ·1.1H ₂ O requires C, 63.07; H, 6.03; N, 9.81%.
92 ^{2d}	 <chem>Oc1ccncc1C2(CCN2)</chem>	58 off- white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.62 (m, 2H), 1.90-2.50 (m, 12H), 2.55-2.82 (m, 3H), 4.40 (d, 1H), 4.69 (d, 1H), 7.05-7.29 (m, 3H), 7.37 (d, 1H), 7.42 (d, 1H), 7.53 (s, 1H), 7.71 (m, 3H), 8.50 (m, 2H). LRMS : m/z (TSP*) 525.1, 527.1 [MH ⁺] Microanalysis found: C, 62.77; H, 5.97; N, 10.19. C ₂₈ H ₃₀ Cl ₂ N ₄ O ₃ ·0.5H ₂ O requires C, 62.92; H, 5.85; N, 10.18%.

93 ^a		14	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.28 (t, 3H), 1.84-2.46 (m, 12H), 2.46-2.73 (m, 4H), 3.19 (s, 2H), 3.96 (d, 1H), 4.19 (q, 2H), 4.66 (d, 1H), 7.16 (m, 1H), 7.24 (d, 1H), 7.42 (d, 1H), 7.49 (s, 1H), 7.71 (m, 2H), 8.50 (d, 1H). LRMS : m/z (TSP ⁺) 519.2, 521.3 [MH ⁺]
94 ^f		36 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.82-2.70 (m, 18H), 3.59 (t, 2H), 3.98 (d, 1H), 4.66 (d, 1H), 7.16 (m, 1H), 7.22 (d, 1H), 7.41 (d, 1H), 7.51 (s, 1H), 7.72 (m, 2H), 8.50 (d, 1H). Microanalysis found: C, 58.37; H, 6.40; N, 10.95. C ₂₄ H ₃₀ Cl ₂ N ₄ O ₂ ·0.9H ₂ O requires C, 58.39; H, 6.40; N, 11.35%.

1 = No triethylamine was added to the reaction mixture.


2 = Tetrahydrofuran was used as the reaction solvent

Starting amines:

- 5 a = 4-(4-(4-piperidinyl)morpholine hydrochloride from preparation 65
 b = N-methyl-N-(4-phenyl-4-piperidinyl)acetamide as prepared in WO 9805640
 c = 4-phenyl-4-piperidinecarboxamide as prepared in WO 9426735
 d = 4-(2-pyridinyl)-4-piperidinol as prepared in DE 2630152
 e = ethyl 1-piperazinylacetate
 10 f = 2-(1-piperazinyl)ethanol


Examples 95 to 96

The following examples of general structure:



were prepared from the aldehyde hydrochloride from preparation 12b and the
 5 appropriate amine, following a similar procedure to that described in example 88, isolating the compounds after azeotroping with diethyl ether.

Example	R	Yield (%)	Data
95 ^a		37 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.98 (m, 2H), 2.07-2.41 (m, 5H), 2.41-2.66 (m, 8H), 2.86 (s, 3H), 3.10 (t, 4H), 3.92 (d, 1H), 4.32 (d, 2H), 4.77 (d, 1H), 5.69 (m, 1H), 6.98-7.06 (m, 2H), 7.29 (d, 1H), 7.44 (m, 2H), 7.57-7.66 (m, 3H), 8.31 (d, 1H). LRMS : m/z (TSP ⁺) 631.3, 633.3 [MH ⁺] Microanalysis found: C, 56.45; H, 6.05; N, 12.41. C ₃₀ H ₃₆ Cl ₂ N ₆ O ₃ S; 0.5(CH ₃ CH ₂) ₂ O; 0.5 H ₂ O requires C, 56.71; H, 6.25; N, 12.40%.

96 ^b		36 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.98 (m, 2H), 2.14-2.30 (m, 10H), 2.30-2.41 (m, 2H), 2.41-2.65 (m, 7H), 3.26 (m, 4H), 3.35 (s, 2H), 3.92 (d, 1H), 4.77 (d, 1H), 6.91 (dd, 1H), 7.01 (d, 1H), 7.31 (d, 1H), 7.42 (d, 2H), 7.56-7.72 (m, 3H), 8.20 (d, 1H). LRMS : m/z (TSP*) 581.1, 583.2 [MH ⁺] Microanalysis found: C, 63.22; H, 6.61; N, 13.72. C ₃₁ H ₃₈ Cl ₂ N ₆ O; 0.1(CH ₃ CH ₂) ₂ O; 0.5H ₂ O requires C, 63.07; H, 6.74; N, 14.05%.
-----------------	---	---------------------	---


Starting amines;

a = N-([2-(1-piperazinyl)-3-pyridinyl]methyl)methanesulphonamide dihydrochloride as prepared in preparation 80

5 b = N,N-dimethyl[2-(1-piperazinyl)-3-pyridyl]methanamine trihydrochloride as prepared in preparation 79

Example 97

10 (5S)-5-(3,4-Dichlorophenyl)-1-[6-(dimethylamino)-2-pyridinyl]-5-[2-[3-(4-morpholinyl)-1-azetidinyl]ethyl]-2-piperidinone



The title compound was obtained as a white foam in 74% yield, after trituration from diethyl ether, from the aldehyde from preparation 17 and 3-morpholinoazetidine dihydrochloride (WO 9725322), following a similar procedure to that described in example 88.


- 5 $^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.63 (m, 1H), 1.79 (m, 1H), 1.98-2.35 (m, 9H), 2.41-2.55 (m, 1H), 2.60-2.73 (m, 2H), 2.83 (t, 1H), 3.06 (s, 6H), 3.24 (q, 2H), 3.63 (s, 4H), 3.70 (d, 1H), 4.59 (d, 1H), 6.32 (d, 1H), 6.81 (d, 1H), 7.28 (d, 1H), 7.36 (d, 1H), 7.43 (dd, 1H), 7.57 (s, 1H).

LRMS : m/z (TSP^+) 532.2, 534.3 [MH^+]

- 10 Microanalysis found: C, 60.06; H, 6.69; N, 12.85. $\text{C}_{27}\text{H}_{35}\text{Cl}_2\text{N}_5\text{O}_2 \cdot 0.35\text{H}_2\text{O}$ requires C, 60.19; H, 6.68; N, 13.00%.

Example 98

- 15 *N*-[1-(2-{(3*S*)-3-(3,4-Dichlorophenyl)-1-[6-(dimethylamino)-2-pyridinyl]-6-oxopiperidinyl}ethyl)-4-piperidinyl]-*N*-methylacetamide



- 20 The title compound was obtained as a white foam in 67% yield, from the aldehyde from preparation 17 and the amine from preparation 60, following the procedure described in example 88.


$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.42-1.60 (m, 3H), 1.69 (m, 1H), 1.77-2.29 (m, 12H), 2.55 (m, 1H), 2.72-2.82 (m, 5H), 3.10 (s, 6H), 4.39 (m, 1H), 4.63 (d, 1H), 6.37 (d, 2H), 6.84 (d, 1H), 7.29-7.42 (m, 2H), 7.48 (dd, 1H), 7.61 (s, 1H).

LRMS : m/z (TSP^+) 546.3, 548.3 [MH^+]

- 25 Microanalysis found: C, 60.20; H, 6.93; N, 12.40. $\text{C}_{28}\text{H}_{37}\text{Cl}_2\text{N}_5\text{O}_2 \cdot 0.5\text{H}_2\text{O}$ requires C, 60.54; H, 6.89; N, 12.61%.

Example 99

tert-Butyl 1-(2-[(3*S*)-3-(3,4-dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl)-4-piperidinyl(methyl)carbamate



Triethylamine (1.5ml, 10.8mmol) was added to a suspension of the aldehyde hydrochloride from preparation 11b (500mg, 1.25mmol) and *tert*-butyl methyl(4-piperidinyl)carbamate (EP 457686) (402mg, 1.88mmol) in dichloromethane (250ml), and the mixture stirred at room temperature for 5 minutes. Acetic acid (1.5ml, 26.2mmol) and sodium triacetoxyborohydride (530mg, 2.5mmol) were added and the reaction stirred at room temperature for 2 hours. The mixture was washed with 2N sodium hydroxide solution (200ml), and the aqueous wash extracted with dichloromethane (2x200ml). The combined organic solutions were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. The crude product was purified by Biotage® column chromatography on silica gel using an elution gradient of dichloromethane:methanol (100:0 to 95:5) to afford the title compound as a white foam, 360mg.


¹Hnmr (CDCl₃, 300MHz) δ: 1.45 (s, 9H), 1.58 (m, 6H), 1.81-2.21 (m, 6H), 2.32 (m, 2H), 2.57-2.64 (m, 1H), 2.70 (s, 3H), 2.80 (m, 2H), 3.95 (d, 1H), 4.64 (d, 1H), 7.16 (m, 1H), 7.23 (m, 1H), 7.41 (m, 1H), 7.50 (s, 1H), 7.72 (m, 2H), 8.50 (m, 1H).

LRMS : m/z (TSP⁺) 561.2, 563.2 [MH⁺]

Microanalysis found: C, 61.50; H, 6.87; N, 9.83. C₂₉H₃₈Cl₂N₄O₃ requires C, 62.03; H, 6.77; N, 9.98%.




Examples 100 to 111




The following examples of general structure:








- were prepared from the aldehyde hydrochloride from preparation 12b and the appropriate amine, following a similar procedure to that described in example 99, except the products were purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia as eluant.

Example	R	Yield (%)	Data
100 ^a		28 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.75 (m, 1H), 1.82 (m, 1H), 2.06-2.18 (m, 2H), 2.20-2.08 (m, 2H), 2.14 (m, 3H), 2.54 (m, 3H), 2.79 (m, 2H), 2.98 (m, 1H), 3.03 (m, 5H), 3.16 (m, 2H), 3.44 (m, 2H), 3.82 (d, 1H), 4.46 (d, 1H), 6.97 (d, 1H), 7.19 (d, 1H), 7.38 (m, 2H), 7.55 (m, 2H). LRMS : m/z (TSP ⁺) 502.8, 504.8 [MH ⁺]

101 ^b		27 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.56 (m, 3H), 1.70 (m, 3H), 1.92 (m, 4H), 2.03 (s, 3H), 2.16 (m, 2H), 2.30 (m, 2H), 2.55 (m, 4H), 2.80 (s, 3H), 3.00 (m, 1H), 3.42 (m, 1H), 4.40 (m, 1H), 4.58 (dd, 1H), 7.00 (d, 1H), 7.22 (d, 1H), 7.40 (m, 2H), 7.59 (d, 2H). LRMS : m/z (TSP ⁺) 517.2, 519.2 [MH ⁺]
102 ^c		61 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.46 (m, 2H), 1.88-2.08 (m, 8H), 2.15 (m, 2H), 2.25 (m, 2H), 2.54 (s, 3H), 2.84 (m, 2H), 3.42 (m, 1H), 3.90 (d, 1H), 4.60 (d, 1H), 6.98 (d, 1H), 7.06 (m, 1H), 7.18 (m, 3H), 7.39 (d, 2H), 7.58 (m, 2H), 8.20 (d, 1H). LRMS : m/z (TSP ⁺) 539.2, 541.1 [MH ⁺]
103 ^{id}		33	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.15 (m, 4H), 2.50 (m, 3H), 2.77 (s, 3H), 2.80-3.00 (m, 4H), 3.08 (m, 2H), 3.17 (m, 2H), 3.63 (m, 2H), 4.20 (d, 1H), 4.34 (d, 1H), 7.42 (d, 1H), 7.59 (d, 1H), 7.68 (m, 2H), 7.80 (m, 1H), 7.92 (m, 1H), 8.22 (m, 1H), 8.42 (m, 1H), 8.72 (m, 1H), 8.91 (s, 1H).

104 ^{1b}		42	<p>¹Hnmr (CD₃OD, 400MHz) δ: 2.15 (m, 4H), 2.39 (m, 2H), 2.50 (m, 3H), 2.77 (s, 3H), 2.80-3.00 (m, 2H), 3.06 (m, 2H), 3.19 (m, 2H), 3.63 (m, 2H), 4.21 (d, 1H), 4.34 (d, 1H), 7.42 (d, 1H), 7.59 (d, 1H), 7.68 (m, 2H), 7.80 (m, 1H), 7.95 (m, 2H), 8.44 (m, 1H); 8.80 (m, 2H).</p> <p>LRMS : m/z (TSP⁺) 539.3 [MH⁺]</p>
105 ^f		23 pale yellow solid	<p>¹Hnmr (CDCl₃, 400MHz) δ: 1.60 (d, 1H), 1.75-2.10 (m, 7H), 2.18 (m, 2H), 2.32 (m, 2H), 2.44 (m, 2H), 2.56 (s, 3H), 2.60 (m, 1H), 2.75 (m, 2H), 3.94 (d, 1H), 4.60 (d, 1H), 7.00 (d, 1H), 7.20 (m, 2H), 7.36 (d, 1H), 7.40 (dd, 2H), 7.60 (m, 2H), 7.71 (dd, 1H), 8.50 (d, 1H).</p> <p>LRMS : m/z (ES⁺) 539, 541 [MH⁺]</p> <p>Microanalysis found: C, 63.12; H, 5.94; N, 9.94.</p> <p>C₂₉H₃₂Cl₂N₄O₂·0.2CH₂Cl₂ requires C, 63.02; H, 6.12; N, 9.92%.</p>
106 ^g		38 white foam	<p>¹Hnmr (CDCl₃, 300MHz) δ: 1.65 (m, 3H), 1.91-2.40 (m, 11H), 2.55 (s, 3H), 2.63 (m, 2H), 3.96 (d, 1H), 4.64 (d, 1H), 7.01 (d, 1H), 7.28 (m, 3H), 7.40 (m, 4H), 7.60 (m, 2H).</p> <p>LRMS : m/z (TSP⁺) 573.0, 575.1 [MH⁺]</p>

107 ^h		42 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.80-2.37 (m, 14H), 2.57 (m, 6H), 3.54 (m, 2H), 3.92 (d, 1H), 4.63 (d, 1H), 7.00 (m, 1H), 7.19-7.43 (m, 8H), 7.60 (m, 2H). LRMS : m/z (ES ⁺) 552, 554 [MH ⁺]
108 ⁱ		10 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.94 (m, 2H), 2.04-2.20 (m, 4H), 2.24 (m, 2H), 2.50 (m, 7H), 3.30 (m, 4H), 3.90 (d, 1H), 4.64 (d, 1H), 6.78 (d, 1H), 6.80 (dd, 1H), 6.97 (d, 1H), 7.17 (m, 1H), 7.21 (m, 1H), 7.40 (m, 2H), 7.58 (m, 2H), 8.12 (d, 1H). Microanalysis found: C, 60.14; H, 5.99; N, 12.28. C ₂₈ H ₃₁ Cl ₂ N ₅ O ₂ ·H ₂ O requires C, 60.22; H, 5.96; N, 12.54%.
109 ^j		72 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.93-2.63 (m, 15H), 3.39 (m, 4H), 3.86 (s, 3H), 3.92 (d, 1H), 4.74 (d, 1H), 6.83 (m, 1H), 7.03 (dd, 2H), 7.28 (m, 1H), 7.43 (d, 2H), 7.61 (m, 2H), 7.88 (d, 1H). LRMS : m/z (ES ⁺) 554, 556 [MH ⁺] Microanalysis found: C, 61.92; H, 5.95; N, 11.86. C ₂₈ H ₃₃ Cl ₂ N ₅ O ₂ ·0.5H ₂ O requires C, 61.81; H, 6.08; N, 12.43%.

110 ^k		76 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.97 (m, 2H), 2.07-2.62 (m, 13H), 3.23 (m, 4H), 3.92 (d, 1H), 4.78 (d, 1H), 5.80 (s, 1H), 7.03 (m, 1H), 7.09 (m, 1H), 7.28 (m, 1H), 7.44 (m, 2H), 7.62 (m, 2H), 8.30 (m, 1H), 8.40 (m, 2H). LRMS : m/z (TSP ⁺) 568.0, 570.1 [MH ⁺] Microanalysis found: C, 60.02; H, 5.85; N, 14.25. C ₂₈ H ₃₂ N ₆ Cl ₂ O ₂ ; 0.2CH ₂ Cl ₂ requires C, 60.00; H, 5.59; N, 14.38%.
111 ^l		71 white foam	¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.96 (m, 2H), 2.04-2.59 (m, 13H), 3.68 (m, 4H), 3.92 (d, 1H), 4.74 (d, 1H), 6.75 (m, 1H), 7.00 (d, 1H), 7.27 (m, 1H), 7.43 (m, 2H), 7.61 (m, 2H), 7.75 (d, 1H), 8.32 (s, 1H). LRMS : m/z (TSP ⁺) 550.0, 552.0 [MH ⁺] Microanalysis found: C, 62.70; H, 5.55; N, 14.79. C ₂₈ H ₃₀ N ₆ Cl ₂ O; 0.1CH ₂ Cl ₂ requires C, 62.64; H, 5.46; N, 15.06%.

1 = isolated as the hydrochloride salt

Starting amines:

- a = 4-(3-azetidiny)-1-piperazine sulphonamide trifluoroacetate as prepared in WO 9725322
- 5 b = N-methyl-N-(4-piperidinyl)acetamide hydrochloride from preparation 60
- c = 2-(4-piperidinyl)pyridine 1-oxide as prepared in WO 0037028
- d = 3-(4-piperidinyl)pyridine 1-oxide dihydrochloride as prepared in preparation 73
- e = 4-(4-piperidinyl)pyridine 1-oxide dihydrochloride as prepared in preparation 72
- f = 4-(2-pyridinyl)-4-piperidinol as prepared in DE 2630152
- 10 g = 4-(4-chlorophenyl)-4-piperidinol

h = (4-phenyl-4-piperidiny)methanol

i = 1-(1-oxido-2-pyridinyl)piperazine dihydrochloride as prepared in preparation 74


j = 1-(3-methoxy-2-pyridinyl)piperazine as prepared in EP 345808

k = 2-(1-piperazinyl)nicotinamide as prepared in J.Med.Chem. 1983; 26(12); 1698.

5 l = 2-(1-piperazinyl)nicotinonitrile

Example 112

(5S)-5-(3,4-Dichlorophenyl)-5-(2-[(1-oxido-2-pyridinyl)methyl]amino)ethyl)-1-(2-pyridinyl)-2-piperidinone



Triethylamine (287 μ l, 2.06mmol) was added to a suspension of the aldehyde from preparation 11a (250mg, 0.68mmol) in dichloromethane (10ml) followed by (1-oxido-2-pyridinyl)methylamine (J.O.C. 1974; 39(9); 1250) (136mg, 0.68mmol) and the mixture stirred at room temperature for 15 minutes. Acetic acid (158 μ l, 2.75mmol) and sodium triacetoxyborohydride (292mg, 1.38mmol) were added and the reaction stirred at room temperature for 18 hours. The mixture was washed with 2N sodium hydroxide solution (5ml), dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (95:5:0.5 to 93:7:1) to afford the title compound as a white foam, 224mg.


¹Hnmr (CDCl₃, 400MHz) δ : 1.90 (m, 1H), 2.00 (m, 1H), 2.15 (m, 1H), 2.28 (m, 3H), 2.47 (m, 1H), 2.58 (m, 1H), 3.85 (s, 2H), 3.90 (d, 1H), 4.58 (d, 1H), 7.15 (dd, 1H), 7.19 (m, 4H), 7.39 (d, 1H), 7.47 (s, 1H), 7.70 (q, 2H), 8.19 (d, 1H), 8.47 (d, 1H).

LRMS : m/z (TSP⁺) 471.1, 473.1 [MH⁺]

Microanalysis found: C, 58.74; H, 5.14; N, 11.28. C₂₄H₂₄Cl₂N₄O₂·0.3CH₂Cl₂ requires C, 58.74; H, 4.99; N, 11.28%.



Examples 113 to 118



The following examples of general structure:




were prepared from the aldehyde from preparation 11b and the appropriate
 5 amines, following a similar procedure to that described in example 112.

Example	R	Yield (%)	Data
113 ^a		43 clear oil	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.82-2.32 (m, 10H), 2.39 (m, 2H), 2.56 (m, 1H), 3.22 (s, 3H), 3.30 (m, 2H), 3.88 (d, 1H), 4.52 (d, 1H), 7.10 (m, 1H), 7.19 (d, 1H), 7.38 (d, 1H), 7.43 (s, 1H), 7.66 (s, 2H), 8.43 (d, 1H). LRMS : m/z (TSP ⁺) 436.1, 438.1 [MH ⁺] Microanalysis found: C, 60.17; H, 6.24; N, 9.48. C ₂₂ H ₂₇ Cl ₂ N ₃ O ₂ requires C, 60.55; H, 6.24; N, 9.63%.

114 ^b		42 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.62 (m, 2H), 1.72 (m, 1H), 1.84 (m, 1H), 2.00 (s, 3H), 2.10 (m, 6H), 2.30 (m, 4H), 2.55-2.75 (m, 4H), 2.91 (t, 1H), 3.41 (t, 2H), 3.88 (d, 1H), 4.54 (d, 1H), 5.45 (s, 1H), 7.17 (m, 1H), 7.21 (m, 2H), 7.30 (m, 2H), 7.37 (m, 2H), 7.40 (d, 1H), 7.45 (s, 1H), 7.70 (s, 2H), 8.50 (d, 1H). LRMS : m/z (TSP ⁺) 620.3, 622.3 [MH ⁺]
115 ^c		48 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.80 (d, 2H), 2.00 (m, 4H), 2.13 (q, 2H), 2.21 (m, 2H), 2.33 (dd, 2H), 2.46 (m, 2H), 2.60 (s, 3H), 2.93 (d, 1H), 3.01 (d, 1H), 4.00 (d, 1H), 4.09 (m, 1H), 4.75 (d, 1H), 7.18 (m, 3H), 7.28 (d, 1H), 7.45 (d, 1H), 7.46 (d, 1H), 7.53 (s, 1H), 7.68 (d, 1H), 7.75 (d, 2H), 8.50 (d, 1H) LRMS : m/z (TSP ⁺) 562.1, 564.1 [MH ⁺] Microanalysis found: C, 62.70; H, 5.64; N, 11.92. C ₃₁ H ₃₃ Cl ₂ N ₅ O; 0.18CH ₂ Cl ₂ requires C, 62.88; H, 5.69; N, 11.66%.

116 ^d		79 white foam	<p>¹Hnmr (CDCl₃, 400MHz) δ: 1.78 (d, 2H), 2.00 (m, 4H), 2.07 (m, 1H), 2.19 (q, 2H), 2.32 (m, 4H), 2.61 (m, 1H), 2.90 (m, 2H), 3.99 (d, 1H), 4.30 (m, 1H), 4.70 (d, 1H), 7.05 (s, 3H), 7.18 (m, 2H), 7.43 (d, 1H), 7.50 (s, 1H), 7.72 (s, 2H), 8.50 (d, 1H), 8.62 (s, 1H).</p> <p>LRMS : m/z (TSP*) 564.1, 566.1 [MH⁺]</p> <p>Microanalysis found: C, 59.45; H, 5.34; N, 11.39.</p> <p>C₃₀H₃₁Cl₂N₅O₂; 0.6CH₂Cl₂ requires C, 59.72; H, 5.27; N, 11.38%.</p>
117 ^e		16 white foam	<p>¹Hnmr (CDCl₃, 400MHz) δ: 1.73-1.85 (m, 3H), 1.85-1.97 (m, 3H), 1.97-2.06 (m, 3H), 2.06-2.20 (m, 2H), 2.28 (m, 2H), 2.55 (m, 1H), 2.80 (m, 3H), 3.90 (d, 1H), 4.41 (d, 1H), 7.09 (m, 1H), 7.18 (m, 3H), 7.20 (s, 1H), 7.38 (d, 1H), 7.46 (s, 1H), 7.65 (m, 3H), 8.43 (d, 1H), 9.33 (bs, 1H).</p> <p>LRMS : m/z (TSP*) 548.1, 550.1 [MH⁺]</p> <p>Microanalysis found: C, 63.76; H, 5.85; N, 11.99.</p> <p>C₃₀H₃₁Cl₂N₅O; 0.25CH₂Cl₂ requires C, 63.77; H, 5.57; N, 12.29%.</p>

118 ¹		66 white foam	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.85-2.05 (m, 11H), 2.30 (m, 2H), 2.59 (dd, 1H), 2.75-2.95 (m, 3H), 3.94 (d, 1H), 4.70 (d, 1H), 7.12 (dd, 1H), 7.23 (d, 1H), 7.29 (d, 2H), 7.41 (d, 1H), 7.49 (d, 1H), 7.50 (s, 1H), 7.70 (m, 3H), 8.50 (d, 1H). LRMS : m/z (TSP ⁺) 549.2, 551.1 [MH ⁺] Microanalysis found: C, 64.55; H, 5.49; N, 9.89. C ₃₀ H ₃₀ Cl ₂ N ₄ O ₂ ·0.06CH ₂ Cl ₂ requires C, 64.41; H, 5.43; N, 9.96%.
------------------	---	---------------------	--

Starting amines:

a = N-(2-methoxyethyl)methylamine

b = N-[1-(3-azetidyl)-4-phenyl-4-piperidinyl]acetamide dihydrochloride from preparation 34

5 c = 2-methyl-1-(4-piperidinyl)-1H-benzimidazole hydrochloride as prepared in J. Heterocycl. Chem. 1983; 20(3); 566

d = 1-(4-piperidinyl)-1,3-dihydro-2H-benzimidazol-2-one


e = 2-(4-piperidinyl)-1H-benzimidazole

f = 2-(4-piperidinyl)-1,3-benzoxazole

10

Example 119

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methoxy-2-pyridinyl)-5-[2-[3-(4-morpholinyl)-1-azetidyl]ethyl]-2-piperidinone



15

The title compound was obtained as a white foam in 80% yield from the aldehyde from preparation 18 and 3-morpholinoazetidine dihydrochloride (WO 9725322), following the procedure described in example 112.


¹Hnmr (CDCl₃, 400MHz) δ: 1.68 (m, 2H), 1.79 (m, 1H), 2.05 (m, 2H), 2.21 (m, 6H), 2.50 (m, 1H), 2.65 (m, 2H), 2.85 (m, 1H), 3.34 (q, 2H), 3.65 (m, 4H), 3.73 (d, 1H), 3.96 (s, 3H), 4.59 (d, 1H), 6.58 (d, 1H), 7.21 (d, 2H), 7.39 (d, 1H), 7.55 (s, 1H), 7.59 (dd, 1H).

LRMS : m/z (TSP⁺) 519.4, 521.4 [MH⁺]

Microanalysis found: C, 58.97; H, 6.18; N, 10.47. C₂₆H₃₂Cl₂N₄O₃·0.15CH₂Cl₂ requires C, 59.02; H, 6.12; N, 10.53%.

Example 120

N-(1-(2-((3S)-3-(3,4-Dichlorophenyl)-1-(6-methoxy-2-pyridinyl)-6-oxopiperidinyl)ethyl)-4-piperidinyl)-N-methylacetamide



The title compound was obtained as a white foam in 63% yield from the aldehyde from preparation 18 and N-methyl-N-(4-piperidinyl)acetamide hydrochloride from preparation 60 following the procedure described in example 112.


¹Hnmr (CDCl₃, 400MHz) δ: 1.50 (m, 4H), 1.80-2.20 (m, 7H), 2.05 (s, 3H), 2.28 (m, 2H), 2.55 (m, 1H), 2.78 (2xs, 3H), 2.80 (m, 2H), 3.41, 4.39 (2xm, 1H), 3.81 (m, 1H), 3.97 (s, 3H), 4.65 (d, 1H), 6.60 (d, 1H), 7.25 (m, 2H), 7.41 (m, 1H), 7.60 (m, 2H).

LRMS : m/z (TSP⁺) 533.4, 535.4 [MH⁺]

Microanalysis found: C, 59.50; H, 6.38; N, 10.16. C₂₇H₃₄Cl₂N₄O₃·0.18CH₂Cl₂ requires C, 59.49; H, 6.31; N, 10.21%.

Example 121

(5S)-5-(3,4-Dichlorophenyl)-5-{2-[(3R)-3-methoxypyrrolidinyl]ethyl}-1-(2-pyridinyl)-2-piperidinone




- 5 Triethylamine (0.1ml, 1.6mmol) was added to a solution of the aldehyde hydrochloride from preparation 11b (200mg, 0.5mmol) in dichloromethane (4ml), followed by (3R)-3-methoxypyrrolidine trifluoroacetate from preparation 47 (186mg, 0.7mmol), and the solution stirred at room temperature for 10 minutes. Sodium triacetoxyborohydride (159mg, 0.75mmol) and acetic acid
10 (60 μ l, 2mmol) were added, and the reaction stirred at room temperature for 18 hours. Methanol was added, the mixture stirred for 10 minutes, then concentrated under reduced pressure. The residue was partitioned between sodium carbonate solution and dichloromethane, the layers separated, and the organic phase evaporated under reduced pressure, to afford the title compound
15 as a white gum, 199mg.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.74 (m, 1H), 1.84-2.00 (m, 3H), 2.10 (m, 2H), 2.25 (m, 4H), 2.44 (m, 1H), 2.55 (m, 3H), 3.20 (s, 3H), 3.80 (m, 1H), 4.10 (d, 1H), 4.54 (d, 1H), 7.08 (m, 1H), 7.19 (d, 1H), 7.39 (d, 1H), 7.42 (s, 1H), 7.65 (d, 2H), 8.46 (d, 1H).

- 20 LRMS : m/z (TSP $^+$) 448.0, 449.9 [MH^+]

Example 122

(5S)-5-(3,4-Dichlorophenyl)-5-[2-[(3S)-3-methoxypyrrolidinyl]ethyl]-1-(2-pyridinyl)-2-piperidinone



5

The title compound was obtained as a clear gum, in 85% yield from the aldehyde hydrochloride from preparation 11b and (3S)-3-methoxypyrrolidine trifluoroacetate from preparation 48, following the procedure described in example 121.


¹Hnmr (CDCl₃, 400MHz) δ: 1.74 (m, 1H), 1.82-2.16 (m, 5H), 2.23 (m, 4H), 2.44 (m, 1H), 2.52 (m, 3H), 3.20 (s, 3H), 3.80 (m, 1H), 3.88 (d, 1H), 4.10 (d, 1H), 4.54 (d, 1H), 7.08 (m, 1H), 7.19 (d, 1H), 7.39 (d, 1H), 7.42 (s, 1H), 7.65 (d, 2H), 8.46 (d, 1H).

LRMS : m/z (TSP⁺) 448.1, 450.1 [MH⁺]

15

Example 123

tert-Butyl 4-[2-[(3S)-3-(3,4-dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl]-1-piperazinecarboxylate



20


The title compound was obtained as a white foam in 73% yield from the aldehyde hydrochloride from preparation 11b and tert-butyl 1-

piperazinecarboxylate, following a similar procedure to that described in example 121.

¹Hnmr (CDCl₃, 400MHz) δ: 1.40 (s, 9H), 1.90 (m, 2H), 1.98 (m, 1H), 2.04-2.34 (m, 8H), 2.56 (m, 1H), 3.30 (m, 4H), 3.92 (d, 1H), 4.62 (d, 1H), 7.10 (m, 1H),
 5 7.19 (d, 1H), 7.38 (d, 1H), 7.42 (s, 1H), 7.66 (d, 2H), 8.44 (d, 1H).
 LRMS : m/z (TSP⁺) 533.3 [MH⁺]

Example 124

10 (5S)-5-(3,4-Dichlorophenyl)-5-[2-[4-(2-methoxyethyl)-1-piperazinyl]ethyl]-1-(2-pyridinyl)-2-piperidinone




1-(2-Methoxyethyl)piperazine hydrochloride (90mg, 0.5mmol), followed by triethylamine (100mg, 1mmol) were added to a solution of the aldehyde
 15 hydrochloride from preparation 11b (200mg, 0.5mmol) in dichloromethane (2.5ml), and stirring continued for 20 minutes. Acetic acid (50mg, 0.83mmol) and sodium triacetoxyborohydride (150mg, 0.71mmol) were then added and the reaction stirred at room temperature for 1 hour. Methanol (2ml) was added, the mixture stirred for 20 minutes and then evaporated under reduced pressure.
 20 The residue was purified by column chromatography on silica gel using dichloromethane:methanol (93:7) as eluant. The product was redissolved in dichloromethane (10ml), washed with 10% aqueous potassium carbonate solution (3ml), then dried (Na₂SO₄) and evaporated under reduced pressure, to afford the title compound as a clear foam, 180mg.
 25 ¹Hnmr (CDCl₃, 400MHz) δ: 1.88 (m, 2H), 1.99 (m, 1H), 2.10 (m, 2H), 2.21-2.60 (m, 13H), 3.30 (s, 3H), 3.42 (t, 2H), 3.92 (d, 1H), 4.60 (d, 1H), 7.10 (m, 1H), 7.18 (d, 1H), 7.37 (d, 1H), 7.42 (s, 1H), 7.64 (m, 2H), 8.44 (m, 1H).
 LRMS : m/z (ES⁺) 491 [MH⁺]

Microanalysis found: C, 61.10; H, 6.56; N, 11.40. $C_{25}H_{32}Cl_2N_4O_2 \cdot 0.5H_2O$ requires C, 60.00; H, 6.65; N, 11.19%.

5 Example 125

N-((3*S*)-1-(2-((3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinylethyl)pyrrolidinyl)-*N*-methylacetamide




A mixture of the aldehyde from preparation 11a (200mg, 0.5mmol), the amine
10 from preparation 44 (256mg, 1.0mmol), triethylamine (0.21ml, 1.5mmol), acetic
acid (0.12ml, 1.55mmol) and sodium triacetoxyborohydride (211mg, 1.0mmol)
in dichloromethane (50ml) were stirred at room temperature in a STEM® block,
for 24 hours. The mixture was washed with aqueous sodium bicarbonate
solution and water, the organic layer filtered, dried ($MgSO_4$) and evaporated
5 under reduced pressure. The crude product was purified by column
chromatography on silica gel using dichloromethane:methanol:0.88 ammonia
(95:5:1) as eluant, to give the title compound.

1H nmr ($CDCl_3$, 400MHz) δ : 1.58 (m, 2H), 1.90-2.70 (m, 17H), 2.96 (m, 2H), 3.96
(d, 1H), 4.62 (m, 1H), 7.16 (d, 1H), 7.24 (m, 1H), 7.42 (d, 1H), 7.50 (s, 1H), 7.74
20 (s, 2H), 8.50 (d, 1H).

LRMS : m/z (TSP⁺) 489.1, 491.2 [MH^+]



Examples 126 to 130

The following compounds of general formula:





- 5 were prepared from the appropriate aldehydes and amines, according to the procedure described in example 125.

Example	R ¹	R ²	Data
126 ^a		H	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.53-1.80 (m, 2H), 1.90-2.60 (m, 15H), 2.78 (m, 1H), 2.92 (m, 2H), 3.92 (d, 1H), 4.36, 5.20 (2xs, 1H), 4.62 (m, 1H), 7.12 (m, 1H), 7.20 (d, 1H), 7.40 (d, 1H), 7.45 (s, 1H), 7.72 (m, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 489.1, 491.1 [MH ⁺] Microanalysis found: C, 60.32; H, 6.11; N, 11.25. C ₂₅ H ₃₀ Cl ₂ N ₄ O ₂ ·0.5H ₂ O requires C, 60.24; H, 6.27; N, 11.24%.

127 ^{1b}		H	<p>¹Hnmr (CDCl₃, 400MHz) δ: 1.90-2.40 (m, 15H), 2.58 (m, 2H), 2.80 (m, 1H), 3.36 (m, 2H), 3.94 (d, 1H), 4.64 (m, 2H), 7.16 (m, 1H), 7.20 (d, 1H), 7.42 (d, 1H), 7.48 (s, 1H), 7.70 (m, 2H), 8.48 (d, 1H).</p> <p>LRMS : m/z (TSP⁺) 501.1, 503.2 [MH⁺]</p> <p>Microanalysis found: C, 60.68; H, 6.16; N, 10.68.</p> <p>C₂₆H₃₀Cl₂N₄O₂:0.2CH₂Cl₂ requires C, 60.70; H, 5.91; N, 10.81%.</p>
128 ^{1c}		H	<p>¹Hnmr (CDCl₃, 400MHz) δ: 1.70-2.70 (m, 18H), 3.38 (m, 2H), 3.92 (d, 1H), 4.62 (d, 1H), 4.70 (bs, 1H), 7.16 (dd, 1H), 7.20 (d, 1H), 7.42 (d, 1H), 7.48 (s, 1H), 7.70 (m, 2H), 8.48 (d, 1H).</p> <p>LRMS : m/z (TSP⁺) 501.1, 503.1 [MH⁺]</p> <p>Microanalysis found: C, 60.74; H, 6.18; N, 10.79.</p> <p>C₂₆H₃₀Cl₂N₄O₂:0.2CH₂Cl₂ requires C, 60.70; H, 5.91; N, 10.81%.</p>

101

129 ^d		CH ₃	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.70-2.80 (m, 22H), 2.90 (s, 2H), 3.90 (m, 1H), 4.58 (dd, 1H), 7.00 (d, 1H), 7.20 (d, 1H), 7.40 (m, 2H), 7.58 (m, 2H). LRMS : m/z (TSP ⁺) 503.1 [MH ⁺]
130 ^e		CH ₃	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.90-2.80 (m, 22H), 2.90 (m, 2H), 4.88 (d, 1H), 4.34, 5.20 (2xbs, 1H), 7.00 (d, 1H), 7.22 (d, 1H), 7.40 (m, 2H), 7.60 (m, 2H). LRMS : m/z (TSP ⁺) 503.2 [MH ⁺] Microanalysis found: C, 60.28; H, 6.69; N, 10.99. C ₂₈ H ₃₂ Cl ₂ N ₄ O ₂ ·0.2CH ₂ Cl ₂ requires C, 60.46; H, 6.27; N, 10.76%.

1 = 1.5mmol amine/1mmol NaBH(OAc)/1mmol Et₃N/1.05mmol AcOH

Starting amines:

a = N-methyl-N-[(3S)-pyrrolidinyl]acetamide trifluoroacetamide from preparation 43

5 b = 1-[(3S)-pyrrolidin-3-yl]-2-pyrrolidine from preparation 53


c = 1-[(3R)-pyrrolidin-3-yl]-2-pyrrolidine from preparation 54

d = N-methyl-N-[(3R)-pyrrolidinyl]acetamide trifluoroacetate from preparation 43

e = N-methyl-N-[(3S)-pyrrolidinyl]acetamide trifluoroacetate from preparation 44

Example 131

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-(2-[3-(4-morpholinyl)-1-azetidiny]ethyl)-2-piperidinone




A mixture of the aldehyde from preparation 12a (2.0g, 5.3mmol), 3-morpholinoazetidine dihydrochloride (WO 9725322) (1.25g, 5.83mmol), triethylamine (1.84ml, 13.3mmol) and titanium isopropoxide (16ml, 53mmol) in ethanol (20ml), was stirred at room temperature for 18 hours. Sodium borohydride (320mg, 7.95mmol) in ethanol (19ml) was then added and the reaction stirred for 30 minutes. Sodium hydroxide was added, the resulting precipitate filtered off, and washed with ethyl acetate. The filtrate was washed with water (2x) and brine (2x), dried (MgSO₄) and evaporated under reduced pressure. The residual gum was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol (100:0 to 99:1) to afford the title compound as a white solid, 1.58g.

¹Hnmr (CDCl₃, 400MHz) δ: 1.72 (m, 1H), 1.82 (m, 1H), 2.14 (m, 2H), 2.24 (m, 7H), 2.56 (m, 1H), 2.58 (s, 3H), 2.75 (m, 2H), 2.90 (m, 1H), 3.40 (m, 2H), 3.65 (m, 4H), 3.88 (d, 1H), 4.50 (d, 1H), 7.00 (d, 1H), 7.20 (d, 1H), 7.40 (d, 2H), 7.55 (s, 1H), 7.60 (dd, 1H).

LRMS : m/z (TSP⁺) 503.6, 505.2 [MH⁺]


Examples 132 to 134

The compounds of the general structure:



were prepared from the aldehyde from preparation 12a and the appropriate
 5 amines, following a similar procedure to that described in example 131.

Example	R	Yield (%)	Data
132 ^a		21 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.10-1.90 (m, 9H), 2.10 (m, 3H), 2.25 (m, 3H), 2.56 (s, 3H), 2.56 (m, 3H), 2.74 (s, 3H), 3.28 (m, 2H), 3.75 (d, 2H), 3.84 (d, 1H), 4.50 (d, 1H), 6.98 (d, 1H), 7.20 (d, 1H), 7.28 (d, 1H), 7.40 (dd, 1H), 7.55 (m, 2H). LRMS : m/z (TSP ⁺) 580.1, 582.1 [MH ⁺]
133 ^b			¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.80 (m, 1H), 2.10-2.40 (m, 11H), 2.55 (s, 4H), 2.75 (m, 5H), 2.94 (m, 1H), 3.20 (m, 4H), 3.40 (d, 1H), 3.88 (d, 1H), 4.64 (d, 1H), 7.00 (d, 1H), 7.20 (d, 1H), 7.40 (d, 2H), 7.54 (s, 1H), 7.60 (dd, 1H). LRMS : m/z (ES ⁺) 580, 582 [MH ⁺]

134 ^{1c}		9 white solid	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.50-2.60 (m, 20H), 3.95 (d, 1H), 4.62 (d, 1H), 7.00 (d, 1H), 7.25 (d, 2H), 7.35 (d, 2H), 7.42 (m, 4H), 7.60 (dd, 2H). LRMS : m/z (TSP ⁺) 538.4 [MH ⁺] Microanalysis found: C, 64.82; H, 6.11; N, 7.49. C ₃₀ H ₃₃ Cl ₂ N ₃ O ₂ ; 0.1(CH ₂ CH ₂) ₂ O; 0.9H ₂ O requires C, 66.91; H, 6.18; N, 7.80%.
-------------------	---	---------------------	--

1 = purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:0.05) as eluant and triturated with diethyl ether.

5 Starting amines:


a = 4-(3-azetidiny)-1-(methylsulfonyl)piperidine hydrochloride as prepared in EP 992493

b = 1-(3-azetidiny)-4-(methylsulphonyl)piperazine trifluoroacetate as prepared in WO 9725322

c = 4-hydroxy-4-phenylpiperidine

Example 135a

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-[2-[3-(4-hydroxypiperidinyl)azetidiny]ethyl]-2-piperidinone



5

3-(4-hydroxypiperidinyl)azetidine trifluoroacetate (WO 96/05193 – preparation 77)(25.2gm, 65mmol) was dissolved in tetrahydrofuran (100ml) and added to a solution of the aldehyde from preparation 12a (24.8gm, 65mmol) in dichloromethane (300ml). Sodium triacetoxyborohydride (21gm, 98mmol) was added and the reaction stirred at room temperature for 18 hours. The reaction was concentrated under reduced pressure and the resulting orange oil was dissolved in ethyl acetate (500ml), washed with 2N sodium hydroxide solution (200ml), the aqueous layer extracted with ethyl acetate (2x 200ml), the combined organic extracts were dried (MgSO₄), and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant to afford the title compound as a white foam (17gm).


¹Hnmr (CDCl₃, 400MHz) δ: 1.52 (m, 3H), 1.70 (m, 1H), 1.85 (m, 3H), 1.95 (m, 2H), 2.15 (m, 2H), 2.25 (m, 3H), 2.55 (m, 6H), 2.66 (m, 2H), 2.86 (t, 1H), 3.42 (m, 2H), 3.70 (m, 1H), 3.84 (d, 1H), 4.50 (d, 1H), 7.00 (d, 1H), 7.20 (d, 1H), 7.40 (dd, 2H), 7.58 (s, 1H), 7.60 (dd, 1H).

LRMS : m/z (TSP*) 517.3 [MH⁺]

20

Example 135b

(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-[2-[3-(4-hydroxypiperidinyl)azetidinyl]ethyl]-2-piperidinone bisfumarate



5

Fumaric acid (8gm, 69mmol) was dissolved in a mixture of water (3.4ml, 190mmol) and tetrahydrofuran (100ml) before adding to a solution of the title compound from example 135a (24.7gm, 47.7mmol) in tetrahydrofuran (50ml).

- 10 Resulting solution was allowed to stand at room temperature for 1 hour, during which time a white precipitate had formed. Mixture filtered, filter cake washed with tetrahydrofuran (30ml), diethyl ether (2x 200ml) and dried at 50 °C for 48 hours to give a white solid (26.4gm). Solid recrystallised from refluxing 2% (vol/vol) aqueous tetrahydrofuran (250ml), upon cooling mixture filtered, filter cake washed with tetrahydrofuran (2x 100ml), diethyl ether (3x 500ml) and air dried for 20 minutes. Resulting solid was slurried in acetone (100ml) for 18 hours, filtered washed with diethyl ether and dried for 48 hours at 50 °C under reduced pressure (6mbar) to give the title compound as a fine white solid.

- 20 ¹Hnmr (CD₃OD, 400MHz) δ: 1.54 (m, 2H), 1.82 (m, 2H), 1.94 (m, 1H), 2.05 (m, 1H), 2.14 (m, 2H), 2.21 (m, 2H), 2.38 (m, 1H), 2.54 (m, 4H), 2.66 (m, 2H), 2.73 (m, 1H), 2.94 (m, 1H), 3.25 (m, 1H), 3.65 (m, 3H), 3.90 (d, 1H), 4.00 (m, 2H), 4.48 (d, 1H), 6.69 (s, 3H), 7.13 (d, 1H), 7.25 (d, 1H), 7.44 (d, 1H), 7.54 (d, 1H), 7.68 (t, 1H), 7.81 (s, 1H).

LRMS : m/z (TSP⁺) 517.3 [MH⁺]


- 25 Microanalysis found: C, 55.73; H, 5.67; N, 7.48.

C₂₇H₃₄Cl₂N₄O₂·2·C₄H₄O₄·0.25H₂O requires C, 55.74; H, 5.68; N, 7.43%.

Mpt. 175.5-177°C

Example 136

(5S)-5-(3,4-Dichlorophenyl)-1-(5-methyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone




- 5 The title compound was prepared as a yellow solid in 6% yield from the aldehyde from preparation 14 and 3-morpholinoazetidine dihydrochloride (WO 9725322), following a similar procedure to that described in example 131, except ethyl acetate:pentane (10:90 to 100:0) was used as the column eluant.
- ¹Hnmr (CDCl₃, 400MHz) δ: 1.75 (m, 1H), 1.85 (m, 1H), 2.20 (m, 9H), 2.38 (s, 3H), 2.58 (m, 1H), 2.76 (m, 2H), 2.95 (m, 1H), 3.42 (m, 2H), 3.68 (m, 4H), 3.90 (d, 1H), 4.50 (d, 1H), 7.12 (d, 1H), 7.42 (d, 1H), 7.50 (s, 1H), 7.58 (m, 2H), 8.32 (s, 1H).
- 10 LRMS : m/z (TSP⁺) 503.3, 504.9 [MH⁺]

15

Example 137

(5S)-5-(3,4-Dichlorophenyl)-1-(6-ethyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone




The title compound was prepared as a white solid in 37% yield, from the aldehyde from preparation 16, and 3-morpholinoazetidine dihydrochloride (WO 9725322), following the procedure described in example 131.

5

Example 138

N-(1-[2-[(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(3-pyridinyl)piperidinyl]ethyl]-4-piperidinyl)-*N*-methylacetamide



10 Potassium carbonate (390mg, 2.82mmol), copper (90mg, 1.41mmol) and 3-bromopyridine (680μl, 7.06mmol) were added to the amine from preparation 83 (300mg, 0.70mmol) and the mixture stirred at 140°C for 20 hours. The mixture was partitioned between water (100ml) and ethyl acetate (100ml), and the layers separated. The aqueous phase was extracted with ethyl acetate, the

15 combined organic solutions washed with water, then brine, and dried (Na₂SO₄) and evaporated under reduced pressure. The crude oil was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol (95:5 to 85:15) to afford the title compound as a yellow oil, 80mg.


20 ¹Hnmr (CDCl₃, 400MHz) δ: 1.46 (m, 3H), 1.63 (m, 1H), 1.88 (m, 3H), 2.00 (s, 3H), 2.20 (m, 1H), 2.37 (m, 2H), 2.58 (m, 1H), 2.75 (m, 5H), 3.40 (s, 3H), 3.88 (d, 1H), 4.03 (m, 1H), 4.37 (m, 1H), 7.10 (d, 1H), 7.30 (m, 2H), 7.40 (d, 1H), 7.60 (d, 1H), 8.45 (s, 1H), 8.53 (s, 1H).

LRMS : m/z (ES⁺) 525, 527 [MH⁺]

25 Microanalysis found: C, 58.52; H, 6.18; N, 10.31. C₂₈H₃₂Cl₂N₄O₂·0.5CH₂Cl₂ requires C, 58.30; H, 6.09; N, 10.26%.

Example 139

N-(1-(2-((3*S*)-3-(3,4-Dichlorophenyl)-1-(6-methyl-3-pyridinyl)-6-oxopiperidinylethyl)-4-piperidiny)-*N*-methylacetamide



The title compound was obtained as a black solid in 71% yield, from the amine from preparation 83 and 5-bromo-2-methylpyridine, following a similar procedure to that described in example 138, except 1-methyl-2-pyrrolidinone was used as the reaction solvent.


¹Hnmr (CD₃OD, 400MHz) δ: 1.45 (m, 1H), 1.58 (m, 2H), 1.65 (m, 1H), 1.85 –
 10 2.14 (m, 10H), 2.24 (m, 2H), 2.42 (m, 1H), 2.56 (s, 3H), 2.74 (s, 1H), 2.82 (m,
 4H), 3.58 (m, 0.2H), 4.0 (d, 1H), 4.10 (d, 1H), 4.24 (m, 0.8H), 7.38 (d, 2H), 7.57
 (d, 1H), 7.60 (s, 1H), 7.63 (dd, 1H), 8.35 (s, 1H).

LRMS : m/z (ES⁺) 539, 541 [MNa⁺]

Microanalysis found: C, 61.39; H, 6.59; N, 10.41. C₂₇H₃₄Cl₂N₄O₂·0.6H₂O
 15 requires C, 61.38; H, 6.72; N, 10.60%.

Example 140

N-(1-(2-[(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyrazinyl)piperidinyl]ethyl)-4-piperidinyl)-*N*-methylacetamide



A mixture of the amine from preparation 83 (300mg, 0.70mmol), potassium *tert*-butoxide (160mg, 1.43mmol) and chloropyrazine (260 μ l, 2.8mmol) in *N*-methylpyrrolidine (5ml) was stirred at 100°C for 72 hours. The cooled mixture was partitioned between water (100ml) and ethyl acetate (100ml), and the layers separated. The aqueous phase was extracted with ethyl acetate, the
10 combined organic solutions washed with water and brine, then dried (MgSO₄) and evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (99:1:0.1 to 90:10:1) to afford the title compound as a brown, glass-like solid, 45mg.


15 ¹Hnmr (CDCl₃, 400MHz) δ : 1.55 (m, 4H), 1.83-2.20 (m, 11H), 2.26-2.43 (m, 2H), 2.60-2.82 (m, 5H), 3.92 (d, 1H), 4.38 (m, 1H), 4.64 (d, 1H), 7.18 (d, 1H), 7.39 (d, 1H), 7.43 (s, 1H), 8.35 (s, 1H), 8.39 (s, 1H), 9.19 (s, 1H).

LRMS : m/z (ES⁺) 526, 528 [MH⁺]

Microanalysis found: C, 52.88; H, 5.29; N, 12.27. C₂₅H₃₁Cl₂N₅O₂·CH₂Cl₂
20 requires C, 52.98; H, 5.64; N, 11.88%.

Example 141

N-(1-(2-[(3*S*)-3-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyrazinyl)-6-oxopiperidinyl]ethyl)-4-piperidinyl)-*N*-methylacetamide




- i The title compound was obtained as a brown solid in 4% yield, from the amine from preparation 83 and 2-chloro-6-methylpyrazine (Tetrahedron 1972; 28; 4155), following a similar procedure to that described in example 140.

¹Hnmr (CDCl₃, 400MHz) δ: 1.56 (m, 3H), 1.81-2.19 (m, 11H), 2.32 (m, 2H), 2.58 (s, 3H), 2.62 (m, 1H), 2.80 (m, 5H), 3.40 (m, 0.2H), 3.85 (dd, 1H), 4.42 (m, 0.8H), 4.58 (dd, 1H), 7.18 (d, 1H), 7.40 (d, 1H), 7.50 (s, 1H), 8.22 (s, 1H), 8.86 (s, 1H).

LRMS : m/z (ES⁺) 540, 542 [MNa⁺]

15 **Example 142**

(5S)-5-(3,4-Dichlorophenyl)-5-(2-{methyl[(1*R*)-1-phenylethyl]amino}ethyl)-1-(2-pyridinyl)-2-piperidinone dihydrochloride




- Formaldehyde (75μl, 37%aq, 0.92mmol) was added to a solution of the amine from example 37 (160mg, 0.30mmol) in dichloromethane (25ml), and the solution stirred for 5 minutes. Sodium triacetoxyborohydride (63mg, 0.30mmol)

- was added and the reaction stirred at room temperature for 18 hours. Tlc analysis showed starting material remaining, so additional formaldehyde (0.4ml, 37%aq, 4.93mmol) and sodium triacetoxyborohydride (62mg, 0.29mmol) were added, and the reaction stirred for an hour. The reaction was washed with
- 5 saturated aqueous sodium bicarbonate solution (25ml), brine (10ml), dried (MgSO_4) and concentrated under reduced pressure. The residual foam was redissolved in dichloromethane (10ml), and treated with 1N ethereal hydrochloric acid (5ml). This solution was then evaporated under reduced pressure, to afford the title compound as a white foam, 158mg.
- 10 $^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 1.63 (t, 3H), 2.06-3.00 (m, 11H), 4.04-4.65 (m, 3H), 7.07-7.73 (m, 9H), 7.90 (m, 1H), 8.40 (m, 1H), 8.59 (m, 1H).
LRMS : m/z (TSP $^+$) 482.1, 484.1 [MH^+]

15 Example 143


(5S)-5-(3,4-Dichlorophenyl)-5-(2-{methyl[(1S)-1-phenylethyl]amino}ethyl)-1-(2-pyridinyl)-2-piperidinone dihydrochloride



- 20 The title compound was prepared in 91% yield from the amine from example 38, following the procedure described in example 142.
- $^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 1.63 (t, 3H), 2.08-2.97 (m, 11H), 4.10-4.23 (m, 1H), 4.28 (m, 0.5H), 4.41 (m, 1H), 4.60 (m, 0.5H), 7.18-7.63 (m, 8H), 7.72 (dd, 1H), 7.92 (dd, 1H), 8.44 (dd, 1H), 8.60 (m, 1H).
- 25 LRMS : m/z (TSP $^+$) 482.1, 484.1 [MH^+]

Example 144


(5S)-5-(3,4-Dichlorophenyl)-5-(2-{methyl[(1R)-1-phenylethyl]amino}ethyl)-1-(6-methyl-2-pyridinyl)-2-piperidinone dihydrochloride



- 5 The title compound was prepared from the amine from example 41 as a white solid, following a similar procedure to that described in example 143.
- $^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 1.63 (t, 3H), 2.07-2.97 (m, 14H), 4.08 (d, 1H), 4.22 (q, 1H), 4.37 (m, 1H), 7.16-7.75 (m, 10H), 8.25-8.40 (m, 1H).
- LRMS : m/z (TSP^+) 496.1, 498.2 [MH^+]
- 10 Microanalysis found: C, 53.27; H, 6.15; N, 6.51. $\text{C}_{28}\text{H}_{31}\text{Cl}_2\text{N}_3\text{O}; 2\text{HCl}; 3.5\text{H}_2\text{O}$ requires C, 53.18; H, 5.90; N, 6.64%

Example 145

- 15 (5S)-5-(3,4-Dichlorophenyl)-5-(2-{methyl[3-(4-morpholinyl)propyl]amino}ethyl)-1-(2-pyridinyl)-2-piperidinone trihydrochloride



- Triethylamine (0.5ml) was added to a suspension of the amine from example 36
- 20 (267mg, 0.44mmol) in dichloromethane (20ml), followed by acetic acid (0.5ml), formaldehyde (0.36ml, 4.44mmol) and finally sodium triacetoxyborohydride


(94.6mg, 0.45mmol), and the reaction stirred at room temperature for 2 hours. The mixture was washed with saturated aqueous sodium bicarbonate solution (50ml), brine (25ml), dried (MgSO_4) and concentrated under reduced pressure. The crude product was redissolved in dichloromethane, 1N ethereal

5 hydrochloric acid added and the solution then evaporated under reduced pressure to afford the title compound as a white foam, 258mg.

$^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 2.22 (m, 2H), 2.32-2.59 (m, 6H), 2.78-2.90 (m, 5H), 3.10-3.20 (m, 5H), 3.23 (m, 1H), 3.52 (m, 2H), 3.86 (m, 2H), 4.14 (d, 2H), 4.30 (m, 1H), 4.48 (m, 1H), 7.46 (d, 1H), 7.60 (d, 1H), 7.65 (s, 1H), 7.80 (m, 1H),
10 8.15 (m, 1H), 8.04-8.56 (m, 2H).

Example 146

15 (5S)-5-(3,4-Dichlorophenyl)-5-[2-[methyl(2-pyridinylmethyl)amino]ethyl]-1-(2-pyridinyl)-2-piperidinone trihydrochloride



The title compound was prepared as a white foam in 90% yield from the amine from example 56, following the procedure described in example 145.

$^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 2.30-2.60 (m, 5H), 2.81 (m, 1H), 2.90 (s, 3H), 2.98 (m, 1H), 3.17 (m, 1H), 4.30 (d, 1H), 4.41 (d, 1H), 4.58 (s, 2H), 7.40 (d, 1H), 7.57 (d, 1H), 7.64 (m, 2H), 7.72 (d, 1H), 7.82 (dd, 1H), 8.08 (m, 2H), 8.62 (m, 3H).


LRMS : m/z (TSP^+) 471.1, 473.1 [MH^+]

Microanalysis found: C, 43.15; H, 5.29; N, 7.64%.

$\text{C}_{25}\text{H}_{26}\text{Cl}_2\text{N}_4\text{O} \cdot 3\text{HCl} \cdot 3.5\text{H}_2\text{O} \cdot \text{CH}_2\text{Cl}_2$ requires C, 42.96; H, 5.27; N, 7.71%.

Example 147


(5S)-5-(2-[3-(4-Amino-1-piperidinyl)-1-azetidiny]ethyl)-5-(3,4-dichlorophenyl)-1-(2-pyridinyl)-2-piperidinone



- 5 Trifluoroacetic acid (10ml) was added to an ice-cooled solution of the protected amine from preparation 84 (760mg, 1.26mmol) in dichloromethane (5ml), and the solution stirred at 0°C for 2 hours. The solution was poured into ice-cooled water (200ml), basified using 2N sodium hydroxide solution, and extracted with dichloromethane (3x200ml). The combined organic solutions were washed with
- 10 brine, dried (MgSO₄), and evaporated under reduced pressure to give the title compound as a white foam, 540mg.
- ¹Hnmr (CDCl₃, 300MHz) δ: 1.34 (m, 2H), 1.81 (m, 8H), 2.13 (m, 2H), 2.30 (m, 3H), 2.52-2.75 (m, 6H), 2.87 (m, 1H), 3.42 (m, 2H), 3.90 (d, 1H), 4.56 (d, 1H), 7.20 (m, 2H), 7.43 (m, 2H), 7.72 (m, 2H), 8.52 (d, 1H).
- 15 LRMS : m/z (ES⁺) 502, 504 [MH⁺]

Example 148

(5S)-5-(3,4-Dichlorophenyl)-5-[2-(1-piperazinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone



A solution of the protected amine from example 123 (923mg, 1.78mmol) in 4M HCl in dioxan (50ml) was stirred at room temperature for 18 hours. The reaction mixture was concentrated under reduced pressure and the residue partitioned between sodium bicarbonate solution and dichloromethane. The layers were separated, the organic phase dried (MgSO₄) and evaporated under reduced pressure to give a yellow solid, 831mg.


A sample (50mg) was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant to afford the title compound as a clear oil.

¹Hnmr (CDCl₃, 400MHz) δ : 1.83-2.16 (m, 6H), 2.23 (m, 5H), 2.58 (m, 1H), 2.89 (m, 4H), 3.89 (d, 1H), 4.60 (d, 1H), 7.12 (s, 1H), 7.19 (d, 1H), 7.38 (d, 1H), 7.44 (s, 1H), 7.68 (s, 2H), 8.45 (d, 1H).
LRMS : m/z (TSP⁺) 433.1, 43.1 [MH⁺]

15

Example 149

(5S)-5-[2-(4-Amino-1-piperidiny)ethyl]-5-(3,4-dichlorophenyl)-1-(2-pyridinyl)-2-piperidinone



A solution of the protected amine from example 54 (577mg, 1.05mmol) in 4M HCl in dioxan (10ml) was stirred at room temperature for 2 hours. The mixture was evaporated under reduced pressure and the residue partitioned between 10% aqueous sodium carbonate solution and diethyl ether, and the layers separated. The aqueous phase was extracted with diethyl ether, then dichloromethane, the combined organic solutions dried (MgSO₄) and evaporated under reduced pressure, to give the title compound as a white foam, 465mg.


117

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.42 (m, 2H), 1.83-1.98 (m, 7H), 2.13 (m, 2H), 2.29 (m, 2H), 2.59 (m, 1H), 2.73 (m, 3H), 3.05 (m, 2H), 3.91 (d, 1H), 4.56 (d, 1H), 7.13 (dd, 1H), 7.21 (d, 1H), 7.40 (d, 1H), 7.46 (s, 1H), 7.68 (m, 2H), 8.48 (d, 1H).

5 LRMS : m/z (TSP $^+$) 447.1, 449.1 [MH^+]

Example 150

10 (5S)-5-(3,4-Dichlorophenyl)-5-[2-[4-(methylamino)-1-piperidinylethyl]-1-(2-pyridinyl)-2-piperidinone dihydrochloride




Hydrogen chloride was bubbled through a solution of the protected amine from example 99 (340mg, 0.607mmol) in dichloromethane (50ml) for 5 minutes. The mixture was evaporated under reduced pressure to afford a quantitative amount of the title compound as a white foam.

15 $^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 1.98 (m, 2H), 1.98-2.58 (m, 8H), 2.70 (s, 3H), 2.81 (m, 2H), 3.00 (m, 2H), 3.38 (m, 2H), 3.62 (m, 2H), 4.22 (d, 1H), 4.39 (d, 1H), 7.41 (m, 1H), 7.58 (d, 1H), 7.66 (s, 1H), 7.78 (m, 1H), 8.02 (d, 1H), 8.58 (m, 1H).

20 LRMS : m/z (TSP $^+$) 461.1, 463.1 [MH^+]

Example 151

N-[1-(1-{2-[1-(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-3-azetidiny)-4-piperidinyl]acetamide



- 5 Triethylamine (85mg, 0.84mmol) and acetic anhydride (74mg, 0.63mmol) were added to a solution of the amine from example 147 (210mg, 0.42mmol) in dichloromethane (100ml), and the reaction stirred at room temperature for 20 minutes. The solution was washed with 2N sodium hydroxide solution, and the aqueous layer extracted with dichloromethane (2x50ml). The combined organic solutions were dried (MgSO₄), evaporated under reduced pressure and the residue azeotroped with dichloromethane (4x200ml), to afford the title compound as a white foam, 206mg.


- ¹Hnmr (CDCl₃, 300MHz) δ: 1.72-1.99 (m, 10H), 2.08-2.23 (m, 3H), 2.23-2.42 (m, 3H), 2.54-2.69 (m, 3H), 2.78 (m, 2H), 2.93 (m, 1H), 3.49 (m, 2H), 3.78 (m, 1H), 3.92 (d, 1H), 4.56 (d, 1H), 5.36 (m, 1H), 7.17 (m, 1H), 7.21 (d, 1H), 7.44 (m, 2H), 7.74 (m, 2H), 8.51 (m, 1H).

LRMS : m/z (ES⁺) 544, 546 [MH⁺]

Microanalysis found: C, 57.75; H, 7.00; N, 11.80. C₂₈H₃₅Cl₂N₅O₂·2H₂O requires C, 57.93; H, 6.77; N, 12.06%.

Example 152


(5S)-5-(3,4-Dichlorophenyl)-5-[2-(4-propionyl-1-piperazinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone






- A mixture of the piperazine from example 148 (168mg, 0.39mmol), propionyl chloride (34 μ l, 0.39mmol) and triethylamine (54 μ l, 0.39mmol) in dichloromethane (10ml) was stirred at room temperature for an hour. The mixture was washed with water, and the aqueous solution extracted with dichloromethane. The combined organic solutions were dried (MgSO₄) and
- 10 evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as eluant to afford the title compound as a clear gum, 52mg.
- ¹Hnmr (CDCl₃, 400MHz) δ : 1.09 (t, 3H), 1.88 (d, 2H), 1.99 (s, 1H), 2.13 (m, 2H), 2.23 (m, 8H), 2.58 (m, 1H), 3.33 (s, 2H), 3.50 (s, 2H), 3.90 (d, 1H), 4.62 (d, 1H),
- 15 7.11 (s, 1H), 7.19 (d, 1H), 7.38 (d, 1H), 7.43 (s, 1H), 7.68 (s, 2H), 8.42 (d, 1H).
LRMS : m/z (TSP⁺) 489.2, 491.2 [MH⁺]

Examples 153 to 156


- 20 The following compounds of general formula:



were prepared from the piperazine from example 148 and the appropriate acid chloride, according to the method described in example 152.

Example	R	Yield (%)	Data
153		42	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.97 (m, 2H), 2.03 (m, 1H), 2.18 (m, 2H), 2.30 (m, 6H), 2.60 (m, 1H), 3.42 (s, 3H), 3.43 (s, 2H), 3.57 (s, 2H), 3.98 (d, 1H), 4.08 (s, 2H), 4.68 (d, 1H), 7.18 (m, 1H), 7.23 (d, 1H), 7.42 (d, 1H), 7.50 (s, 1H), 7.75 (s, 2H), 8.51 (d, 1H). LRMS : m/z (TSP ⁺) 505.1, 507.2 [MH ⁺]
154		33	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.08 (d, 6H), 1.90-2.35 (m, 11H), 2.55 (s, 1H), 3.33 (m, 4H), 3.86 (d, 1H), 4.60 (d, 1H), 4.82 (m, 1H), 7.08 (m, 1H), 7.20 (d, 1H), 7.38 (d, 1H), 7.41 (s, 1H), 7.63 (s, 2H), 8.41 (s, 1H). LRMS : m/z (TSP ⁺) 519.2, 521.2 [MH ⁺]
155			¹ Hnmr (CDCl ₃ , 300MHz) δ: 1.86-2.10 (m, 3H), 2.10-2.40 (m, 9H), 2.60 (m, 1H), 2.81 (m, 6H), 3.19 (m, 3H), 3.97 (d, 1H), 4.64 (d, 1H), 7.12-7.32 (m, 2H), 7.40 (d, 1H), 7.49 (s, 1H), 7.74 (m, 2H), 8.48 (d, 1H). LRMS : m/z (TSP ⁺) 504.2, 506.2 [MH ⁺]


121

156		24	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.11 (t, 6H), 1.90-2.40 (m, 11H), 2.60 (m, 1H), 3.19 (m, 8H), 3.97 (d, 1H), 4.68 (d, 1H), 7.18 (m, 1H), 7.22 (m, 1H), 7.41 (d, 1H), 7.50 (s, 1H), 7.73 (d, 2H), 8.51 (d, 1H). LRMS : m/z (TSP ⁺) 532.3, 534.2 [MH ⁺]
-----	---	----	---

Example 157

N-(1-(2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl)-4-piperidinyl)-N,N-diethylurea

5




Triethylamine (40μl, 0.32mmol) and diethylcarbonyl chloride (40μl, 0.30mmol) were added to a solution of the amine from example 149 (120mg, 0.27mmol) in tetrahydrofuran (5ml), and the reaction stirred at 40°C for 18 hours. The mixture was concentrated under reduced pressure and the residue purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) to afford the title compound, 129mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.11 (t, 6H), 1.27 (m, 2H), 1.96 (m, 6H), 2.14 (m, 2H), 2.30 (m, 2H), 2.58 (m, 1H), 2.68 (d, 2H), 3.22 (q, 4H), 3.62 (m, 1H), 3.92 (d, 1H), 4.06 (d, 1H), 4.61 (d, 1H), 7.14 (dd, 1H), 7.22 (d, 1H), 7.40 (d, 1H), 7.48 (s, 1H), 7.70 (s, 2H), 8.48 (d, 2H).
LRMS : m/z (ES⁺) 546, 548 [MH⁺]

Example 158

N-(1-{2-[*(3S)*-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-piperidinyl)-*N*-methylacetamide




- 5 The title compound was obtained as a white foam in 89% yield, from the amine from example 150 and acetyl chloride, following a similar procedure to that described in example 157, except that dichloromethane was used as the reaction solvent.
- ¹Hnmr (CDCl₃, 400MHz) δ: (mixture of rotamers) 1.48 (m, 4H), 1.68 (m, 1H), 1.88 (s, 3H), 1.97 (m, 1H), 2.02 (2xs, 3H), 2.11 (m, 2H), 2.26 (m, 2H), 2.54 (m, 1H), 2.72 (s, 2H), 2.76 (s, 3H), 3.40, 4.36 (2xm, 1H), 3.91 (d, 1H), 4.58 (d, 1H), 7.10 (dd, 1H), 7.19 (d, 1H), 7.37 (d, 1H), 7.44 (s, 1H), 7.67 (s, 2H), 8.44 (s, 1H).
- 10 LRMS : m/z (ES⁺) 503, 505 [MH⁺]
- Microanalysis found: C, 60.66; H, 6.55; N, 10.89. C₂₆H₃₂Cl₂N₄O₂·0.5H₂O
- 15 requires C, 60.94; H, 6.49; N, 10.93%.

Example 159

N-(1-{2-[*(3S)*-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-piperidinyl)-*N,N*-diethyl-*N*-methylurea

20

123



Triethylamine (55 μ l, 0.40mmol) and diethylcarbamoyl chloride (50 μ l, 0.36mmol) were added to a solution of the amine from example 150 (152mg, 0.33mmol) in dichloromethane (5ml), and the solution stirred at room temperature for 2 hours.

- 5 Tlc analysis showed starting material remaining, so additional diethylcarbamoyl chloride (50 μ l, 0.36mmol) was added and the reaction stirred at room temperature for 18 hours. The mixture was concentrated under reduced pressure and the residue purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant, to afford the title compound as a white foam, 152mg.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.09 (t, 6H), 1.63 (m, 4H), 1.93 (m, 5H), 2.12 (m, 2H), 2.30 (m, 2H), 2.58 (m, 1H), 2.66 (s, 3H), 2.80 (m, 2H), 3.12 (q, 4H), 3.54 (m, 1H), 3.92 (d, 1H), 4.62 (d, 1H), 7.13 (dd, 1H), 7.22 (d, 1H), 7.40 (d, 1H), 7.48 (s, 1H), 7.70 (s, 2H), 8.48 (dd, 1H).


- 5 LRMS : m/z (ES^+) 560, 562 [MH^+]

Microanalysis found: C, 61.07; H, 7.04; N, 12.21. $\text{C}_{29}\text{H}_{39}\text{Cl}_2\text{N}_5\text{O}_2 \cdot 0.15\text{CH}_2\text{Cl}_2$ requires C, 61.07; H, 6.91; N, 12.22%.

20 Example 160

N-(1-(2-[(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinylethyl]-4-piperidinyl)methanesulfonamide

124




Triethylamine (43 μ l, 0.31mmol) and methanesulphonyl chloride (20 μ l, 0.26mmol) were added to an ice-cooled solution of the amine from example 149 (115mg, 0.26mmol) in dichloromethane (5ml), and the solution stirred at room temperature for an hour. The mixture was washed with water, then brine, dried (MgSO₄) and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) to afford the title compound.

¹Hnmr (CDCl₃, 300MHz) δ : 1.60 (m, 4H), 1.85-2.44 (m, 11H), 2.63 (m, 1H), 2.80 (m, 1H), 2.99 (s, 3H), 3.32 (m, 1H), 3.97 (d, 1H), 4.65 (d, 1H), 7.17 (m, 1H), 7.23 (d, 1H), 7.44 (d, 1H), 7.51 (s, 1H), 7.73 (m, 2H), 8.50 (d, 1H).

LRMS : m/z (ES⁺) 525, 527 [MH⁺]

15 Example 161

N-(1-(2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl)-4-piperidinyl)benzenesulfonamide



The title compound was obtained, from the amine from example 149 and phenylsulphonyl chloride, following the procedure described in example 160.

125


¹Hnmr (CDCl₃, 400MHz) δ: 1.33 (m, 3H), 1.47-2.17 (m, 8H), 2.18-2.36 (m, 2H), 2.52 (m, 3H), 3.09 (bs, 1H), 3.86 (d, 1H), 4.35 (bs, 1H), 4.60 (d, 1H), 7.10 (m, 1H), 7.18 (d, 1H), 7.28-7.59 (m, 5H), 7.66 (s, 2H), 7.81 (m, 2H), 8.43 (d, 1H).
LRMS : m/z (TSP⁺) 587.2, 589.2 [MH⁺]

- 5 Microanalysis found: C, 57.25; H, 5.54; N, 8.84. C₂₉H₃₂Cl₂N₄O₃S;1.2H₂O requires C, 57.18; H, 5.69; N, 9.20%.

10

Example 162

N-(1-(2-((3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinylethyl)-4-piperidinyl)-N-methylmethanesulfonamide



15

Triethylamine (0.21ml, 1.51mmol) followed by methanesulphonyl chloride (0.1ml, 1.28mmol) were added to an ice-cooled solution of the amine from example 150 (340mg, 0.6mmol) in dichloromethane (10ml) and the reaction stirred at room temperature for 2 hours. The mixture was washed with water, the aqueous wash extracted with dichloromethane and the combined organic solutions washed with brine, dried (MgSO₄) and concentrated under reduced pressure. The residual oil was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as eluant to afford the title compound as a white foam, 102mg.

- 25 ¹Hnmr (CD₃Cl, 400MHz) δ: 1.50 (m, 4H), 1.60 (m, 2H), 1.80-2.18 (m, 6H), 2.28 (m, 2H), 2.58 (m, 1H), 2.72 (m, 4H), 2.78 (s, 3H), 3.61 (m, 1H), 3.95 (d, 1H),


126

4.59 (d, 1H), 7.10 (m, 1H), 7.20 (m, 1H), 7.39 (m, 1H), 7.42 (s, 1H), 7.69 (m, 2H), 8.43 (d, 1H).

LRMS : m/z (TSP⁺) 539.1, 541.2 [MH⁺]

Example 163

N-[1-(1-[2-[(3*S*)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl]-3-azetidiny]-4-piperidinyl]methanesulfonamide




- 5 Triethylamine (127mg, 1.26mmol) and methanesulphonyl chloride (120mg, 1.05mmol) were added to a solution of the amine from example 147 (210mg, 0.42mmol) in dichloromethane (100ml), and the reaction stirred for 20 minutes. The mixture was washed with 0.88 ammonia (20ml), the layers separated, and the aqueous phase was extracted with dichloromethane (2x100ml). The
- 10 combined organic solutions were washed with brine, dried (MgSO₄), evaporated under reduced pressure and azeotroped with dichloromethane (4x100ml), to afford the title compound as a white foam, 210mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.56 (m, 2H), 1.87-2.04 (m, 5H), 2.08-2.39 (m, 6H), 2.46 (m, 1H), 2.62 (m, 3H), 2.81-3.00 (m, 5H), 3.32 (m, 1H), 3.60 (m, 2H), 3.92 (d, 1H), 4.57 (d, 1H), 4.67 (m, 1H), 7.16 (m, 1H), 7.23 (d, 1H), 7.43 (m, 2H), 7.73 (m, 2H), 8.49 (m, 1H).

LRMS : m/z (ES⁺) 580, 582 [MH⁺]

Example 164

5-(3,4-Dichlorophenyl)-5-[2-[(2-phenoxyethyl)amino]ethyl]-1-(2-pyridinyl)-2-piperidinone




7 Triethylamine was added to a suspension of 2-phenoxyethylamine in dichloromethane (8ml) until a solution was obtained, and then sufficient acetic acid was added to achieve a pH of 4. A solution of the aldehyde hydrochloride from preparation 11b (250mg, 0.63mmol) in dichloromethane (5ml) was added followed by sodium triacetoxyborohydride (132mg, 0.63mmol), and the reaction
10 stirred for 20 minutes. 4N Sodium hydroxide solution was added, the mixture stirred for 30 minutes, then filtered through a phase separation membrane. The organic filtrate was concentrated under reduced pressure, and the residue purified by column chromatography using dichloromethane:methanol:0.88 ammonia (95:5:0.5) as eluant. The product was redissolved in dichloromethane,
15 treated with 1N ethereal hydrochloric acid, and the mixture evaporated under reduced pressure to afford the title compound as a white solid, 199mg.

¹Hnmr (CD₃OD, 400MHz) δ: 2.20-2.40 (m, 3H), 2.43-2.60 (m, 2H), 2.73-2.92 (m, 2H), 2.90-3.05 (m, 1H), 3.40 (m, 2H), 4.15-4.27 (m, 3H), 4.46 (d, 1H), 6.88-7.03 (m, 3H), 7.28 (dd, 2H), 7.44 (d, 1H), 7.59 (d, 1H), 7.68-7.75 (m, 2H), 7.93 (d,
20 1H), 8.43 (dd, 1H), 8.60 (d, 1H).

LRMS : m/z (TSP⁺) 484.1, 486.1 [MH⁺]



Examples 165 to 167

The following examples of general structure:



- 5 were prepared as white solids, from the aldehyde hydrochloride from preparation 11b and the appropriate amine, according to the procedure described in example 164.

Example	R	Yield	Data
165 ^a		37	¹ Hnmr (CD ₃ OD, 400MHz) δ: 2.17-2.38 (m, 3H), 2.40-2.57 (m, 2H), 2.70-2.78 (m, 1H), 2.84 (m, 1H), 3.01 (m, 1H), 3.41 (t, 2H), 4.17 (d, 1H), 4.26 (t, 2H), 4.50 (d, 1H), 6.97-7.16 (m, 4H), 7.45 (d, 1H), 7.52-7.60 (m, 2H), 7.73 (d, 1H), 7.80 (d, 1H), 8.20 (dd, 1H), 8.58 (d, 1H). LRMS : m/z (TSP ⁺) 504.2, 506.2 [MH ⁺]

166 ^b		39	<p>¹Hnmr (CD₃OD, 400MHz) δ: 2.18-2.42 (m, 3H), 2.42-2.60 (m, 2H), 2.72-2.89 (m, 2H), 2.95 (m, 1H), 3.37 (t, 2H), 4.15-4.36 (m, 3H), 4.43 (d, 1H), 6.60-6.82 (m, 3H), 7.24 (dd, 1H), 7.40 (d, 1H), 7.52 (d, 1H), 7.68 (d, 1H), 7.75 (dd, 1H), 7.99 (d, 1H), 8.50 (dd, 1H), 8.55 (d, 1H).</p> <p>Microanalysis found: C, 50.39; H, 5.26; N, 6.60.</p> <p>C₂₆H₂₈Cl₂FN₃O₂·2HCl·2.5H₂O requires C, 49.78; H, 5.30; N, 6.70%.</p>
167 ^c		17	<p>¹Hnmr (CD₃OD, 400MHz) δ: 2.21-2.46 (m, 3H), 2.46-2.61 (m, 2H), 2.72-2.90 (m, 2H), 2.99 (m, 1H), 3.39 (t, 2H), 4.12-4.35 (m, 3H), 4.46 (d, 1H), 6.93 (m, 2H), 7.02 (m, 2H), 7.46 (d, 1H), 7.60 (d, 1H), 7.74 (s, 1H), 7.77 (dd, 1H), 8.01 (d, 1H), 8.51 (dd, 1H), 8.60 (d, 1H).</p> <p>Microanalysis found: C, 50.01; H, 5.36; N, 6.89.</p> <p>C₂₆H₂₆Cl₂FN₃O₂·2HCl·2.5H₂O requires C, 49.78; H, 5.30; N, 6.70%.</p>

Starting amines:

a = 2-(2-(2-fluorophenoxy)ethyl)amine hydrochloride as prepared in WO 0020401


b = 2-(2-(3-fluorophenoxy)ethyl)amine hydrochloride from preparation 19

5 c = 2-(2-(4-fluorophenoxy)ethyl)amine hydrochloride as prepared in WO 0020401

131

Example 168


5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-oxo-1-piperidinyl)-1-azetidiny]ethyl}-2-piperidinone



- J Dess-Martin periodinane (165mg, 0.39mmol) was added to a solution of the alcohol from example 132 (200mg, 0.386mmol) in dichloromethane (10ml), and the solution stirred at room temperature for an hour. Tlc analysis showed starting material remaining, so additional Dess-Martin periodinane (82.5mg, 0.19mmol) was added and the reaction stirred for a further 30 minutes. Sodium
- 10 thiosulphate (100mg), and aqueous sodium bicarbonate solution (10ml) were added, and the mixture stirred for 10 minutes. The layers were separated, the organic phase dried (Na_2SO_4) and evaporated under reduced pressure. The residual gum was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (95:5:0.5 to
- 5 90:10:1) to afford the title compound as a white foam, 125mg.
- $^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 1.60-2.60 (m, 25H), 3.95 (d, 1H), 4.50 (d, 1H), 7.00 (d, 1H), 7.24 (d, 1H), 7.44 (d, 2H), 7.60 (m, 2H).
- LRMS : m/z (TSP $^+$) 515.1, 517.2 [MH^+]

Example 169


(5S)-5-(3,4-Dichlorophenyl)-1-(4-methyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone



- 5 The title compound was prepared as a yellow solid in 32% yield from the aldehyde from preparation 15 and 3-morpholinoazetidine dihydrochloride (WO 9725322), following a similar procedure to that described in example 131.
- $^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.82 (m, 1H), 1.97 (m, 1H), 2.12 (m, 2H), 2.18-2.27 (m, 10H), 2.57 (m, 2H), 2.88-3.10 (m, 3H), 3.63 (m, 5H), 3.88 (d, 1H), 4.42 (d, 1H), 6.96 (d, 1H), 7.20 (m, 1H), 7.40 (m, 2H), 7.48 (s, 1H), 8.30 (d, 1H).
- 10 LRMS : m/z (TSP $^+$) 503.2, 504.9 [MH^+]

Example 170

- 15 (5S)-5-(3,4-Dichlorophenyl)-1-(3-methyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone



- The title compound was prepared as a yellow solid in 8% yield from the aldehyde from preparation 13 and 3-morpholinoazetidine dihydrochloride (WO 9725322), following a similar procedure to that described in example 131.
- 20

133


¹Hnmr (CDCl₃, 400MHz) δ: 1.72-1.92 (m, 2H), 2.01 (m, 2H), 2.12-2.40 (m, 8H), 2.57 (m, 1H), 2.80 (m, 2H), 2.98 (m, 1H), 3.43 (m, 3H), 3.70 (m, 5.5H), 3.90 (m, 0.5H), 4.22 (m, 0.5H), 4.42 (m, 0.5H), 7.19 (m, 2H), 7.38 (m, 0.5H), 7.44 (m, 1H), 7.58 (m, 1H), 7.80 (m, 0.5H), 8.40 (m, 1H).

5

LRMS : m/z (TSP⁺) 503.6, 505.7 [MH⁺]

Example 171

10 (5S)-5-(3,4-Dichlorophenyl)-5-[2-(4-phenyl-3,6-dihydro-1(2H)-pyridinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone



The title compound was prepared in 35% yield as a white foam, from the aldehyde hydrochloride from preparation 11b and 1,2,3,6-tetrahydro-4-phenylpyridine, following a similar procedure to that described in example 17.

15


¹Hnmr (CDCl₃, 400MHz) δ: 1.98-2.38 (m, 7H), 2.50 (m, 2H), 2.58 (m, 3H), 3.01 (s, 2H), 3.95 (d, 1H), 4.62 (d, 1H), 5.98 (s, 1H), 7.15 (m, 1H), 7.20-7.38 (m, 7H), 7.40 (d, 1H), 7.52 (s, 1H), 7.72 (d, 2H), 8.47 (d, 1H).

LRMS : m/z (TSP⁺) 506.1, 508.0 [MH⁺]

20

Example 172

(5S)-5-(3,4-Dichlorophenyl)-5-[2-(1,1-dioxido-4-thiomorpholinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone



- 5 The title compound was prepared in 83% yield as a white foam, from the aldehyde hydrochloride from preparation 11b and thiomorpholine 1,1-dioxide (WO 9605193) following a similar procedure to that described in example 17. ¹Hnmr (CDCl₃, 400MHz) δ: 1.90 (t, 2H), 2.10-2.38 (m, 5H), 2.58 (m, 1H), 2.78 (m,

10

2H), 2.86 (m, 2H), 2.99 (m, 4H), 3.88 (d, 1H), 4.90 (d, 1H), 7.18 (m, 2H), 7.41 (d, 1H), 7.45 (s, 1H), 7.72 (dd, 1H), 7.80 (d, 1H), 8.52 (d, 1H).

LRMS : m/z (TSP⁺) 482.1, 484.1 [MH⁺]


Microanalysis found: C, 54.53; H, 5.31; N, 8.50. C₂₂H₂₅Cl₂N₃O₃S requires C,

15 54.77; H, 5.22; N, 8.71%.

Example 173

(5S)-5-(3,4-Dichlorophenyl)-5-[2-(2,6-dimethyl-4-morpholinyl)ethyl]-1-(2-pyridinyl)-2-piperidinone

20



135

The title compound was prepared in 30% yield as a white foam, from the aldehyde hydrochloride from preparation 11b and 2,6-dimethylmorpholine following a similar procedure to that described in example 17.


¹Hnmr (CDCl₃, 400MHz) δ: 1.06 (2xd, 6H), 1.55 (m, 2H), 1.92 (m, 3H), 2.02-2.18 (m, 2H), 2.26 (m, 2H), 2.55 (m, 3H), 3.54 (m, 2H), 3.90 (d, 1H), 4.58 (d, 1H), 7.11 (m, 1H), 7.19 (d, 1H), 7.38 (d, 1H), 7.42 (s, 1H), 7.66 (m, 2H), 8.45 (m, 1H).

LRMS : m/z (ES⁺) 462, 464 [MH⁺]

10

Example 174


(5S)-5-{2-[3-(4-Amino-1-piperidinyl)-1-azetidiny]ethyl}-5-(3,4-dichlorophenyl)-1-(6-methyl-2-pyridinyl)-2-piperidinone tetratrifluoroacetate



- 15 A mixture of the protected amine from preparation 85 (59mg, 0.1mmol) and trifluoroacetic acid (0.5ml) in dichloromethane (3ml) was stirred at room temperature for 1 hour. The mixture was concentrated under reduced pressure, and the residue azeotroped with toluene (3x) and diethyl ether to afford the title compound as a brown solid, 60mg.
- 20 ¹Hnmr (CD₃OD, 400MHz) δ: 1.74 (m, 2H), 1.99-2.32 (m, 8H), 2.40 (m, 1H), 2.58 (m, 5H), 2.84-3.30 (m, 6H), 3.53 (m, 1H), 3.98 (d, 1H), 4.06 (m, 1H), 4.20 (m, 2H), 4.48 (d, 1H), 7.20 (d, 1H), 7.30 (d, 1H), 7.45 (d, 1H), 7.60 (d, 1H), 7.78 (dd 1H), 7.83 (s, 1H).
- Microanalysis found: C, 42.44; H, 4.38; N, 6.48.
- 25 C₂₇H₃₅Cl₂N₅O; 4CF₃CO₂H; 0.2(C₂H₅)₂O; 1.5H₂O requires C, 42.39; H, 4.37; N, 6.90%.

Example 175

5-(3,4-difluorophenyl)-1-(6-methyl-2-pyridinyl)-5-(2-[3-(4-morpholinyl)-1-azetidiny]ethyl)-2-piperidinone



5

The title compound was prepared as a white foam in 63% yield from the aldehyde from preparation 12c and 3-morpholinoazetidine dihydrochloride (WO 9725322), following a similar procedure to that described in example 112, except dichloromethane:methanol:0.88 ammonia (96:4:0.4) was used as the column eluant.

10

¹Hnmr (CDCl₃, 400MHz) δ: 1.50-2.00 (m, 6H), 2.05-2.46 (m, 7H), 2.58 (s, 3H), 2.80 (m, 1H), 2.95 (m, 1H), 3.44 (m, 2H), 3.64 (m, 4H), 3.86 (d, 1H), 4.44 (d, 1H), 6.97(d, 1H), 7.12 (m, 2H), 7.24 (m, 1H), 7.41 (d, 1H), 7.58 (t, 1H).


LRMS : m/z (TSP⁺) 471.3 [MH⁺]

15 Microanalysis found: C, 65.42; H, 6.90; N, 11.68. C₂₆H₃₂F₂N₄O₂; 0.4H₂O; 0.1 Et₂O requires C, 65.36; H, 7.02; N, 11.55.

137

Example 176

5-(3,4-difluorophenyl)-5-(2-[4-hydroxy-4-phenyl-1-piperidinyl]ethyl)-1-(6-methyl-2-pyridinyl)-2-piperidinone




- 5 The title compound was prepared as a white solid in 82% yield from the aldehyde from preparation 12c and 4-hydroxy-4-phenylpiperidine, following a similar procedure to that described in example 1, except using a elution gradient of dichloromethane:methanol:0.88 ammonia (97:3:0.3-96:4:0.4).
- ¹Hnmr (CDCl₃, 400MHz) δ: 1.55 (m, 3H), 1.72 (m, 2H), 1.90-2.40 (m, 10H), 2.44-2.80 (m, 5H), 3.94 (d, 1H), 4.61 (d, 1H), 6.99(d, 1H), 7.15 (m, 2H), 7.27 (m, 1H), 7.36 (m, 3H), 7.44 (m, 3H), 7.60 (t, 1H).
- 10 LRMS : m/z (TSP⁺) 506.3 [MH⁺]
- Microanalysis found: C, 69.78; H, 6.63; N, 8.13. C₃₀H₃₃F₂N₃O₂; 0.3H₂O; 0.1 CH₂Cl₂ requires C, 69.59; H, 6.56; N, 8.09.

15

Example 177

5-(3,4-difluorophenyl)-5-(2-[1,4-dioxo-8-azaspiro[4.5]dec-8-yl]ethyl)-1-(6-methyl-2-pyridinyl)-2-piperidinone



20

The title compound was prepared as a white foam in 69% yield from the aldehyde from preparation 12c and 1,4-dioxa-8-azaspiro[4.5]decane, following a similar procedure to that described in example 1, except using a elution gradient of dichloromethane:methanol:0.88 ammonia (97:3:0.3-95:5:0.5).

5 $^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.40-2.80 (m, 19H), 3.92 (m, 5H), 4.60 (m, 1H), 6.96 (d, 1H), 7.12 (m, 2H), 7.24 (m, 1H), 7.40 (d, 1H), 7.57 (t, 1H).


LRMS : m/z (TSP $^+$) 472.5 [MH^+]

Microanalysis found: C, 65.35; H, 6.74; N, 8.70. $\text{C}_{26}\text{H}_{31}\text{F}_2\text{N}_3\text{O}_3$; $0.4\text{H}_2\text{O}$; $0.05\text{Et}_2\text{O}$ requires C, 65.23; H, 6.75; N, 8.71.

10

Preparation 1

(5S)-5-(3,4-Dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-1-(2-pyridinyl)-2-piperidinone



15

Potassium *tert*-butoxide (68g, 0.606mol) was added to a suspension of (5S)-5-(3,4-dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-2-piperidinone (WO 9807722) (200g, 0.606mol) in 1,2-dimethoxyethane (700ml), and the mixture heated under reflux for 1

20

hour. 2-Fluoropyridine (59g, 0.606mol) was then added and the mixture stirred under reflux for 1 hour. Additional potassium *tert*-butoxide (34g, 0.303mol) and 2-fluoropyridine (30g, 0.303mol) were added and the reaction mixture stirred under reflux for a further hour. The cooled mixture was partitioned with water (500ml), and the aqueous layer then extracted with ethyl acetate (2x500ml). The combined organic extracts were dried (MgSO_4), and evaporated under reduced pressure. The crude product was purified by column chromatography

25


139

on silica gel using an elution gradient of ethyl acetate:pentane (20:80 to 100:0) to afford the title compound as a clear gum, 129g.

$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.98 (dd, 1H), 2.16-2.39 (m, 4H), 2.55-2.70 (m, 1H), 3.68 (m, 2H), 3.85 (m, 2H), 3.98 (d, 1H), 4.43 (t, 1H), 4.70 (d, 1H), 7.15 (dd, 1H), 7.29 (dd, 1H), 7.42 (d, 1H), 7.53 (s, 1H), 7.65 (m, 2H), 8.52 (d, 1H).
LRMS : m/z (TSP $^+$) 407.2, 409.5 [MH^+]

Preparation 2

10 (5S)-5-(3,4-Dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-1-(6-methyl-2-pyridinyl)-2-piperidinone



A mixture of (5S)-5-(3,4-dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-2-piperidinone (WO 9807722) (6.5g, 19.7mmol), potassium carbonate (3.05g, 21.7mmol), copper (I) iodide (400mg, 2.1mmol) and 2-bromo-6-methylpyridine (10.2g, 60mmol) in 1-methyl-2-pyrrolidinone (200ml) was stirred at 140°C for 24 hours. The cooled mixture was partitioned between ethyl acetate and 10% aqueous ammonia, and the layers separated. The aqueous phase was extracted with ethyl acetate, and the combined organic extracts were washed with water, then brine (3x), dried (MgSO_4) and evaporated under reduced pressure to give a gum. The crude product was purified


by column chromatography on silica gel using an elution gradient of ethyl acetate:pentane (0:100 to 100:0) to afford the title compound as a brown solid, 3.02g.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 2.00 (m, 1H), 2.15-2.38 (m, 4H), 2.58 (m, 4H), 3.69 (m, 2H), 3.85 (m, 2H), 3.95 (d, 1H), 4.45 (t, 1H), 4.62 (d, 1H), 7.00 (d, 1H), 7.26 (m, 1H), 7.40 (dd, 2H), 7.61 (m, 2H).

140

LRMS : m/z (TSP⁺) 421.0, 423.0 [MH⁺]**Preparation 2a**

5-(3,4-Difluorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-1-(6-methyl-2-pyridinyl)-2-
piperidinone



The title compound was obtained as a brown foam in 63% yield from 5-(3,4-difluorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-2-piperidinone (EP 992493) and 2-bromo-6-methylpyridine, following the procedure described in preparation 2.


¹Hnmr (CDCl₃, 400MHz) δ : 1.98 (m, 1H), 2.15-2.38 (m, 4H), 2.60 (m, 4H), 3.69 (m, 2H), 3.85 (m, 2H), 3.95 (d, 1H), 4.45 (t, 1H), 4.62 (d, 1H), 7.00 (d, 1H), 7.18 (m, 2H), 7.38 (m, 2H), 7.61 (t, 1H).

LRMS : m/z (TSP⁺) 389.1 [MH⁺]

15





Preparations 3 to 6


The following compounds of the general formula:



were prepared from (5S)-5-(3,4-dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-2-piperidinone (WO 9807722) and the appropriate bromide, according to the procedure described in preparation 2.


141

Prep. No.	R	Yield (%)	Data
3		17	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.90-2.35 (m, 9H), 3.62 (m, 2.5H), 3.80 (m, 2H), 4.04 (d, 0.5H), 4.22 (d, 0.5H), 4.36 (t, 1H), 4.50 (d, 0.5H), 7.18 (m, 1.5H), 7.40 (m, 1.5H), 7.55 (m, 1.5H), 7.80 (s, 0.5H), 8.36 (d, 1H). LRMS : m/z (TSP ⁺) 421.4, 423.3 [MH ⁺]
4		28	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.92 (m, 1H), 2.18-2.38 (m, 7H), 2.56 (m, 1H), 3.64 (m, 2H), 3.84 (m, 2H), 3.92 (dd, 1H), 4.40 (m, 1H), 4.58 (m, 1H), 7.12 (m, 2H), 7.36 (dd, 1H), 7.44 (m, 2H), 8.28 (s, 1H). LRMS : m/z (TSP ⁺) 421.1, 423.4 [MH ⁺]
5		19	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.94 (dd, 1H), 2.22 (m, 4H), 2.36 (s, 3H), 2.58 (m, 1H), 3.62 (m, 2H), 3.84 (m, 2H), 3.94 (d, 1H), 4.40 (t, 1H), 4.60 (d, 1H), 6.95 (d, 1H), 7.22 (dd, 1H), 7.40 (d, 2H), 7.48 (s, 1H), 8.35 (d, 1H). LRMS : m/z (TSP ⁺) 421.1, 422.7 [MH ⁺]
6		16	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.32 (t, 3H), 1.93 (dd, 1H), 2.20 (m, 4H), 2.56 (m, 1H), 2.80 (q, 2H), 3.68 (m, 2H), 3.82 (m, 3H), 4.41 (t, 1H), 4.80 (d, 1H), 6.96 (d, 1H), 7.30 (d, 1H), 7.38 (m, 2H), 7.58 (dd, 1H), 7.62 (d, 1H). LRMS : m/z (TSP ⁺) 435.2, 437.2 [MH ⁺]

Preparation 7**2-Bromo-6-methoxypyridine**

Dried sodium (970mg, 42.2mmol) was added portionwise to cooled methanol
5 (50ml) under nitrogen, and once addition was complete, 2,6-dibromopyridine
(10g, 42.2mmol) was added portionwise. The resulting suspension was heated
under reflux for 24 hours. The cooled reaction was concentrated under reduced
pressure, the residue diluted with water (100ml), and extracted with ethyl
acetate (2x75ml). The organic extracts were dried (MgSO₄), and evaporated
10 under reduced pressure to give a pale yellow oil. The crude product was
purified by column chromatography on silica gel using an elution gradient of
pentane:ethyl acetate (100:0 to 98:2) to give the title compound as a clear oil,
4.23g.

¹Hnmr (CDCl₃, 400MHz) δ: 3.93 (s, 3H), 6.67 (d, 1H), 7.05 (d, 1H), 7.40 (dd,
15 1H).

Preparation 8**(5S)-5-(3,4-Dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-1-(6-methoxy-2-
pyridinyl)-2-piperidinone**

A mixture of (5S)-5-(3,4-dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-2-
piperidinone (WO 9807722) (4.7g, 14.3mmol), potassium carbonate (2.97g,
25 21.4mmol), copper (I) iodide (3.0g, 15.7mmol) and the bromide from
preparation 7 (4.03g, 21.4mmol) in 1-methyl-2-pyrrolidinone (25ml) was stirred

between dichloromethane (600ml) and water (300ml), the layers separated, and the aqueous phase extracted with further dichloromethane (2x100ml). The combined organic extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give a pale yellow oil. This oil was
5 triturated with ether, and the resulting precipitate was filtered and dried to afford the title compound as a solid, 72g.

¹Hnmr (CDCl₃, 400MHz) δ: 2.04 (t, 2H), 2.15-2.40 (m, 3H), 2.60 (m, 1H), 3.17 (d, 6H), 3.92 (d, 1H), 4.01 (t, 1H), 4.68 (d, 1H), 7.15 (t, 1H), 7.24 (d, 1H), 7.43 (d, 1H), 7.52 (s, 1H), 7.68 (m, 2H), 8.52 (d, 1H).


10 LRMS : m/z (ES⁺) 409.0, 411.0 [MH⁺]

Preparation 10

(5S)-5-(3,4-Dichlorophenyl)-5-(2,2-dimethoxyethyl)-1-(6-methoxy-2-pyridinyl)-2-


15

piperidinone



The title compound was obtained as a golden oil (86%) from the acetal of preparation 8, following a similar procedure to that described in preparation 9.

¹Hnmr (CDCl₃, 400MHz) δ: 1.95-2.30 (m, 5H), 2.58 (m, 1H), 3.17 (s, 3H), 3.19 (s, 3H), 3.72 (d, 1H), 3.92 (m, 1H), 4.00 (s, 3H), 4.90 (dd, 1H), 6.61 (d, 1H),
20 7.25 (d, 1H), 7.30 (dd, 1H), 7.42 (d, 1H), 7.61 (dd, 1H), 7.68 (s, 1H).

Preparation 11a**[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]acetaldehyde**

A solution of the acetal from preparation 9 (72g, 0.176mol) in tetrahydrofuran (250ml) was added to an ice-cooled solution of hydrochloric acid (2N, 880ml), and the solution stirred at room temperature for 18 hours. The mixture was re-cooled in ice, neutralised by the addition of sodium bicarbonate, then basified to pH 9 using 2N sodium hydroxide solution. This aqueous solution was extracted with ethyl acetate (2x1.5L), the combined extracts washed with 2N sodium hydroxide (5x300ml), dried (MgSO₄) and evaporated under reduced pressure to afford the title compound as a pale yellow gum, 47g.

¹Hnmr (CDCl₃, 300MHz) δ: 2.20-2.46 (d, 3H), 2.63 (m, 1H), 2.79 (d, 1H), 2.95 (d, 1H), 4.09 (d, 1H), 4.70 (d, 1H), 7.16 (m, 1H), 7.30 (m, 1H), 7.44 (d, 1H), 7.56 (s, 1H), 7.72 (m, 2H), 8.48 (d, 1H), 9.55 (s, 1H).

Preparation 11b**[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]acetaldehyde hydrochloride**


Amberlyst® 15 resin (13g) was added to a solution of the acetal from preparation 1 (1.34g, 3.4mmol) in methanol (50ml), and the reaction stirred at room temperature for 18 hours. The mixture was filtered, the resin washed with a solution of dichloromethane:methanol:0.88 ammonia (90:10:1), and the combined filtrate evaporated under reduced pressure. The residue was partitioned between diethyl ether and sodium hydroxide solution, the layers separated, and the ether extract was washed with brine, dried (MgSO₄) and

evaporated under reduced pressure. The residue was dissolved in hydrochloric acid (3N), and the solution stirred at room temperature for 2 hours. The solution was carefully basified using 2N sodium hydroxide solution, and extracted with diethyl ether (3x). The combined organic extracts were dried (MgSO₄) and concentrated under reduced pressure. The product was treated with ethereal hydrochloric acid, then evaporated under reduced pressure, to afford the title compound as a white foam, 460mg.

¹Hnmr (CDCl₃, 300MHz) δ: 2.42 (t, 2H), 2.55 (m, 1H), 2.75 (m, 1H), 3.10 (m, 3H), 4.55 (d, 1H), 4.70 (d, 1H), 7.34 (m, 1H), 7.46 (m, 2H), 7.58 (dd, 1H), 7.90 (d, 1H), 8.20 (dd, 1H), 8.64 (d, 1H), 9.58 (s, 1H).

Preparation 12a

[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(6-methyl-2-pyridinyl)piperidinyl]acetaldehyde



15

5N Hydrochloric acid (40ml) was added dropwise to an ice-cooled solution of the acetal from preparation 2 (3.8g, 9mmol) in tetrahydrofuran (40ml), and the reaction stirred at room temperature for 24 hours. The mixture was evaporated, and the residue neutralised by the addition of aqueous sodium bicarbonate solution. The aqueous solution was extracted with ethyl acetate (2x), the combined organic extracts dried (MgSO₄), and evaporated under reduced pressure to give the title compound, 2.05g.

¹Hnmr (CDCl₃, 300MHz) δ : 2.24 (m, 1H), 2.38 (m, 2H), 2.56 (s, 3H), 2.58 (m, 1H), 2.88 (dd, 1H), 2.96 (dd, 1H), 4.05 (d, 1H), 4.66 (d, 1H), 7.00 (d, 1H), 7.30 (d, 1H), 7.42 (d, 1H), 7.44 (d, 1H), 7.60 (dd, 1H), 7.65 (d, 1H), 9.56 (s, 1H).


Preparation 12b**[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(6-methyl-2-pyridinyl)piperidinyl]acetaldehyde hydrochloride**

Amberlyst® 15 resin (320g) was added to an ice-cooled solution of the acetal
5 from preparation 2 (156g, 352mmol) in methanol (1L), and the reaction then
stirred at room temperature for 20 hours. The mixture was filtered through
Arbocel®, and the resin washed with a (80:20) methanolic ammonia solution
(4x1L) and then dichloromethane (2x1L), filtering between each wash. The
combined filtrates were concentrated under reduced pressure and the residue
10 partitioned between water (1L) and ethyl acetate (2L), and the layers separated.
The organic phase was washed with water (2x500ml), and the combined
aqueous solutions extracted with ethyl acetate (500ml). This organic extract
was washed with brine, dried (MgSO₄) and evaporated under reduced pressure
to give a brown oil, 136g.

15 ¹Hnmr (CDCl₃, 300MHz) δ : 2.02 (m, 2H), 2.15-2.39 (m, 3H), 2.59 (m, 4H), 3.18
(s, 3H), 3.22 (s, 3H), 3.83 (d, 1H), 4.00 (m, 1H), 4.78 (dd, 1H), 7.00 (dd, 1H),
7.30 (dd, 1H), 7.42 (m, 2H), 7.60 (dd, 1H), 7.65 (d, 1H).

A solution of the oil in tetrahydrofuran (280ml) was added to an ice-cooled
solution of 3N hydrochloric acid (530ml), and the solution stirred at room
20 temperature for 3 hours. The solution was diluted with ethyl acetate (500ml),
and neutralised using sodium bicarbonate. 2N Sodium hydroxide was added to
give pH 9, the phases separated, and the organic layer washed with 2N sodium
hydroxide (3x100ml). The combined aqueous washes were extracted with ethyl
acetate, then the combined organic solutions washed with brine, dried (MgSO₄)
25 and evaporated under reduced pressure. The residual oil was purified by
column chromatography on silica gel using ethyl acetate as eluant to give a
yellow oil, 94g. This was dissolved in dichloromethane, and treated with an
excess of ethereal hydrochloric acid, and the solution evaporated under
reduced pressure. The residue was azeotroped with dichloromethane, to give
30 the title compound as a yellow foam.


¹Hnmr (CDCl₃, 400MHz) δ : 2.27-2.52 (m, 3H), 2.66 (m, 1H), 2.94 (s, 3H), 3.17
(s, 2H), 4.38 (d, 1H), 4.68 (d, 1H), 7.22 (d, 1H), 7.44 (m, 4H), 8.15 (dd, 1H),
9.52 (s, 1H).

Preparation 12c**[3-(3,4-difluorophenyl)-6-oxo-1-(6-methyl-2-pyridinyl)piperidinyl]acetaldehyde**

- 5 6N Hydrochloric acid (16ml) was added dropwise to an ice-cooled solution of the acetal from preparation 2a (3.42g, 9mmol) in methanol (40ml), and the reaction stirred at room temperature for 24 hours. The mixture was evaporated, and the residue neutralised by the addition of aqueous sodium bicarbonate solution. The aqueous solution was extracted with ethyl acetate (2x), the
- 10 combined organic extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give a yellow oil, 3.5g. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (98:2:0.2) as eluant to give the title compound as a yellow gum, (750mg).
- 15 ¹Hnmr (CDCl₃, 300MHz) δ : 2.21-2.42 (m, 3H), 2.48-2.64 (m, 4H), 2.78 (dd, 1H), 2.96 (dd, 1H), 4.05 (d, 1H), 4.66 (d, 1H), 7.00 (d, 1H), 7.18 (m, 2H), 7.36 (m, 1H), 7.46 (d, 1H), 7.60 (m, 1H), 9.55 (s, 1H).
- LRMS: m/z (TSP+) 345.1 [MH⁺].

Preparations 13 to 16


The following compounds of the general formula:



5 were prepared from the corresponding acetals following the procedure described in preparation 12a.

Prep. No.	R	Yield (%)	Data
13		69	LRMS : m/z (TSP ⁺) 377.2, 378.7 [MH ⁺]
14		78	LRMS : m/z (TSP ⁺) 377.5, 379.5 [MH ⁺]
15		56	¹ Hnmr (CDCl ₃ , 400MHz) δ: 2.00-2.46 (m, 6H), 2.58 (m, 1H), 2.80 (m, 1H), 3.02 (m, 1H), 4.05 (m, 1H), 4.30 (d, 0.5H), 4.60 (d, 0.5H), 7.20 (m, 1H), 7.41-7.83 (m, 4H), 8.39 (m, 1H), 9.55 (s, 1H). LRMS : m/z (TSP ⁺) 377.2, 378.2 [MH ⁺]


150

16		90	¹ Hnmr (CDCl ₃ , 400MHz) δ: 1.30 (t, 3H), 2.20-2.40 (m, 2H), 2.38 (m, 2H), 2.60 (m, 1H), 2.80 (q, 2H), 2.92 (d, 1H), 4.00 (d, 1H), 4.76 (d, 1H), 6.96 (d, 1H), 7.32 (dd, 1H), 7.40 (d, 1H), 7.44 (d, 1H), 7.60 (dd, 1H), 7.74 (d, 1H), 9.54 (s, 1H). LRMS : m/z (TSP ⁺) 391.0, 393.2 [MH ⁺]
----	---	----	--

Preparation 17

[(3S)-3-(3,4-Dichlorophenyl)-1-[6-(dimethylamino)-2-pyridinyl]-6-oxopiperidinyl]acetaldehyde hydrochloride

5



Copper (I) iodide (6.4g, 33.3mmol), potassium carbonate (6.3g, 45.4mmol) and 6-bromo-2-(dimethylamino)pyridine (WO 9843971) (9g, 45.4mmol) were added consecutively to a solution of (5S)-5-(3,4-dichlorophenyl)-5-(1,3-dioxolan-2-ylmethyl)-2-piperidinone (WO 9807722) (10g, 30.3mmol), in 1-methyl-2-pyrrolidinone (50ml), and the mixture was stirred at 140°C for 4 hours. The cooled mixture was poured into 4N hydrochloric acid; then carefully basified using 10% aqueous ammonia. The aqueous mixture was extracted with ethyl acetate (3x200ml), and the combined organic extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give an orange oil. The crude product was purified by column chromatography on silica gel using methanol:dichloromethane (5:95) as eluant, and repeated using ethyl acetate as eluant. The product was treated with 1N ethereal hydrochloric acid, and evaporated under reduced pressure to afford the title compound.

151


¹Hnmr (CDCl₃, 400MHz) δ: 2.21-2.43 (m, 2H), 2.55-2.81 (m, 2H), 3.05 (d, 1H), 3.18 (d, 1H), 3.38 (s, 6H), 4.39 (q, 2H), 6.90 (d, 1H), 6.95 (d, 1H), 7.25 (d, 1H), 7.42 (d, 1H), 7.48 (s, 1H), 7.86 (dd, 1H), 9.53 (s, 1H).

LRMS : m/z (TSP⁺) 406.1, 408.1 [MH⁺]

5

Preparation 18

{(3S)-3-(3,4-Dichlorophenyl)-1-[6-methoxy-2-pyridinyl]-6-oxopiperidinyl}acetaldehyde hydrochloride




10

A solution of the acetal from preparation 10 (2.47g, 5.62mmol) in tetrahydrofuran (10ml) was added dropwise to cooled (5°C) hydrochloric acid (14ml, 2N, 28mmol), and the reaction stirred at room temperature for 18 hours. Tlc analysis showed starting material remaining, so additional hydrochloric acid (10ml, 2N), and tetrahydrofuran (10ml) were added and the reaction stirred for a further 24 hours at room temperature. The solution was cooled in ice, neutralised by the addition of sodium bicarbonate, and basified by the addition of 1N sodium hydroxide solution (10ml). The mixture was extracted with ethyl acetate (3x50ml), and the combined organic extracts washed with brine (2x20ml), dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using ethyl acetate:pentane (75:25) as eluant, to afford the title compound as a foam, 760mg.

¹Hnmr (CDCl₃, 400MHz) δ: 2.24-2.40 (m, 3H), 2.60 (m, 1H), 2.70-3.00 (ABq, 2H), 3.95 (d, 1H), 3.98 (s, 3H), 4.81 (d, 1H), 6.62 (d, 1H), 7.27 (d, 1H), 7.35 (dd, 1H), 7.45 (d, 1H), 7.63 (dd, 1H), 7.69 (d, 1H), 9.52 (s, 1H).


25

Preparation 19**2-(3-Fluorophenoxy)ethylamine hydrochloride**

A solution of 3-fluorophenol (20g, 178mmol) in tetrahydrofuran (100ml) and
5 N,N-dimethylformamide (100ml), was added dropwise to a cooled (10°C)
suspension of sodium hydride (13.38g, 80% dispersion in mineral oil, 446mmol)
in tetrahydrofuran (200ml). Once addition was complete, the mixture was stirred
at room temperature for 45 minutes. 2-Bromoethylamine hydrobromide (36.56g,
178mmol) was added portionwise over 30 minutes, and then the reaction stirred
10 at 45°C for 18 hours. Water (800ml) was carefully added to the cooled solution,
and the mixture extracted with ethyl acetate (3x250ml). The organic solutions
were extracted with 2M hydrochloric acid (3x200ml), and these acidic fractions
then basified to pH 10 using 2N sodium hydroxide solution. This was re-
extracted with ethyl acetate (4x250ml), these combined organic solutions dried
15 (MgSO₄) and evaporated under reduced pressure. The residual oil was
dissolved in ethyl acetate (200ml), and the solution treated with 1N ethereal
hydrochloric acid (150ml), and the suspension stirred for 2 hours. The resulting
precipitate was filtered and dried *in vacuo*, to afford the title compound as a
white solid, 8.0g.

20 ¹Hnmr (CD₃OD, 400MHz) δ: 3.35 (s, 2H), 4.20 (s, 2H), 6.65-6.82 (m, 3H), 7.25
(m, 1H).


LRMS : m/z (ES⁺) 156 [MH⁺]

Preparation 20**1-Methyl-2-oxo-1,2-dihydro-4-pyridinecarbonitrile**

- 5 Oxalyl chloride (250 μ l, 2.87mmol) was added to an ice-cooled solution of N,N-dimethylformamide (243 μ l, 3.14mmol) in acetonitrile (3ml). A suspension of 1-methyl-2-oxo-1,2-dihydro-4-pyridinecarboxamide (J.O.C. 24; 1959; 196) (201.5mg, 1.32mmol) and pyridine (470 μ l, 5.82mmol) in acetonitrile (20ml) was added to the resulting white suspension, and the mixture stirred at room
- 10 temperature for 18 hours. The mixture was diluted with water (20ml), and extracted with ethyl acetate (2x100ml). The combined organic solutions were washed with brine, dried (MgSO₄) and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel using ethyl acetate:pentane (75:25), then dichloromethane:methanol (90:10) as
- 15 eluants, to afford the title compound as a pale yellow solid, 112.3mg.
- ¹Hnmr (CDCl₃, 300MHz) δ : 3.40 (s, 3H), 6.27 (d, 1H), 6.93 (s, 1H), 7.41 (d, 1H).

Preparation 21

20

4-(Aminomethyl)-1-methyl-2(1H)-pyridinone

Raney Nickel® (10mg) was added to a solution of the nitrile from preparation 20 (112.3mg, 0.84mmol) and potassium hydroxide (69.7mg, 1.24mmol) in ethanol


(15ml) and the mixture hydrogenated at 60psi for 18 hours. The mixture was filtered through Arbocel®, and washed through with ethanol. The filtrate was evaporated under reduced pressure, and the residue partitioned between water and dichloromethane, and the phases separated. The aqueous layer was
 5 evaporated under reduced pressure and the residue triturated with methanol (70ml). This organic solution was concentrated under reduced pressure and the residue purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (90:10:1) as eluant to give the title compound as a crystalline solid, 66.3mg.

10 ¹Hnmr (CDCl₃, 300MHz) δ: 3.50 (s, 3H), 3.69 (s, 2H), 6.11 (d, 1H), 6.49 (s, 1H), 7.21 (d, 1H).

LRMS : m/z 277.3 [2MH⁺]

15 Preparation 22

1-Acetyl-4-piperidinamine



0.88 Ammonia (85ml) was added to a solution of N-acetylperldone (15g, 106mmol) in methanol (120ml), followed by palladium hydroxide (2g) and the
 20 mixture hydrogenated at room temperature and 60 psi for 18 hours. The reaction mixture was filtered through Arbocel®, the filtrate concentrated under reduced pressure and the residue azeotroped with toluene to give a yellow oil. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (96:3.5:0.5 to
 25 84:14:2) to afford the title compound as a clear oil, 9.6g.


¹Hnmr (CDCl₃, 400MHz) δ: 1.20 (m, 2H), 1.80 (m, 2H), 2.04 (s, 3H), 2.66 (m, 1H), 2.88 (m, 1H), 3.03 (m, 1H), 3.75 (m, 1H), 4.42 (m, 1H).

LRMS : m/z (ES⁺) 165 [MNa⁺]

Preparation 23

30 tert-Butyl 3-[(acetylamino)methyl]-1-azetidinecarboxylate

155




Triethylamine (0.76ml, 5.45mmol) and acetic anhydride (0.43ml, 4.56mmol) were added to an ice-cooled solution of *tert*-butyl 3-(aminomethyl)-1-azetidinecarboxylate (J. Med. Chem. 2001; 44(1); 94) (850mg, 4.56mmol) in dichloromethane (50ml), and the reaction stirred for 30 minutes. The solution was washed with water (50ml), brine (50ml), dried (MgSO₄) and evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel using ethyl acetate as eluant to afford the title compound as a yellow oil.

- 10 ¹Hnmr (CDCl₃, 300MHz) δ : 1.43 (s, 9H), 2.00 (s, 3H), 2.72 (m, 1H), 3.44 (m, 2H), 3.61 (m, 2H), 4.00 (m, 2H), 5.74 (bs, 1H).
LRMS : m/z (ES⁻) 227 (M-H)⁻

15 Preparation 24

tert-Butyl 3-[[acetyl(methyl)amino]methyl]-1-azetidinecarboxylate



- Sodium hydride (154.7mg, 60% dispersion in mineral oil, 3.87mmol) was added to a solution of the acetamide from preparation 23 (740mg, 3.24mmol) in tetrahydrofuran (20ml), and the mixture stirred for 45 minutes. Methyl iodide
- 20


156

(0.40ml, 6.42mmol) was added and the reaction stirred at room temperature for 18 hours. Water (10ml) was then carefully added, and the tetrahydrofuran removed under reduced pressure. The aqueous solution was extracted with dichloromethane (2x30ml), the combined organic solutions washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give the title compound as a yellow oil.

¹Hnmr (CDCl₃, 300MHz) δ (mixture of rotamers in 8:3 ratio): 1.43 (s, 9H), 2.08, 2.16 (s, 3H), 2.81 (m, 1H), 2.90, 3.02 (s, 3H), 3.50-3.75 (m, 4H), 3.96-4.10 (m, 2H).
LRMS : m/z (ES⁺) 243.2 [MH⁺]

Preparation 25


N-(3-Azetidinylmethyl)-N-methylacetamide



Trifluoroacetic acid (3ml) was added to an ice-cooled solution of the protected amine from preparation 24 (720mg, 2.97mmol) in dichloromethane (20ml), and the reaction stirred for 3 hours. The solution was diluted with toluene (30ml), then concentrated under reduced pressure. The residue was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (90:10:1 to 80:20:3), and the product triturated with a solution of ether:ethyl acetate (50:50, 3x30ml), to give the title compound as a foam.


¹Hnmr (CD₃OD, 300MHz) δ: 2.10 (s, 3H), 3.08 (s, 3H), 3.21 (m, 1H), 3.64 (d, 2H), 3.95 (m, 2H), 4.09 (m, 2H).
LRMS : m/z (TSP⁺) 285.3 [MH⁺]

157

Preparation 26**4-Methyl-4-piperidinol**

A mixture of 1-benzyl-4-methyl-4-piperidinol (Tet.Lett. 37; 8;1996; 1297) (1.58g, 7.7mmol) and palladium hydroxide (500mg) in ethanol (50ml) was hydrogenated at 50psi and 50°C for 18 hours. The cooled mixture was filtered through Arbocel®, and the filtrate evaporated under reduced pressure to give the title compound as a solid, 880mg.


¹Hnmr (CDCl₃, 400MHz) δ: 1.22 (s, 3H), 1.58 (m, 4H), 2.82 (m, 2H), 2.98 (m, 2H).

Preparation 27**tert-Butyl 3-(4-hydroxy-4-methyl-1-piperidinyl)-1-azetidinecarboxylate**

A mixture of the piperidine from preparation 26 (220mg, 1.91mmol), *tert*-butyl 3-iodo-1-azetidinecarboxylate (600mg, 2.0mmol) (EP 992493) and potassium carbonate (276mg, 2.0mmol) in 1-methyl-2-pyrrolidinone (10ml) was stirred at 80°C for 48 hours. The mixture was partitioned between water and ethyl acetate, and the layers separated. The organic phase was washed with water, then brine, dried (Na₂SO₄), and evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (100:0:0 to 90:10:0.5) to afford the title compound, 162mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.20 (s, 3H), 1.38 (s, 9H), 1.60 (m, 5H), 2.20 (t, 2H), 2.42 (m, 2H), 3.05 (m, 1H), 3.88 (m, 2H), 3.90 (t, 2H).

158


LRMS : m/z (TSP⁺) 271.1 [MH⁺]**Preparation 28****5** 1-(3-Azetidinyl)-4-methyl-4-piperidinol trifluoroacetate

A mixture of the protected azetidine from preparation 27 (160mg, 0.6mmol) in trifluoroacetic acid (1ml) and dichloromethane (1ml), was stirred at room temperature for 2 hours. The solution was concentrated under reduced pressure and azeotroped with toluene, to afford the title compound as a yellow gum, 155mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.25 (s, 3H), 1.76 (d, 2H), 1.90 (m, 2H), 3.16 (t, 2H), 3.25 (m, 3H), 4.35 (m, 4H), 4.50 (m, 2H).

LRMS : m/z (ES⁺) 172 [MH⁺]

15

Preparation 291-Benzhydryl-3-(2-methoxyphenyl)-3-azetidinol

20 2-Methoxyphenylmagnesium bromide (5.9ml, 1M solution in tetrahydrofuran, 5.9mmol) was added to a cooled (-78°C) solution of 1-benzhydryl-azetidin-3-one (WO 9412181) (1g, 4.2mmol) in tetrahydrofuran (20ml), and the reaction stirred at -78°C for 15 minutes, then allowed to warm to room temperature over 30 minutes. The mixture was partitioned between water (100ml) and ethyl acetate (100ml), the layers separated, and the aqueous phase extracted with

25

ethyl acetate (100ml). The combined organic solutions were washed with brine (100ml), dried (MgSO_4) and evaporated under reduced pressure. The residual yellow oil was purified by column chromatography on silica gel using pentane:ethyl acetate (50:50) as eluant to afford the title compound as a white foam.


$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 3.58 (m, 4H), 3.95 (s, 3H), 4.42 (s, 1H), 6.94 (d, 1H), 7.00 (dd, 1H), 7.20 (m, 2H), 7.28 (m, 6H), 7.46 (m, 4H).

LRMS : m/z (TSP^+) 346.1 [MH^+]

10

Preparation 30

3-(2-Methoxyphenyl)-3-azetidinol hydrochloride




A mixture of palladium hydroxide (1g, 7.14mmol), and the azetidine from preparation 29 (950mg, 2.75mmol) in methanol (100ml) was hydrogenated at 50°C and 50 psi for 18 hours. The cooled reaction mixture was filtered through Arbocel®, and 1N ethereal hydrochloric acid was added to the filtrate. The filtrate was evaporated under reduced pressure, azeotroped with dichloromethane, and the product triturated with diethyl ether, to afford the title compound as a white solid.


$^1\text{Hnmr}$ (CD_3OD , 400MHz) δ : 3.95 (s, 3H), 4.15 (d, 2H), 4.62 (d, 2H), 6.99 (dd, 1H), 7.07 (d, 1H), 7.30 (d, 1H), 7.38 (dd, 1H).

LRMS: m/z (TSP^+) 180.1 [MH^+]

25

Preparation 31**3-Tetrahydro-2H-pyran-4-yl-azetidine p-toluenesulfonate**

- Sodium (390mg, 16.6mmol) was added to a solution of naphthalene (2.61g, 20.4mmol) in 1,2-dimethoxyethane (25ml), and the solution stirred at room temperature for 4 hours. This solution was then added dropwise to a cooled (-70°C) solution of 4-methylphenyl 3-tetrahydro-2H-pyran-4-yl-1-azetidinesulfonate (EP 962457) (1g, 3.39mmol) in 1,2-dimethoxyethane (25ml), and once the addition was complete, the mixture was stirred at -70°C for 20 minutes. The reaction was allowed to warm to room temperature, the reaction quenched by the addition of water and concentrated under reduced pressure. The residual gum was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (80:20:3) as eluant to give the title compound as an off-white gum, 690mg.
- ¹Hnmr (CD₃OD, 400MHz) δ: 1.18 (m, 2H), 1.58 (m, 2H), 1.81 (m, 1H), 2.38 (s, 3H), 2.68 (m, 1H), 3.40 (t, 2H), 3.90 (m, 4H), 4.04 (t, 2H), 7.22 (d, 2H), 7.55, 7.72 (d, 2H).

Preparation 32**N-(4-Phenyl-4-piperidiny)acetamide**

- A mixture of *N*-(1-Benzyl-4-phenyl-4-piperidiny)acetamide (Bioorg. Med. Chem. Lett. 1996; 6(19); 2307) (49g, 158mmol), and palladium hydroxide (5g) in methanol (600ml) was hydrogenated at 50 psi and room temperature for 18

161


hours. The mixture was filtered, the filtrate concentrated under reduced pressure and the residue azeotropered with dichloromethane to give the title compound as a foam.

¹Hnmr (CDCl₃, 300MHz) δ: 2.00 (m, 5H), 2.38 (m, 2H), 2.97 (m, 4H), 7.19-7.40

5 (m, 5H).

Preparation 33

N-[1-(1-Benzhydryl-3-azetidiny)-4-phenyl-4-piperidiny]acetamide




10

A mixture of 1-(diphenylmethyl)-3-azetidinyldimethanesulfonate (8g, 25.2mmol), the amine from preparation 32 (7.1g, 27.7mmol) and triethylamine (4.6ml, 32.8mmol) in acetonitrile (80ml) was heated under reflux for 18 hours. The cooled mixture was concentrated under reduced pressure, the residue partitioned between sodium bicarbonate solution and ethyl acetate, and the layers separated. The organic phase was washed with water, then brine, dried (MgSO₄) and evaporated under reduced pressure. The residual foam was purified by column chromatography on silica gel using an elution gradient (hexane:ethyl acetate:methanol 80:20:0 to 0:100:0 to 0:93:7) to afford the title compound as a white solid, 2.31g.

20


¹Hnmr (CDCl₃, 300MHz) δ: 2.00 (s, 3H), 2.01-2.20 (m, 4H), 2.37 (m, 2H), 2.62 (m, 2H), 2.84-3.06 (m, 3H), 3.41 (t, 2H), 4.41 (s, 1H), 5.44 (s, 1H), 7.12-7.54 (m, 15H).

Preparation 34**N-[1-(3-Azetidinyl)-4-phenyl-4-piperidinyl]acetamide dihydrochloride**

5 α -Chloroethylchloroformate (630 μ l, 8.63mmol) was added dropwise to an ice-cooled solution of the azetidine from preparation 33 (2.3g, 5.23mmol) in dichloromethane (25ml), and the solution stirred at room temperature for 18 hours. The mixture was concentrated under reduced pressure the residue suspended in methanol (37ml), potassium carbonate (2.2g, 15.7mmol) added, and the mixture heated under reflux for an hour. The cooled mixture was
10 filtered, the filtrate acidified to pH 3 using ethereal hydrochloric acid, then re-filtered. The filtrate was concentrated under reduced pressure, the residual gum triturated with diethyl ether, to give the title compound as a pale brown solid, 1.5g.

¹Hnmr (DMSO-d₆, 300MHz) δ : 1.92 (s, 2H), 2.30 (m, 1H), 2.48 (s, 3H), 2.63 (m, 1H), 3.10 (m, 1H), 4.08 (m, 7H), 4.50 (m, 2H), 7.12-7.65 (m, 5H), 8.30 (s, 1H),
15 9.15 (bs, 1H), 10.10 (bs, 1H).

LRMS : m/z (TSP⁺) 274.3 [MH⁺]

20 Preparation 35**Ethyl 4-cyanotetrahydro-2H-pyran-4-carboxylate**


Ethyl cyanoacetate (22.6g, 0.2mol) and bis(2-chloroethyl)ether (14g, 0.1mol) were added to a suspension of potassium carbonate (70g, 0.5mol) in N,N-

dimethylformamide (150ml), and the mixture stirred at 100°C for 72 hours. The cooled mixture was partitioned between water and diethyl ether (500ml), the layers separated and the aqueous solution extracted with diethyl ether (3x200ml). The combined organic solutions were washed with 2N hydrochloric acid (3x100ml), brine (2x100ml), dried (MgSO₄) and evaporated under reduced pressure to give a yellow oil. This crude product was purified by column chromatography on silica gel using ethyl acetate:cyclohexane (50:50) as eluant, to afford the title compound, 9.3g.

¹Hnmr (CDCl₃, 400MHz) δ: 1.34 (t, 3H), 2.00 (m, 2H), 2.14 (m, 2H), 3.74 (m, 2H), 3.98 (m, 2H), 4.26 (q, 2H).

Preparation 36

4-([[(4-Methylphenyl)sulfonyl]amino)methyl]tetrahydro-2H-pyran-4-yl)methyl 4-methylbenzenesulfonate



15

A solution of the ester from preparation 35 (9.3g, 51mmol) in tetrahydrofuran (50ml) was added dropwise to a suspension of lithium aluminium hydride (10g, 263mmol) in tetrahydrofuran (200ml), and once addition was complete, the mixture was heated under reflux for an hour. The mixture was cooled in ice, and water (15ml) in tetrahydrofuran (50ml) was added dropwise. Additional tetrahydrofuran (200ml) was added, the mixture poured into a suspension of MgSO₄ (200g) in tetrahydrofuran (300ml), and the resulting slurry filtered. The solids were washed well with tetrahydrofuran, and the combined filtrates evaporated under reduced pressure. The residual gum was dissolved in 1,2-dimethoxyethane (200ml), and triethylamine (20ml), p-toluenesulfonyl chloride (29g, 153mmol) and pyridine (21ml) added, and the reaction heated under reflux for 18 hours. The cooled mixture was partitioned between dichloromethane (300ml) and 2N hydrochloric acid (300ml), the layers

20

25

separated, and the aqueous phase extracted with dichloromethane (5x300ml). The combined organic extracts were washed with brine, dried (MgSO_4) and the mixture filtered through a plug of silica gel. The filtrate was discarded, the silica washed well with ethyl acetate, and this filtrate concentrated under reduced pressure. The residue was triturated with diethyl ether, to afford the title compound as a beige solid, 6.0g.


$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.43 (m, 1H), 1.55 (m, 3H), 2.46 (2xs, 6H), 3.00 (d, 2H), 3.60 (m, 4H), 3.88 (s, 2H), 4.88 (t, 1H), 7.37 (2xd, 4H), 7.77 (2xd, 4H).

LRMS : m/z (ES^+) 454 [MH^+]

10

Preparation 37

2-[(4-Methylphenyl)sulfonyl]-7-oxa-2-azaspiro[3.5]nonane




15 Potassium *tert*-butoxide (2.0g, 18mmol) was added to a suspension of the compound from preparation 36 (5.5g, 12mmol) in 1,2-dimethoxyethane (300ml), and the reaction stirred at 100°C for 30 minutes. The cooled mixture was concentrated under reduced pressure and the residue partitioned between dichloromethane and 2N sodium hydroxide solution, and the layers separated.

20 The aqueous phase was extracted with further dichloromethane (2x250ml), and the combined organic solutions washed with brine, dried (MgSO_4) and evaporated under reduced pressure to give the title compound as a white solid, 3.3g.

$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.57 (m, 4H), 2.48 (s, 3H), 3.51 (m, 8H), 7.39 (d, 2H),

25 7.75 (d, 2H).

LRMS : m/z (ES^+) 282 [MH^+]


Preparation 38**7-Oxa-2-azaspiro[3.5]nonane p-toluenesulphonate**

Freshly cut sodium (2.3g, 100mmol) was added to a solution of naphthalene
 5 (15.4g, 120mmol) in 1,2-dimethoxyethane (100ml), and the mixture stirred at room temperature for 3 hours.

The compound from preparation 37 (6.0g, 21mmol) was dissolved in 1,2-dimethoxyethane (100ml), and the solution cooled to -70°C . The prepared solution of sodium naphthalenide was then added, the solution stirred for 10
 10 minutes, and then quenched by the addition of water (5ml). The solution was allowed to warm to room temperature, potassium carbonate (200g) and additional 1,2-dimethoxyethane (500ml) added, and the mixture filtered and evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia
 15 (80:20:5) as eluant, and the product azeotroped with toluene (5x100ml) to give a gum. This was triturated with ether to afford the title compound as a white solid, 4.1g.

$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.65 (m, 4H), 2.38 (s, 3H), 3.44 (m, 4H), 3.52 (s, 4H), 7.22 (d, 2H), 7.56 (d, 2H).

20

Preparation 39**tert-Butyl (3R)-3-(acetilamino)-1-pyrrolidinecarboxylate**

25

Di-tert-butyl dicarbonate (1.92g, 8.8mmol) was added to a solution of *N*-[(3*R*)-pyrrolidinyl]acetamide (1g, 8mmol) in dichloromethane (30ml), and the solution

166


cooled in ice. Hünig's base (1.5ml, 8.8mmol) was added dropwise, and once addition was complete the reaction was stirred for 2 hours. The mixture was washed with sodium bicarbonate solution (3x), then brine (3x), dried (MgSO₄) and evaporated under reduced pressure to give the title compound, 1.75g.

- 5 ¹Hnmr (CDCl₃, 300MHz) δ: 1.46 (s, 9H), 1.96 (s, 3H), 2.16 (m, 2H), 3.18 (m, 1H), 3.40 (m, 2H), 3.60 (m, 1H), 4.42 (m, 1H), 5.70 (bs, 1H).

Preparation 40

10

tert-Butyl (3S)-3-(acetlamino)-1-pyrrolidinecarboxylate




The title compound was obtained as a colourless gum, from *N*-[(3*S*)-pyrrolidinyl]acetamide, following the procedure described in preparation.

- 15 ¹Hnmr (CDCl₃, 300MHz) δ: 1.46 (s, 9H), 1.96 (s, 3H), 2.10 (m, 2H), 3.16 (m, 1H), 3.38 (m, 2H), 3.58 (m, 1H), 4.42 (m, 1H), 5.52 (bs, 1H).

Preparation 41

20

tert-Butyl (3*R*)-3-[acetyl(methyl)amino]-1-pyrrolidinecarboxylate



- A solution of the pyrrolidine from preparation 39 (1.75g, 7.7mmol) in *N,N*-dimethylformamide (5ml), was added dropwise to a mixture of sodium hydride (470mg, 11.7mmol) in *N,N*-dimethylformamide (10ml), and the solution stirred
- 25

167

for 30 minutes. Iodomethane (0.8ml, 8.4mmol) was added, and the reaction stirred at room temperature for 2 hours. Aqueous ammonium chloride solution was added and the mixture extracted with ethyl acetate. The combined organic solutions were washed with water, brine, then dried (MgSO_4) and evaporated under reduced pressure to give the title compound as a yellow gum, 725mg.


$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.44 (s, 9H), 1.90-2.16 (m, 5H), 2.95 (s, 3H), 3.12 (m, 1H), 3.30 (m, 1H), 3.50 (m, 2H), 4.38, 5.20 (2xm, 1H).

LRMS : m/z (TSP^+) 243.2 [MH^+]

10

Preparation 42


tert-Butyl (3*S*)-3-[acetyl(methyl)amino]-1-pyrrolidinecarboxylate




15 The title compound was obtained as a yellow gum in 44% yield, from the pyrrolidine from preparation 40, following the procedure described in preparation 41.

$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.42 (s, 9H), 1.82-2.15 (m, 5H), 2.84 (s, 3H), 3.14 (m, 1H), 3.25 (m, 1H), 3.54 (m, 2H), 4.38, 5.18 (2xm, 1H).


20 LRMS : m/z (TSP^+) 243.2 [MH^+]

Preparation 43**N-Methyl-N-[(3R)-pyrrolidinyl]acetamide trifluoroacetate**

- 5 A mixture of the protected pyrrolidine from preparation 41 (720mg, 2.96mmol) and trifluoroacetic acid (4ml) in dichloromethane (4ml) was stirred at room temperature for 1 hour. The mixture was concentrated under reduced pressure, the residue azeotroped with toluene (3x), dichloromethane (3x) and diethyl ether (3x) to afford the title compound as a gum.
- 10 ¹Hnmr (CD₃OD, 400MHz) δ: 2.10 (s, 3H), 2.16 (m, 1H), 2.38 (m, 1H), 3.06 (s, 3H), 3.20 (m, 1H), 3.40 (m, 2H), 3.60 (m, 1H), 4.50 (m, 1H).
LRMS : m/z (TSP⁺) 285.2 [MH⁺]

15 Preparation 44**N-Methyl-N-[(3S)-pyrrolidinyl]acetamide trifluoroacetate**

- The title compound was obtained as a gum, from the protected pyrrolidine from preparation 42, following the procedure described in preparation 43.
- 20 ¹Hnmr (CD₃OD, 400MHz) δ: 2.08 (s, 3H), 2.18 (m, 1H), 2.38 (m, 1H), 3.06 (s, 3H), 3.20 (m, 1H), 3.40 (m, 2H), 3.60 (m, 1H), 4.50 (m, 1H).
LRMS : m/z (TSP⁺) 285.2 [MH⁺]

Preparation 45**tert-Butyl (3R)-3-methoxy-1-pyrrolidinecarboxylate**


Sodium hydride (2.2g, 80% dispersion in mineral oil, 73.2mmol) was added to
5 an ice-cooled solution of (3R)-N-*tert*-butoxycarbonylpyrrolidin-3-ol (J. Med.
Chem. 41; 25; 1998; 4983) (12.5g, 66.7mmol) in tetrahydrofuran (330ml), and
the solution stirred at room temperature for an hour. Methyl iodide (14.5g,
100mmol) was then added and the reaction stirred for 18 hours. Water (100ml)
10 was added and the mixture concentrated under reduced pressure to remove the
organic solvents. The aqueous was extracted with ethyl acetate, the combined
organic solutions dried (MgSO₄) and evaporated under reduced pressure to
give the title compound as an oil, 12.48g.

¹Hnmr (CDCl₃, 400MHz) δ: 1.42 (s, 9H), 1.86-2.01 (m, 2H), 3.32 (s, 3H), 3.40
(m, 4H), 3.93 (m, 1H).

15 Microanalysis found: C, 59.71; H, 9.63; N, 6.71. C₁₀H₁₉NO₃ requires C, 59.68;
H, 9.52; N, 6.96%.

Preparation 46

20

tert-Butyl (3S)-3-methoxy-1-pyrrolidinecarboxylate

The title compound was prepared from (3S)-N-*tert*-butoxycarbonylpyrrolidin-3-ol
(US 6180627), following the procedure described in preparation 45.


170

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.42 (s, 9H), 1.84-2.00 (m, 2H), 3.32 (s, 3H), 3.40 (m, 4H), 3.92 (m, 1H).

Microanalysis found: C, 59.72; H, 9.62; N, 6.63. $\text{C}_{10}\text{H}_{19}\text{NO}_3$ requires C, 59.68; H, 9.52; N, 6.96%.

5 Preparation 47

(3R)-3-Methoxypyrrolidine trifluoroacetate




Hydrogen chloride was bubbled through a solution of the protected amine from preparation 45 (24.8g, 123mmol) in diethyl ether (615ml), until saturated, and the solution then stirred for 1 hour at room temperature. The reaction was concentrated under reduced pressure, the residue resuspended in diethyl ether, the solution stirred for 2 hours the ether decanted off, and the residue evaporated under reduced pressure. The product was dissolved in ethanol, trifluoroacetic acid (200ml) added, and the solution evaporated under reduced pressure to afford the title compound.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 2.00 (m, 1H), 2.18 (m, 1H), 3.24-3.50 (m, 7H), 4.05 (m, 1H), 8.80 (bs, 1H), 9.37 (bs, 1H).

Microanalysis found: C, 33.76; H, 5.35; N, 5.54.
 $\text{C}_5\text{H}_{11}\text{NO} \cdot 1.25\text{CF}_3\text{CO}_2\text{H} \cdot 1.25\text{H}_2\text{O}$ requires C, 33.84; H, 5.59; N, 5.26%.

Preparation 48


(3S)-3-Methoxypyrrolidine trifluoroacetate



The title compound was obtained, from the protected amine from preparation 46, following the procedure described in preparation 47.

171

¹Hnmr (CDCl₃, 400MHz) δ: 2.00 (m, 1H), 2.18 (m, 1H), 3.25-3.48 (m, 7H), 4.06 (m, 1H), 8.75 (bs, 1H), 9.24 (bs, 1H).


Preparation 49**N-[(3S)-1-Benzylpyrrolidinyl]-4-chlorobutanamide**

5 4-Chlorobutyl chloride (0.31ml, 3.1mmol) was added to a mixture of (3S)-1-benzyl-3-pyrrolidinamine (500mg, 2.8mmol) in tetrahydrofuran (30ml), and the reaction was stirred at room temperature for 2 hours. The mixture was washed with water, then brine, dried (MgSO₄) and evaporated under reduced pressure to give the title compound as a yellow gum, 823mg.

10 ¹Hnmr (CDCl₃, 400MHz) δ: 1.62 (m, 1H), 2.06 (m, 2H), 2.24 (m, 4H), 2.56 (m, 1H), 2.62 (m, 1H), 2.94 (m, 1H), 3.58 (m, 4H), 4.44 (m, 1H), 6.05 (bs, 1H), 7.20-7.35 (m, 5H).

LRMS : m/z (ES⁺) 281, 283 [MH⁺]

15

Preparation 50**N-[(3R)-1-Benzylpyrrolidinyl]-4-chlorobutanamide**

20


The title compound was obtained as a gum from (3R)-1-benzyl-3-pyrrolidinamine following the procedure described in preparation 49.

¹Hnmr (CDCl₃, 400MHz) δ: 1.62 (m, 1H), 2.10 (m, 2H), 2.232 (m, 4H), 2.58 (m, 1H), 2.64 (m, 1H), 2.96 (m, 1H), 3.62 (m, 4H), 4.50 (m, 1H), 6.02 (bs, 1H), 7.15-
25 7.35 (m, 5H).

173

LRMS : m/z (TSP⁺) 281.1, 283.1 [MH⁺]**Preparation 51**

5

1-[(3S)-1-Benzylpyrrolidinyl]-2-pyrrolidone

The pyrrolidine from preparation 49 (825mg, 3mmol) was added to a mixture of sodium hydride (176mg, 60% dispersion in mineral oil, 4.4mmol) in 1-methyl-2-pyrrolidine (10ml), and the reaction stirred at room temperature for 18 hours.


10 Aqueous ammonium chloride was added to quench the reaction, then the mixture extracted with ethyl acetate. The combined organic solutions were washed with water (3x), brine (3x), dried (MgSO₄) and evaporated under reduced pressure. The residual gum was purified by column chromatography on silica gel using dichloromethane:methanol (95:5) as eluant, to afford the title
15 compound as a yellow oil, 250mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.70 (m, 1H), 1.94 (m, 2H), 2.10 (m, 1H), 2.28 (m, 3H), 2.44 (m, 1H), 2.56 (m, 1H), 2.84 (m, 1H), 3.38-3.68 (m, 4H), 4.76 (m, 1H), 7.20-7.35 (m, 5H).

LRMS : m/z (TSP⁺) 245.1 [MH⁺]

20

174

Preparation 521-[(3R)-1-Benzylpyrrolidinyl]-2-pyrrolidone


5 The title compound was obtained as a yellow gum in 18% yield, from the compound from preparation 50, following the procedure described in preparation 51.

¹Hnmr (CDCl₃, 400MHz) δ: 1.74 (bs, 1H), 1.98 (m, 2H), 2.08 (m, 1H), 2.16 (m, 3H), 2.50 (m, 1H), 2.62 (m, 1H), 2.90 (m, 1H), 3.40-3.75 (m, 4H), 4.78 (s, 1H),
10 7.20-7.35 (m, 5H).

LRMS : m/z (TSP⁺) 245.2 [MH⁺]

Preparation 53

15

1-[(3S)-Pyrrolidinyl]-2-pyrrolidone


A mixture of the protected pyrrolidine from preparation 51 (246mg, 1mmol) and palladium hydroxide (150mg) in ethanol (10ml), was hydrogenated at 60 psi and 60°C for 18 hours. The cooled mixture was filtered through Arbocel®, washing
20 through with ethanol, and the filtrate evaporated under reduced pressure, to give the title compound as a yellow gum, 156mg.

175

¹Hnmr (CDCl₃, 400MHz) δ: 1.86 (m, 1H), 1.97-2.17 (m, 3H), 2.38 (t, 2H), 3.02 (m, 2H), 3.20 (m, 2H), 3.41 (m, 2H), 4.60 (m, 1H) 4.70-4.92 (bs, 1H).

Preparation 541-[(3R)-pyrrolidin-3-yl]-2-pyrrolidone

5



The title compound was obtained as a gum, from the protected pyrrolidine from preparation 52, following the procedure described in preparation 53.


¹Hnmr (CDCl₃, 400MHz) δ: 1.78 (m, 1H), 2.05 (m, 3H), 2.38 (t, 2H), 2.8-3.05 (m, 3H), 3.16 (m, 2H), 3.42 (t, 2H), 4.60 (m, 1H).

10

LRMS : m/z (TSP⁺) 155.2 [MH⁺]

Preparation 55

15

1-Benzyl-4-ethyl-4-piperidinol

Ethylmagnesium bromide (18ml, 3M solution in diethyl ether, 54mmol) was added dropwise over 30 minutes to a cooled (-78°C) solution of 1-benzyl-4-piperidinone (5g, 26.4mmol) in diethyl ether (50ml). Once addition was complete, the mixture was allowed to warm to room temperature and then stirred for 18 hours.

20

The residual gum was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (97:3:0.5 to 90:10:1) to afford the title compound as an oil, 848mg.


25

176

¹Hnmr (CDCl₃, 400MHz) δ: 0.94 (t, 3H), 1.52 m, 4H), 1.63 (m, 2H), 2.35 (m, 2H), 2.62 (m, 2H), 3.55 (s, 2H), 7.32 (m, 5H).


Preparation 56

5

4-Ethyl-4-piperidinol

A mixture of the amine from preparation 55 (848mg, 3.87mmol) and palladium hydroxide (300mg) in ethanol (30ml) was hydrogenated at 60 psi and room temperature for 18 hours. The mixture was filtered through Arbocel®, and the filtrate evaporated under reduced pressure to give the title compound as an oil, 380mg.

¹Hnmr (CDCl₃, 400MHz) δ: 0.93 (t, 3H), 1.45-1.66 (m, 6H), 2.88-3.04 (m, 4H).

15 Preparation 57**tert-Butyl 4-hydroxy-4-(trifluoromethyl)-1-piperidinecarboxylate**

Tetrabutylammonium fluoride (50mg, 1M solution in tetrahydrofuran) was added to an ice-cooled solution of (trifluoromethyl)trimethylsilane (2.1g, 15mmol) and *tert*-butyl 4-oxo-1-piperidinecarboxylate (2g, 10mmol) in tetrahydrofuran (20ml), and the reaction stirred at room temperature for 18 hours. The mixture was concentrated under reduced pressure, the residue suspended in ethyl acetate, hydrochloric acid (20ml, 1N) was added, the mixture stirred for an hour, and then neutralised using sodium bicarbonate. The solution was washed with water, then brine, dried (MgSO₄) and evaporated under reduced pressure to give a gum. This was purified by column chromatography on silica gel using an


elution gradient of dichloromethane:methanol (100:0 to 98:2) to afford the title compound, 1.94g.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.42 (s, 9H), 1.68 (d, 2H), 1.76 (m, 2H), 3.02 (m, 2H), 4.02 (m, 2H).

5 LRMS : m/z (TSP $^+$) 270.2 [MH^+]

Preparation 58

4-Trifluoromethylpiperidinol trifluoroacetate




10 A mixture of the piperidine from preparation 57 (950mg, 3.6mmol) and trifluoroacetic acid (5ml) in dichloromethane (5ml) was stirred at room temperature for 90 minutes. The mixture was concentrated under reduced pressure, and the residue azeotroped with toluene and dichloromethane. The product was triturated with diethyl ether to afford the title compound as a yellow solid, 866mg.

15 $^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.94 (m, 4H), 3.18-3.35 (m, 4H).
LRMS : m/z (TSP $^+$) 170.0 [MH^+]

Preparation 59

tert-Butyl 4-[acetyl(methyl)amino]-1-piperidinecarboxylate



25 Triethylamine (18ml, 132mmol), followed by acetic anhydride (8.8ml, 93mmol) were added to a solution of *tert*-butyl 4-[(methyl)amino]-1-piperidinecarboxylate (WO 9639385) (9g, 89mmol) in dichloromethane (300ml), and the reaction stirred at room temperature for 1 hour. The solution was diluted with dichloromethane (200ml), washed with 2N citric acid (2x200ml), brine, dried (MgSO_4) and evaporated under reduced pressure. The crude product was


178

purified by column chromatography on silica gel using dichloromethane:methanol (90:10), and the product azeotrope with dichloromethane to afford the title compound as a yellow oil, 20.5g.

¹Hnmr (CDCl₃, 300MHz) δ: (mixture of rotamers) 1.40-1.74 (m, 13H), 2.09, 2.13 (2xs, 3H), 2.67-2.83 (m, 5H), 3.62, 4.60 (2xm, 1H), 4.12-4.25 (m, 2H).

Preparation 60

N-methyl-N-(4-piperidiny)acetamide hydrochloride




- 10 A solution of the protected amine from preparation 59 (20g, 78mmol) in dichloromethane (200ml) was saturated with hydrogen chloride, and the reaction then stirred at room temperature for 1 hour. The mixture was concentrated under reduced pressure, the residue azeotrope with dichloromethane (3x300ml), and the resulting solid triturated with diethyl ether.
- 15 The solid was filtered off, and dried under vacuum, to give the title compound as a white solid, 16.3g.

¹Hnmr (DMSO-d₆, 300MHz) δ: (mixture of rotamers) 1.55 (m, 1H), 1.70 (m, 1H), 1.96-2.15 (m, 5H), 2.60, 2.78 (2xs, 3H), 2.82-3.02 (m, 2H), 3.22 (m, 2H), 3.98, 4.47 (2xm, 1H), 9.23-9.42 (bs, 2H).

20

Preparation 61

1-Benzyl-4-piperidiny methylcarbamate



- 25 Methyl isocyanate (8.94g, 157mmol) was added to a solution of 1-benzyl-4-piperidinol (10g, 52.3mmol) in chloroform (80ml), and the reaction stirred under reflux for 16 hours. The cooled mixture was concentrated under reduced

179


pressure, the residual solid was triturated from 40-60 petroleum ether, and the product filtered and dried to afford the title compound as a white solid, 11g.

m.p.- 106-108°C

Microanalysis found: C, 67.86; 8.14; N, 11.25. $C_{14}H_{20}N_2O_2$ requires C, 67.71; H, 8.12; N, 11.28%.

Preparation 62

4-Piperidinyl methylcarbamate



10 A mixture of the piperidine from preparation 61 (8.2g, 33.0mmol) and 10% palladium on charcoal (1g) in ethanol (200ml) was hydrogenated at 50°C and 50 psi for 18 hours. The cooled reaction was filtered through Hyflo®, and the filtrate evaporated under reduced pressure to afford the title compound as a white solid, 5.5g.


15 $^1\text{Hnmr}$ (CDCl_3 , 90MHz) δ : 1.0-2.10 (m, 4H), 2.30-3.30 (m, 7H), 4.40-5.20 (m, 2H).

Microanalysis found: C, 53.29; H, 8.87; N, 17.78. $C_7H_{14}N_2O_2$ requires C, 53.14; H, 8.91; N, 17.71%.

20

Preparation 63

tert-Butyl 4-(4-hydroxypiperidin-1-yl)-1-piperidinecarboxylate



Acetic acid (3.25ml, 41.3mmol) followed by *tert*-butyl 4-oxo-1-piperidinecarboxylate (1g, 5.0mmol) and sodium triacetoxyborohydride (2.11g, 10mmol) were added to a solution of 4-piperidinol (757mg, 7.5mmol) and


triethylamine (5.5ml, 37.5mmol) in dichloromethane (50ml), and the reaction stirred at room temperature for 18 hours. The reaction was washed with sodium bicarbonate solution, dried (MgSO_4) and evaporated under reduced pressure. The residual oil was purified by column chromatography on silica gel using an
5 elution gradient of dichloromethane:methanol:0.88 ammonia (95:5:0.5 to 93:7:0.5) to give the title compound as a clear gum, 700mg.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.40 (m, 11H), 1.58 (m, 2H), 1.78 (m, 2H), 1.95 (m, 2H), 2.24-2.48 (m, 3H), 2.68 (m, 2H), 2.82 (m, 2H), 3.70 (m, 1H), 4.16 (m, 2H).

Preparation 64

10

1-(Piperidin-4-yl)-4-piperidinol ditrifluoroacetate




Trifluoroacetic acid (0.58ml, 7.5mmol) was added dropwise to an ice-cooled solution of the amine from preparation 63 (700mg, 2.5mmol) in dichloromethane
15 (10ml), and the reaction stirred at room temperature for 18 hours. Tlc analysis showed starting material remaining, so additional trifluoroacetic acid (0.97ml, 12.5mmol) was added, and the reaction stirred for a further 2 hours. The mixture was concentrated under reduced pressure and the residue azeotroped with toluene to afford the title compound as an oil, 1.25g.

20 $^1\text{Hnmr}$ (CD_3OD , 400Hz) δ : 1.97 (m, 4H), 2.17 (m, 1H), 2.37 (m, 3H), 3.15 (m, 3H), 3.28-3.61 (m, 7H).

Preparation 65

25

4-(4-piperidinyl)morpholine hydrochloride



181

Hydrogen chloride was bubbled through an ice-cooled solution of *tert*-butyl 4-(4-morpholinyl)-1-piperidinecarboxylate (J.O.C. 1990; 55(8); 2552) (6.5g, 24mmol) in dichloromethane (100ml), and the solution then stirred at 0°C for 2 hours. The reaction was degassed under nitrogen, allowed to warm to room temperature, and evaporated under reduced pressure to afford the title compound as a white solid, 5.91g.


¹Hnmr (DMSOd₆, 400MHz) δ: 1.90 (m, 2H), 2.22 (m, 2H), 2.80 (m, 2H), 3.00 (m, 2H), 3.38 (m, 5H), 3.90 (m, 4H), 8.92 (bs, 1H), 9.20 (bs, 1H).

LRMS : m/z (ES⁺) 171.2 [MH⁺]

10

Preparation 66

3-(4-Pyridyl)-2,4-imidazolidinedione



15 A mixture of 4-aminopyridine (25g, 266mmol) and ethylisocyanoacetate (35g, 271mmol) in N,N-dimethylformamide (250ml) was heated under reflux for 90 minutes, and allowed to cool. The resulting precipitate was filtered off, and the filtrate heated under reflux for a further 5 hours. The cooled mixture was concentrated under reduced pressure, and the residue triturated with hot ethanol (500ml). The resulting solid was filtered, and recrystallised from N,N-dimethylformamide to afford the title compound as a yellow crystalline solid, 47.8g.

m.p. 232-234°C


Microanalysis found: C, 53.98; H, 3.99; N, 23.63. C₈H₇N₃O₂ requires C, 54.23;

25 H, 3.98; N, 23.63%

Preparation 67

3-(4-Piperidiny)-2,4-imidazolidinedione

182



The pyridyl compound from preparation 66 (42g, 0.24mol) was dissolved in 5N hydrochloric acid, then evaporated under reduced pressure. The solid was dissolved in water, 5% rhodium on alumina (15g) added, and the mixture
5 hydrogenated at 50°C and 750 psi. The cooled mixture was filtered, the filtrate concentrated under reduced pressure and the residue dissolved in water. The solution was basified and evaporated under reduced pressure. The resulting solid was extracted into ethyl acetate using a Soxhlet apparatus over 2 days. The organic solution was evaporated under reduced pressure and the residual
10 yellow solid recrystallised from methanol/butanol to afford the title compound as a white solid, 6.9g.


m.p. 214-217°C

Microanalysis found: C, 52.20; H, 7.21; N, 23.04. $C_7H_{12}N_3O_2$ requires C, 52.44; H, 7.15; N, 22.94%

15

Preparation 68

tert-Butyl 4-(4-pyridinyl)-1-piperidinecarboxylate



20 Dichloromethane (0.1ml) was added to a suspension of zinc (2g, 31.7mmol) in N,N-dimethylformamide (5ml), and the mixture warmed until gas evolution occurred. *tert*-Butyl 4-iodo-1-piperidinecarboxylate (EP 1078928) (4.9g, 15.8mmol), and hydroquinone (35mg, 0.32mmol) in N,N-dimethylformamide (5ml) was added, and the mixture warmed until an exotherm was evident. 4-
25 Bromopyridine (1g, 6.33mmol), tris(dibenzylideneacetone)dipalladium (0)


183

(73mg, 0.127mmol) and tri(2-furyl)phosphine (59mg, 0.25mmol) in N,N-dimethylformamide (5ml) were added, and the reaction stirred at 60°C for 30 minutes. The cooled mixture was partitioned between water (100ml) and diethyl ether (50ml), and the layers separated. The aqueous phase was extracted with
5 diethyl ether (2x50ml), the combined organic solutions washed with brine (50ml), dried (MgSO₄) and evaporated under reduced pressure, to give a brown oil. This was purified by column chromatography on silica gel using ethyl acetate:pentane (50:50) as eluant to afford the title compound as a yellow oil, 1.11g.

10 ¹Hnmr (CDCl₃, 400MHz) δ: 1.43 (s, 9H), 1.58 (m, 2H), 1.80 (m, 2H), 2.60 (m, 1H), 2.78 (m, 2H), 4.25 (m, 2H), 7.09 (d, 2H), 8.49 (d, 2H).
LRMS : m/z (TSP⁺) 263.2 [MH⁺]

15 Preparation 69

tert-Butyl 4-(3-pyridinyl)-1-piperidinecarboxylate



The title compound was obtained as an oil in 40% yield, from *tert*-butyl 4-iodo-1-piperidinecarboxylate (EP 1078928) and 3-bromopyridine, according to the
20 procedure described in preparation 68.


¹Hnmr (CDCl₃, 400MHz) δ: 1.42 (s, 9H), 1.58 (m, 2H), 1.78 (m, 2H), 2.62 (m, 1H), 2.78 (m, 2H), 4.21 (m, 2H), 7.20 (m, 1H), 7.45 (d, 1H), 8.42 (m, 2H).
LRMS : m/z (TSP⁺) 263.1 [MH⁺]

25

Preparation 70

tert-Butyl 4-(1-oxido-4-pyridinyl)-1-piperidinecarboxylate

184




Phthalic anhydride (1.41g, 9.53mmol) was added to a suspension of urea hydrogen peroxide addition compound (2.87g, 30.5mmol) in dichloromethane (10ml), and the mixture stirred at room temperature for 15 minutes. The pyridyl compound from preparation 68 (1g, 3.82mmol) in dichloromethane (10ml) was added and the reaction stirred at room temperature for 72 hours. The mixture was washed with water (100ml), and the aqueous solution was extracted with further dichloromethane (3x75ml). The combined organic solutions were washed with brine (75ml), dried (MgSO₄) and concentrated under reduced pressure. The residual oil was purified by column chromatography on silica gel using dichloromethane:methanol (95:5) as eluant to afford the title compound as an off-white oil, 978mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.46 (s, 9H), 1.58 (m, 2H), 1.82 (m, 2H), 2.66 (m, 1H), 2.80 (m, 2H), 4.28 (m, 2H), 7.14 (d, 2H), 8.18 (d, 2H).

LRMS : m/z (TSP⁺) 279.2 [MH⁺]

Preparation 71

tert-Butyl 4-(1-oxido-3-pyridinyl)-1-piperidinecarboxylate




20

The title compound was obtained as a white solid in 62% yield from the pyridyl compound from preparation 69, according to the procedure described in preparation 70.

185

¹Hnmr (CDCl₃, 400MHz) δ: 1.46 (s, 9H), 1.58 (m, 2H), 1.82 (m, 2H), 2.63 (m, 1H), 2.78 (m, 2H), 4.26 (m, 2H), 7.14 (d, 1H), 7.23 (d, 1H), 8.11 (m, 2H).

LRMS : m/z (TSP⁺) 279.1 [MH⁺]


Preparation 72**4-(4-Piperidinyl)pyridine 1-oxide hydrochloride**

5 Hydrogen chloride was bubbled through a solution of the pyridyl compound from preparation 70 (978mg, 3.52mmol) in dichloromethane (100ml) for 15 minutes. The reaction mixture was then evaporated under reduced pressure to afford the title compound as a white solid, 1.01g.

¹Hnmr (CD₃OD, 400MHz) δ: 2.00 (m, 2H), 2.18 (m, 2H), 3.18 (m, 2H), 3.28 (m,
10 1H), 3.54 (m, 2H), 8.00 (d, 2H), 8.84 (d, 2H).
LRMS : m/z (TSP⁺) 179.2 [MH⁺]

Preparation 73

15

3-(4-Piperidinyl)pyridine 1-oxide hydrochloride

The title compound was obtained as a white solid in quantitative yield, from the pyridyl compound from preparation 71, following the procedure described in preparation 72.


20 ¹Hnmr (CD₃OD, 400MHz) δ: 2.00 (m, 2H), 2.18 (m, 2H), 3.18 (m, 2H), 3.24 (m, 1H), 3.54 (m, 2H), 7.98 (dd, 1H), 8.32 (d, 1H), 8.78 (d, 1H), 8.90 (s, 1H).
LRMS : m/z (TSP⁺) 179.2 [MH⁺]

Preparation 74

25

1-(1-Oxido-2-pyridinyl)piperazine dihydrochloride

187




A mixture of 2-chloropyridine 1-oxide (1g, 6.06mmol) and piperazine hexahydrate (6g, 30.9mmol) were heated at 160-180°C for 4 hours, using a Dean and Stark apparatus. The cooled mixture was diluted with methanol (15ml) and dichloromethane (100ml), silica gel (12g) added, and the mixture evaporated under reduced pressure. This was purified by column chromatography on silica gel using dichloromethane:methanol:0.88 ammonia (94:12:2) as eluant to give a yellow oil. This was suspended in ethereal hydrochloric acid, and the mixture evaporated under reduced pressure to afford the title compound as a white powder, 0.94g.

¹Hnmr (DMSOd₆, 400MHz) δ: 3.20 (m, 4H), 3.58 (m, 4H), 7.02 (dd, 1H), 7.16 (d, 1H), 7.37 (dd, 1H), 8.18 (d, 1H), 9.42 (bs, 2H).

15 Preparation 75

tert-Butyl 4-(3-cyano-2-pyridinyl)-1-piperazinecarboxylate



A mixture of 2-chloro-3-cyanopyridine (15.2g, 0.11mol), *tert*-butyl 1-piperazinecarboxylate (25g, 0.13mol) and triethylamine (18ml, 0.13mol) in toluene (200ml), was heated under reflux for 24 hours. The cooled mixture was concentrated under reduced pressure and the residue suspended in ethyl acetate (250ml), and washed with water (3x). The organic solution was dried (MgSO₄), and concentrated under reduced pressure. The residue was triturated with pentane, filtered and dried *in vacuo*, at 60°C, to afford the title compound as a cream coloured solid, 24.2g.

188


$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.47 (s, 9H), 3.59 (m, 4H), 3.66 (m, 4H), 6.78 (dd, 1H), 7.78 (d, 1H), 8.35 (d, 1H).

LRMS : m/z -(TSP $^+$) 289.2 [MH^+]

5

Preparation 76

tert-butyl 4-[3-(aminomethyl)-2-pyridinyl]-1-piperazinecarboxylate




A mixture of the nitrile from preparation 75 (708mg, 2.45mmol) and Raney Nickel® (170mg) in ethanolic ammonia (20ml) was hydrogenated at 60 psi for 16 hours. The mixture was filtered through Arbocel®, and the filtrate evaporated under reduced pressure to give the title compound as an oil.

$^1\text{Hnmr}$ (CDCl_3 , 400MHz) δ : 1.43 (s, 9H), 3.07 (m, 4H), 3.53 (m, 4H), 3.85 (bs, 2H), 6.95 (dd, 1H), 7.63 (d, 1H), 8.19 (s, 1H).

LRMS : m/z (TSP $^+$) 292.43 [MH^+]

Preparation 77

tert-Butyl 4-[3-[(dimethylamino)methyl]-2-pyridinyl]-1-piperazinecarboxylate



20


Formaldehyde (2ml, 33% aqueous solution) and sodium triacetoxyborohydride (525.6mg, 2.48mmol) were added to a solution of the amine from preparation 76 (362.1mg, 1.24mmol) in dichloromethane (10ml), and the solution stirred at

room temperature for 45 minutes. The mixture was washed with saturated aqueous sodium bicarbonate solution, brine, dried (MgSO_4) and evaporated under reduced pressure to afford the title compound as a yellow oil, 375mg.

$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.52 (s, 9H), 2.28 (s, 6H), 3.15 (m, 4H), 3.41 (s, 2H),
5 3.59 (m, 4H), 6.97 (dd, 1H), 7.72 (d, 1H), 8.22 (d, 1H).
LRMS : m/z (TSP^+) 321.3 [MH^+]


Preparation 78

10 *tert*-Butyl 4-(3-(((methylsulfonyl)amino)methyl)-2-pyridinyl)-1-piperazinecarboxylate



Triethylamine (0.2ml, 1.43mmol), followed by methanesulfonyl chloride (0.1ml, 1.29mmol) were added to an ice-cooled solution of the amine from preparation
15 76 (362.1mg, 1.24mmol) in dichloromethane (10ml), and the solution stirred for 30 minutes. The mixture was washed with 10% aqueous citric acid solution, then brine, dried (MgSO_4) and evaporated under reduced pressure to afford the title compound as a yellow oil, 306mg.


$^1\text{Hnmr}$ (CDCl_3 , 300MHz) δ : 1.49 (s, 9H), 2.93 (s, 3H), 3.12 (s, 4H), 3.61 (m, 4H),
20 4.39 (s, 2H), 5.51 (bs, 1H), 7.06 (dd, 1H), 7.69 (d, 1H), 8.32 (s, 1H).
LRMS : m/z (TSP^+) 371.2 [MH^+]

Preparation 79***N,N*-Dimethyl[2-(1-piperazinyl)-3-pyridinyl]methanamine trihydrochloride**

Hydrogen chloride was bubbled through a solution of the protected amine from preparation 77 (375.8mg, 1.17mmol) in dichloromethane (50ml), and the solution stirred for 30 minutes. The mixture was evaporated under reduced pressure and the residue dried *in vacuo*, to afford the title compound, 371mg.

$^1\text{Hnmr}$ (CD_3OD , 300MHz) δ : 2.94 (s, 6H), 3.41-3.58 (m, 10H), 7.42 (dd, 1H), 8.17 (d, 1H), 8.52 (d, 1H).


LRMS : m/z (TSP^+) 221.2 [MH^+]

Preparation 80***N*-Methyl-*N*-{[2-(1-piperazinyl)-3-pyridinyl]methyl}methanesulfonamide dihydrochloride**


The title compound was obtained as a white solid in 97% yield from the protected amine from preparation 78, according to the procedure described in preparation 79.

$^1\text{Hnmr}$ (DMSO-d_6 , 300MHz) δ : 2.94 (s, 3H), 3.18 (m, 4H), 3.31 (m, 4H), 4.12 (s, 2H), 7.17 (dd, 1H), 7.59 (bs, 1H), 7.87 (d, 1H), 8.19 (d, 1H), 9.33 (bs, 2H).


LRMS : m/z (TSP^+) 271.2 [MH^+]

Preparation 81**tert-Butyl 4-(methylsulfonyl)-1,4-diazepane-1-carboxylate**

- Methanesulfonyl chloride (0.64ml, 8.24mmol) was added to a solution of *tert*-butyl 1,4-diazepane-1-carboxylate (1.5g, 7.49mmol) and triethylamine (1.6ml, 11mmol) in dichloromethane (20ml), and the reaction stirred at room temperature for 18 hours. The solution was washed with sodium bicarbonate solution, then brine, dried (MgSO₄) and evaporated under reduced pressure to give the title compound, 2.01g.
- ¹Hnmr (CDCl₃, 300MHz) δ: 1.45 (s, 9H), 1.95 (m, 2H), 2.83 (s, 3H), 3.38 (m, 4H), 3.52 (m, 4H).

Preparation 82**1-(Methylsulfonyl)-1,4-diazepane hydrochloride**


- Hydrogen chloride was bubbled through an ice-cooled solution of the compound from preparation 81 (2.0g, 7.2mmol) in dichloromethane (50ml), for 15 minutes. The solution was allowed to warm to room temperature and concentrated under reduced pressure. The residue was azeotroped with dichloromethane and triturated with diethyl ether to afford the title compound as a solid, 1.45g.
- ¹Hnmr (DMSO-d₆, 300MHz) δ: 2.00 (m, 2H), 2.98 (s, 3H), 3.17 (m, 4H), 3.35 (t, 2H), 3.54 (t, 2H), 9.35 (bs, 2H).
- LRMS : m/z (TSP⁺) 179.2 [MH⁺]

Preparation 83***N*-(1-[2-[(*(3S)*-3-(3,4-Dichlorophenyl)-6-oxopiperidinyl]ethyl)-4-piperidinyl)-*N*-methylacetamide**

- 5 Triethylamine (4.2ml, 30mmol) was added to a suspension of [(*(3S)*-3-(3,4-dichlorophenyl)-6-oxopiperidinyl]acetaldehyde (WO 9605193) (5g, 17.5mmol) and the amine from preparation 60 (5g, 26.2mmol) in dichloromethane (400ml) and the mixture stirred at room temperature, until all solids had dissolved. Acetic acid, was added (ca. 5ml) to the reaction to give pH 4, the solution stirred
- 10 for 30 minutes, then sodium triacetoxyborohydride (7.4g, 35mmol) added, and the reaction stirred for 3 hours. The mixture was diluted with dichloromethane (300ml), and washed with sodium hydroxide solution (400ml, 1N). The aqueous phase was extracted with further dichloromethane (x2), the combined organic solutions washed with brine, dried (Na₂SO₄) and evaporated under reduced
- 15 pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol:0.88 ammonia (99:1:0.1 to 85:15:1.5) to give the title compound, 4.5g.
- ¹Hnmr (CD₃OD, 400MHz) δ: 1.35-2.17 (m, 15H), 2.17-2.41 (m, 3H), 2.75 (m, 1H), 2.78-2.93 (m, 4H), 3.40 (dd, 1H), 3.75 (dd, 1H), 3.56, 4.25 (m, 1H), 7.33
- 20 (m, 1H), 7.49 (d, 1H), 7.56 (s, 1H).
- LRMS : m/z (ES⁺) 448, 450 [MNa⁺]

Preparation 84

tert-Butyl 1-(1-(2-[(3*S*)-3-(3,4-dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinylethyl)-3-azetidinyl)-4-piperidinylcarbamate




- 5 A mixture of the aldehyde from preparation 11a (250mg, 0.62mmol), *tert*-butyl-1-(3-azetidinyl)-4-piperidinylcarbamate trifluoroacetate (WO 9605193) (450mg, 0.93mmol), triethylamine (1ml) and acetic acid (1.1ml) and sodium triacetoxyborohydride (250mg, 1.24mmol) in dichloromethane (50ml) was stirred at room temperature for 90 minutes. The reaction mixture was washed
- 10 with water, dried (MgSO₄) and evaporated under reduced pressure. The crude product was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol (100:0 to 95:5) to afford the title compound, 252mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.44 (s, 9H), 1.82-2.70 (m, 16H), 2.80-3.05 (m, 3H), 3.48 (m, 1H), 3.62 (m, 2H), 3.92 (d, 1H), 4.40 (d, 1H), 4.58 (d, 1H), 7.18 (m, 1H), 7.22 (d, 1H), 7.42 (s, 1H), 7.44 (d, 1H), 7.74 (d, 2H), 8.54 (d, 1H).

LRMS : m/z (TSP⁺) 502.1, 503.1 [MH⁺]

Preparation 85

tert-Butyl 1-(1-(2-[(3*S*)-3-(3,4-dichlorophenyl)-1-(6-methyl-2-pyridinyl)-6-oxopiperidinyl]ethyl)-3-azetidinyl)-4-piperidinylcarbamate




- 5 A mixture of the aldehyde from preparation 12a (260mg, 0.71mmol), *tert*-butyl-1-(3-azetidinyl)-4-piperidinylcarbamate trifluoroacetate (WO 9605193) (350mg, 0.78mmol), triethylamine (0.26ml, 1.86mmol) and titanium isopropoxide (2.3ml, 0.78mmol) in ethanol (3ml), was stirred at room temperature for 18 hours. Sodium borohydride (50mg, 1.35mmol) in ethanol (5ml) was then added and the reaction stirred for 30 minutes. Sodium hydroxide was added, the resulting precipitate filtered off, and washed with ethyl acetate. The filtrate was washed with water (2x) and brine (2x), dried (MgSO₄) and evaporated under reduced pressure. The residual gum was purified by column chromatography on silica gel using an elution gradient of dichloromethane:methanol (100:0 to 99:1) to afford the title compound as a yellow/white solid, 59mg.

¹Hnmr (CDCl₃, 400MHz) δ: 1.35 (m, 3H), 1.40 (s, 9H), 1.64-1.95 (m, 8H), 2.08 (m, 2H), 2.22 (m, 2H), 2.55 (m, 7H), 2.80 (t, 1H), 3.40 (m, 2H), 3.80 (d, 1H), 4.36 (m, 1H), 4.44 (d, 1H), 6.96 (d, 1H), 7.16 (d, 1H), 7.38 (dd, 2H), 7.50 (d, 1H), 7.58 (dd, 1H).

- 20 LRMS : m/z (TSP⁺) 617.2 [MH⁺]

CLAIMS:

1. A compound of formula (I):



- 5 or a pharmaceutically acceptable salt, prodrug, solvate or polymorph thereof, wherein:

R is het^a;

- 10 R¹ is phenyl optionally substituted by one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, and C₁₋₆ alkyl optionally substituted by one or more halogen;

m is 1-4;

15

Z is selected from:

- a) N(R³)(R⁴X)

wherein X is NR³R⁵, OR³, Oaryl¹, Ohet^b, Ohet^c, aryl¹, het^b or het^c;

- c) N(R³)Y

- 20 wherein Y is aryl¹, het^b or het^c; and

c) a 4-7 membered N containing saturated or partially saturated heterocycle said heterocycle attached to the alkylene link via said nitrogen atom, said heterocycle optionally containing an additional 1-3 groups, each independently selected from C=O, NH, S(O)_p and O; optionally, said heterocycle is:

- 25 (i) spirofused with het^b, such that both rings share 1 atom; or
 (ii) substituted by 1-3 groups each independently selected from het^b, het^c, aryl¹, R³, R⁴OR³, R⁴C(O)R³, OR³, OR⁷OR³, OR⁴OC(O)R³, OR⁴OC(O)NR³R⁶, S(O)_pR⁴, C(O)R³, C(O)NR³R⁶, C(O)OR³, R⁷C(O)OR³, C(O)R⁷OR³, C(O)OR⁷OR³, CF₃, NR³R⁶, R⁴NR³R⁵,
 30 OC(O)NR³R⁴ and NR³R⁵;

wherein R^3 and R^6 are both independently selected from H and C_{1-8} alkyl;
wherein R^4 and R^7 are both independently selected from C_{1-6} alkylene;
wherein R^5 is selected from $C(O)OR^3$, $S(O)_pR^3$, $S(O)_p\text{aryl}^1$, $C(O)R^3$, and
5 $C(O)NR^3R^6$;

het^b is a 4-7 membered heterocycle containing 1-3 heteroatoms, each independently selected from N, O and S, said N being optionally substituted with O, said ring optionally containing 1-2 C=O groups, said ring being saturated
10 or partially saturated, said ring being optionally benzofused, said ring being optionally substituted by 1-3 substituents selected from halo, R^3 , OR^3 , $C(O)NR^3R^6$, $R^7NR^3R^6$, NR^3R^5 , $NHS(O)_pR^4$, $S(O)_pNR^3R^6$, $S(O)_pR^4$, CN, NR^3R^6 and aryl^1 ;

15 het^a and het^c independently represent a 5-7 membered aromatic heterocycle containing 1-3 heteroatoms each independently selected from N, O and S, said ring being optionally benzofused, said ring system as a whole being optionally substituted by 1-3 substituents, each independently selected from: halo, R^3 , OR^3 , $C(O)NR^3R^6$, $R^4NR^3R^6$, NR^3R^5 , $NHS(O)_pR^4$, $S(O)_pNR^3R^6$, $S(O)_pR^4$, CN,
20 NR^3R^6 and $R^4NR^3S(O)_pR^3$;


aryl^1 is phenyl or naphthyl, each optionally substituted by 1-3 substituents, each independently selected from: halo, R^3 , OR^3 , $C(O)NR^3R^6$, $R^7NR^3R^6$, NR^3R^5 , $NHS(O)_pR^4$, $S(O)_pNR^3R^6$, $S(O)_pR^4$, CN;
25

p is 0, 1 or 2; and

n is 0-4.

30 2. A compound according to claim 1 having the following stereochemistry:

197



3. A compound according to claim 1 or claim 2, wherein R is pyridyl, optionally substituted by NR^3R^6 , R^3 or OR^3 .
- 5 4. A compound according to any of claims 1 to 3, wherein R^1 is phenyl optionally substituted by 1 or 2 halo substituents.
5. A compound according to any of claims 1 to 4, wherein m is 2-3.
- 10 6. A compound according to any of claims 1 to 5, wherein n is 1-4.
7. A compound according to any of claims 1 to 6, wherein R^3 is H or C_{1-4} alkyl.
- 15 8. A compound according to any of claims 1 to 7, wherein R^4 is C_{1-4} alkylene.
9. A compound according to any of claims 1 to 8, wherein R^5 is $\text{C}(\text{O})\text{OR}^3$, $\text{C}(\text{O})\text{R}^3$, $\text{C}(\text{O})\text{NR}^3\text{R}^6$.
- 20 10. A compound according to any of claims 1 to 9, wherein Z is a piperidine or azetidine group optionally substituted by one or more of het^b , het^c , aryl^1 , OR^3 , R^3 and NR^3R^5 , wherein;
 het^b is a 5-6 membered saturated or partially saturated nitrogen containing heterocycle, said heterocycle optionally incorporating 1-2 groups each
25 independently selected from O, C=O and N, said heterocycle being optionally benzofused, said heterocycle being optionally substituted by 1-2 substituents, each independently selected from OR^3 , R^3 , NR^3R^6 , NR^3R^5 , aryl^1 , SO_2R^4 and $\text{SO}_2\text{NR}^3\text{R}^6$;

het^c is pyridyl, optionally substituted by 1 or 2 substituents each independently selected from halo and OR³;


5 aryl¹ is phenyl, optionally substituted by 1 or 2 substituents each independently selected from halo and OR³; and

R³, R⁴, R⁵ and R⁶ are as defined in the previous claims.

11. A compound according to any of claims 1 to 10, wherein Z is a piperidine
10 or azetidine group, optionally substituted by het^b, aryl¹ and NR³R⁵; wherein het^b is a morpholine or piperidine, optionally substituted at the 4 position by OH and or methyl; wherein;
aryl¹ is phenyl optionally substituted by OH; and
R³ is H or methyl and R⁵ is C(O)CH₃

15

12. A compound according to any of claims 1 to 11, wherein Z is




13. A compound according to any of claims 1 to 10 selected from:
20 (5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 131)
(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-hydroxypiperidinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 135a)
(5S)-5-(3,4-Dichlorophenyl)-5-[2-(4-methoxy-1-piperidinyl)ethyl]-1-(2-pyridinyl)-
25 2-piperidinone (Example 61)
(5S)-5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{{2-[4-hydroxy-4-phenyl]-1-piperidinyl}ethyl}-2-piperidinone (Example 134)
(5S)-5-(3,4-Dichlorophenyl)-5-[2-[4-hydroxy-4-(2-pyridyl)-1-piperidinyl]ethyl]-1-(2-pyridinyl)-2-piperidinone (Example 92)
30 N-(1-{2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-phenyl-4-piperidinyl)-acetamide (Example 90)

199


- (5S)-5-(3,4-Dichlorophenyl)-1-(6-methoxy-2-pyridinyl)-5-{2-[3-(4-morpholinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 119)
- 5-(3,4-Dichlorophenyl)-1-(6-methyl-2-pyridinyl)-5-{2-[3-(4-oxo-1-piperidinyl)-1-azetidiny]ethyl}-2-piperidinone (Example 168)
- 5 (5S)-5-(3,4-Dichlorophenyl)-5-{2-[3-(4-hydroxy-1-piperidinyl)-1-azetidiny]ethyl}-1-(2-pyridinyl)-2-piperidinone (Example 73)
- N*-(1-{2-[(3S)-3-(3,4-Dichlorophenyl)-6-oxo-1-(2-pyridinyl)piperidinyl]ethyl}-4-piperidinyl)-*N*-methylacetamide (Example 158)

- 10 14. A process of preparing a compound according to any of claims 1 to 13 comprising subjecting a compound of formula (II) to a reductive amination to give a compound of formula (I):



- 15 wherein R, R¹, m, n and Z are as defined in the previous claims.

15. A process for preparing a compound according to any of claims 1 to 13, comprising subjecting a compound (VIII) to an alkylation reaction to give a compound of formula (I):



wherein R, R¹, m, n and Z are as defined in the previous claims.

16. Use of a compound according to any of claims 1 to 13, as a medicament.
- 5
17. Use of a compound according to any of claims 1 to 13, in the preparation of a medicament for the treatment of a condition selected from: inflammatory disease, a central nervous system (CNS) disorder, a gastro-intestinal (GI) disorder, a disease caused by *Helicobacter pylori* or other urease positive Gram negative bacteria, urological conditions, a pulmonary disorder, an allergy, a hypersensitivity disorder, a vasospastic disease, a proliferative disorder, a fibrosing or collagen disease, reflux sympathetic dystrophy, an addiction disorder, a stress-related somatic disorder, a peripheral neuropathy, a neuropathological disorder, a disorder related to immune enhancement or suppression, a rheumatic disease, an opthalmic disease, acute and chronic pain or a viral disease.
- 10
18. Use of a compound according to any of claims 1 to 13 in the preparation of a medicament for the treatment of a condition selected from: urological conditions or acute and chronic pain.
- 15
19. A use according to claim 18, wherein said urological condition is overactive bladder.
- 20

20. A use according to claim 18, wherein said pain is neuropathic pain.
21. A method of treating or preventing a condition for which an NK₂ antagonist is efficacious which comprises administering a therapeutically effective amount
5 of a compound according to any of claims 1 to 13 to a patient in need of treatment.
22. Use of a compound according to any of claims 1 to 13 in the preparation of a medicament in combination with an agent selected from: muscarinic
10 antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); noradrenalin reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; PDE₅ inhibitors; COX₂ inhibitors; non-selective COX inhibitors; vanilloid receptor agonists; HMG-CoA reductase inhibitors; and estrogenic modulators and selective estrogen receptor modulators for the
15 treatment of urological conditions.
23. Use of a compound according to any of claims 1 to 13 in the preparation of a medicament in combination with an agent selected from: NSAIDs, opioids, muscarinic antagonists; cholinergic analgesics; alpha-adrenoceptor antagonists;
20 serotonin/noradrenalin reuptake inhibitors (SNRI); COX₂ inhibitors; non-selective COX inhibitors; tricyclic antidepressants, anticonvulsants, serotonin reuptake inhibitors, serotonin receptor agonists and antagonists, sedatives, skeletal muscle relaxant and NMDA receptor antagonists for the treatment of pain.
25
24. A composition comprising a compound according to any of claims 1 to 13 and a pharmaceutically acceptable diluent or carrier.
25. A composition comprising a compound according to any of claims 1 to 13
30 and an agent selected from: Muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; PDE₅ inhibitors; COX₂ inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); vanilloid

receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier.

- 5 26. A composition comprising a compound according to any of claims 1 to 13 and an agent selected from: NSAIDs, opioids, muscarinic antagonists; cholinergic analgesics; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); COX₂ inhibitors; non-selective COX inhibitors; tricyclic antidepressants, anticonvulsants, serotonin reuptake inhibitors, 10 serotonin receptor agonists and antagonists, sedatives, skeletal muscle relaxant and NMDA receptor antagonists, and a pharmaceutically acceptable diluent or carrier.
27. A kit comprising
- 15 a) a composition comprising a compound according to any of claims 1 to 13 and a pharmaceutically acceptable diluent or carrier;
- b) a composition comprising an agent selected from: compounds which modulate the action of atrial natriuretic factor (also known as atrial natriuretic peptide), such as inhibitors of neutral endopeptidase; compounds which inhibit 20 angiotensin-converting enzyme, and combined inhibitors of angiotensin-converting enzyme and neutral endopeptidase; angiotensin receptor antagonists; substrates for NO-synthase; calcium-channel blockers; antagonists of endothelin receptors and inhibitors of endothelin-converting enzyme; cholesterol lowering agents; antiplatelet and antithrombotic agents, 25 thromboplastin activating factor inhibitors; insulin sensitising agents and hypoglycaemic agents; acetylcholinesterase inhibitors; non-steroidal anti-inflammatory agents (NSAIDs); cGMP PDE₅ inhibitors; muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); noradrenalin reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} 30 agonists/antagonists; COX₂ inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); opioids; tricyclic antidepressants; anticonvulsants, serotonin reuptake inhibitors; serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, sedatives, skeletal muscle

relaxants; NMDA receptor antagonists; vanilloid receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier; and

c) a container.

5

28. A kit comprising:

a) a composition comprising a compound according to any of claims 1 to 13 and a pharmaceutically acceptable diluent or carrier; and

b) a composition comprising a compound of formula (I) or a pharmaceutically acceptable salt, solvate or prodrug thereof and an agent selected from: muscarinic antagonists; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); reuptake inhibitors; NK₁ antagonists; 5-HT_{1A} agonists/antagonists; PDE₅ inhibitors; COX₂ inhibitors; non-selective COX inhibitors (preferably with 'GI protection'); vanilloid receptor agonists; HMG-CoA reductase inhibitors; estrogenic modulators and selective estrogen receptor modulators, and a pharmaceutically acceptable diluent or carrier; and

c) a container.

29. A kit comprising:

a) a composition comprising a compound according to any of claims 1 to 13 and a pharmaceutically acceptable diluent or carrier; and

b) a composition comprising a compound according to any of claims 1 to 13 and an agent selected from: NSAIDs, opioids, muscarinic antagonists; cholinergic analgesics; alpha-adrenoceptor antagonists; serotonin/noradrenalin reuptake inhibitors (SNRI); COX₂ inhibitors; non-selective COX inhibitors; tricyclic antidepressants, anticonvulsants, serotonin reuptake inhibitors, serotonin receptor agonists and antagonists, sedatives, skeletal muscle relaxant and NMDA receptor antagonists, and a pharmaceutically acceptable diluent or carrier; and

c) a container.

INTERNATIONAL SEARCH REPORT

Original Application No.

PCT/IB 02/05234

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/14 C07D401/04 C07D491/10 C07D405/14 C07D413/14
 A61K31/4545 A61P13/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 968 923 A (MARCHINGTON ALLAN PATRICK ET AL) 19 October 1999 (1999-10-19) cited in the application claim 1	1-29
A	US 6 242 438 B1 (MARCHINGTON ALLAN PATRICK ET AL) 5 June 2001 (2001-06-05) cited in the application claim 1	1, 14, 16-18, 21-29

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

1 April 2003

Date of mailing of the international search report

10/04/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3018

Authorized officer

Diederens, J

INTERNATIONAL SEARCH REPORT

national application No.
PCT/IB 02/05234

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 21 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 02/05234

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5968923	A	19-10-1999	AT 200083 T	15-04-2001
			AU 689303 B2	26-03-1998
			AU 3254995 A	07-03-1996
			DE 69520502 D1	03-05-2001
			DE 69520502 T2	12-07-2001
			DK 775132 T3	30-04-2001
			EP 0775132 A1	28-05-1997
			FI 970523 A	07-02-1997
			GR 3035726 T3	31-07-2001
			JP 3159389 B2	23-04-2001
			JP 9508646 T	02-09-1997
			KR 218626 B1	01-09-1999
			NO 970566 A	07-02-1997
			NZ 291423 A	25-03-1998
			PL 318534 A1	23-06-1997
			BR 9503582 A	30-04-1996
			CA 2197086 A1	22-02-1996
			CN 1154699 A , B	16-07-1997
			CZ 9700381 A3	14-01-1998
			WO 9605193 A1	22-02-1996
			ES 2155894 T3	01-06-2001
			HU 77771 A2	28-08-1998
			IL 114826 A	31-12-1999
			PT 775132 T	31-07-2001
			RU 2150468 C1	10-06-2000
			TR 960127 A2	21-06-1996
			TW 432061 B	01-05-2001
US 6242438	B1	05-06-2001	AT 232526 T	15-02-2003
			AU 708282 B2	29-07-1999
			AU 1195097 A	01-08-1997
			BG 102589 A	30-09-1999
			BR 9612412 A	13-07-1999
			DE 69626220 D1	20-03-2003
			EP 0871623 A1	21-10-1998
			JP 11501667 T	09-02-1999
			JP 3123611 B2	15-01-2001
			KR 275402 B1	15-12-2000
			NO 982651 A	09-06-1998
			NZ 324712 A	28-05-1999
			PL 327665 A1	21-12-1998
			SK 89598 A3	14-02-2000
			AP 709 A	22-12-1998
			CZ 9802093 A3	14-04-1999
			WO 9725322 A1	17-07-1997
			HR 970006 A1	30-06-1998
			HU 9903590 A2	28-05-2000
			JP 3254205 B2	04-02-2002
			JP 2000344741 A	12-12-2000
			RU 2158264 C2	27-10-2000
			TR 9801268 T2	21-10-1998
			TW 472054 B	11-01-2002
			ZA 9700047 A	03-07-1998